



# TELDOR®

Safety Information

**TELDOR**

UFI: K6R1-J09H-W009-79S3

Contains 50% w/w fenhexamid



**Toxic to aquatic life with long lasting effects.**

Wear protective gloves/protective clothing/eye protection.

Dispose of contents/container to a licenced hazardous-waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**Contains 4-amino-2,3-dichlorophenol. May cause an allergic reaction.**

**To avoid risks to human health and the environment, comply with the instructions for use**

PCS No. 02003

IE80487169f rA3g

**PROTECT FROM FROST**

**For 24 hour Emergency Information contact Bayer CropScience Limited**

**Freephone: 00800 1020 3333**

Authorisation holder: Bayer CropScience Limited  
230 Cambridge Science Park, Milton Road, Cambridge, CB4 0WB, United Kingdom  
Marketing Company: Bayer CropScience Ltd, Bayer Ltd, 1st Floor, The Grange Offices  
The Grange, Brewery Road, Stillorgan, Co. Dublin A94 H2K7 Freephone: 1800 818534

**Bayer**



# TELDOR®

10 x 1Kg e

**GROUP 17 FUNGICIDE**

For professional use only

For use only as a horticultural fungicide for the control of *Botrytis* (grey mould) on strawberries, raspberries, loganberries, blackberries and other *Rubus* hybrids, blackcurrants, redcurrants, whitecurrants and gooseberries.

Water dispersible granule formulation containing 50% w/w fenhexamid



Specimen-  
2022-10-10 date

4085\_D



# TELDOR®

# TELDOR®

# TELDOR®

# TELDOR®

### SAFETY PRECAUTIONS

#### Operator Protection

Wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.  
Wash all protective clothing after use.  
Wash concentrate from skin or eyes immediately.  
Do not breathe spray.

Wash hands and exposed skin before meals and after work.

If swallowed seek medical advice immediately and show this label or container.

#### Environmental Protection

Do not contaminate water with the product or its container.  
(Do not clean application equipment near surface water / avoid contamination via drains from farmyards and roads).

#### Storage and Disposal

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. Keep in original container tightly closed in a safe place. Keep out of reach of children.

PROTECT FROM FROST  
READ ALL INSTRUCTIONS CAREFULLY BEFORE USE  
FOR USE ONLY AS A HORTICULTURAL FUNGICIDE



To access the **Safety Data Sheet** for this product scan the code or use the link below:  
[www.bayercropscience.ie/sds/teldor.pdf](http://www.bayercropscience.ie/sds/teldor.pdf)  
or alternatively contact your supplier

IEB4060275d rA3g

Bayer

#### DIRECTIONS FOR USE

For detailed instructions, including rate of use, see enclosed instruction leaflet.

© Bayer CropScience Limited 2022

1Kg e

GROUP 17 FUNGICIDE

For professional use only.

For use only as a horticultural fungicide for the control of *Botrytis* (grey mould) on strawberries, raspberries, loganberries, blackberries and other *Rubus* hybrids, blackcurrants, redcurrants, whitecurrants and gooseberries.

A water dispersible granule formulation containing 50% w/w fenhexamid

Authorisation holder:  
Bayer CropScience Ltd.  
230 Cambridge Science Park  
Milton Road  
Cambridge  
CB4 0WB  
United Kingdom

Freephone: 1800 818534

For 24 hour emergency information contact  
Bayer CropScience Ltd Freephone: 00800 1020 3333

Marketing company:  
Bayer CropScience Ltd  
Bayer Ltd, 1st Floor,  
The Grange Offices,  
The Grange,  
Brewery Road,  
Stillorgan, Co. Dublin  
A94 H2K7



**Toxic to aquatic life with long lasting effects.**

Wear protective gloves/protective clothing/eye protection.  
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains 4-amino-2,3-dichlorophenol. May cause an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 02003



**GROUP 17 FUNGICIDE**

For professional use only.

For use only as a horticultural fungicide for the control of *Botrytis* (grey mould) on strawberries, raspberries, loganberries, blackberries and other *Rubus* hybrids, blackcurrants, redcurrants, whitecurrants and gooseberries.

A water dispersible granule formulation containing 50% w/w fenhexamid

**Authorisation holder:**  
Bayer CropScience Ltd.  
230 Cambridge Science Park  
Milton Road  
Cambridge  
CB4 0WB  
United Kingdom

**Marketing company:**  
Bayer CropScience Ltd  
Bayer Ltd, 1st Floor,  
The Grange Offices,  
The Grange,  
Brewery Road,  
Stillorgan, Co. Dublin  
A94 H2K7

Freephone: 1800 818534

**For 24 hour emergency information contact  
Bayer CropScience Ltd**

Freephone: 00800 1020 3333

Risk and Safety Information:

**TELDOR**

UFI: K6R1-J09H-W009-79S3

Contains 50% w/w fenhexamid



**Toxic to aquatic life with long lasting effects.**

Wear protective gloves/protective clothing/eye protection.

Dispose of contents/container to a licenced hazardous-waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**Contains 4-amino-2,3-dichlorophenol. May cause an allergic reaction.**

**To avoid risks to human health and the environment, comply with the instructions for use**

**PCS No. 02003**

**SAFETY PRECAUTIONS**

**Operator Protection**

Wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.

Wash all protective clothing after use.

Wash concentrate from skin or eyes immediately.

Do not breathe spray.

Wash hands and exposed skin before meals and after work.

If swallowed seek medical advice immediately and show this label or container.

**Environmental Protection**

Do not contaminate water with the product or its container (Do not clean application equipment near surface water / avoid contamination via drains from farmyards and roads).

**Storage and Disposal**

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. Keep in original container tightly closed in a safe place.

Keep out of reach of children.

PROTECT FROM FROST

**READ ALL INSTRUCTIONS CAREFULLY BEFORE USE.  
USE ONLY AS A HORTICULTURAL FUNGICIDE.**

**DIRECTIONS FOR USE**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Teldor controls *Botrytis cinerea* (grey mould) on strawberries, cane fruit and bush fruit.

**RATE OF USE**

CROP	Maximum individual dose	Maximum number of applications	Latest time of application
Strawberries	1.5 kg/ha	4	1 day before harvest
Raspberries, blackberries, loganberries and other <i>Rubus</i> hybrid berries	1.5 kg/ha	4	7 days before harvest
Currants, (black, red or white)	1.5 kg/ha	4	3 days before harvest
Gooseberries	1.5 kg/ha	4	7 days before harvest

For successful control of *Botrytis* in all listed crops it is vital to achieve complete cover on all flowers and fruitlets throughout the blossom period. Sprayers should be set up in such a way as to achieve maximum cover on these parts, with sprays directed over the crop rows (not applied overall).

Do not use finer than 80 mesh filters.

Apply Teldor as a **MEDIUM** quality (BCPC definition) spray.

**DISEASES CONTROLLED**

For control of *Botrytis cinerea* (grey mould)

Alternating or mixing fungicides with different modes of action is a recognised disease resistance management strategy.

To minimise the likelihood of the development of *Botrytis* strains resistant to Teldor no more than 2 sprays of Teldor alone should be applied consecutively within a programme of sprays. Afterwards change to botryticides from different chemical groups for at least two consecutive sprays. See also 'Maximum number of Applications' above.

The possible development of disease strains resistant to Teldor cannot be excluded or predicted. Where such resistant strains occur, Teldor is unlikely to give satisfactory control.

**CROP SPECIFIC INFORMATION**

**Strawberries**

The first spray of a fungicide programme should be made at first flower (white bud).

Apply 1.5 kg Teldor in at least 300 litres of water per hectare, with a 10-14 day interval between sprays – see 'Maximum number of Applications' above.

**Raspberries, Blackberries, Loganberries and other *Rubus* hybrid berries**

The first spray of a fungicide programme should be made at early flower (white bud).

Apply 1.5 kg Teldor in at least 1,000 litres of water per hectare, with a 7 day interval between sprays – see 'Maximum number of Applications' above

**Currants, (black, red or white) and Gooseberries**

The first spray of fungicide programme should be made at the late grape stage for currants and at first open flowers for gooseberries.


Apply 1.5 kg Teldor in at least 1,000 litres of water per hectare, with a 10-14 day interval between sprays – see 'Maximum number of Applications' above.

**MIXING**

Half-fill the spray tank with clean water. Commence agitation. Steadily add the recommended quantity of Teldor directly into the water in the spray tank. Complete filling and maintain agitation of the suspension before and during spraying until the tank is empty.

® Teldor is a Registered Trade Mark of Bayer

© Bayer CropScience Limited 2022



To access the **Safety Data Sheet** for this product scan the code or use the link below:  
[www.bayercropscience.ie/sds/teldor.pdf](http://www.bayercropscience.ie/sds/teldor.pdf)  
or alternatively contact your supplier

IE84060275d rA3g  
Bayer CropScience Ltd  
Bayer Ltd, 1st Floor,  
The Grange Offices,  
The Grange,  
Brewery Road,  
Stillorgan, Co. Dublin  
A94 H2K7

**Bayer**