

FRUPICA® SC

MAPP 12067/PCS 02128

GROUP 9 FUNGICIDE


For the control of *Botrytis* in outdoor and protected strawberries.

A suspension concentrate formulation containing 440 g/L (40% w/w) mepaniprym.

Risk and Safety Information

FRUPICA SC

A suspension concentrate formulation containing 440 g/L (40% w/w) mepaniprym.



UFI NI: NM99-Y074-K00Y-YJKP UFI IE: SCW9-90PP-C00V-6J2F

WARNING

SUSPECTED OF CAUSING CANCER. VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS.

Obtain special instructions before use.

Do not handle until all safety precautions have been read and understood.

Avoid release to the environment.

Wear protective gloves/protective clothing.

IF ON SKIN: Wash with plenty of soap and water.

IF exposed or concerned: Get medical advice/attention.

Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

Contains 1,2 benzisothiazolin-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

MAPP 12067/PCS 02128

IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Crops	Maximum individual dose	Maximum number of treatments	Latest time of application	Other specific restrictions
See full details in leaflet attached to the pack				

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

FUNGICIDE

F

Net content

1L

READ CAREFULLY BEFORE USE

PROTECT FROM FROST SHAKE WELL BEFORE USE

Production date / Batch number: see packaging

For advice on medical emergencies, fires, spillages or chemical hazards ONLY - phone Carechem 24 EU: +44 (0)1235 239670

For technical and non-emergency calls phone +44 (0)845 373 0305

Authorisation Holder:
Certis Belchim B.V.

Stadsplateau 16, 3521 AZ Utrecht
PO Box 607, 3500 Ap Utrecht
Netherlands

®Registered trademark of
Kumiai Chemical Industry Co. Ltd

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK – UK only.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (IMPERMEABLE COVERALLS) AND SUITABLE PROTECTIVE GLOVES when using handheld equipment. However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH HANDS before meals and after work. Wash all protective clothing after use.

Environmental protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from handheld sprayers to fall within 1 m of the top of a bank of a static or flowing water body. Aim spray away from water – UK only.

Storage and disposal

EMPTY CONTAINER completely and dispose of safely. DO NOT RE-USE CONTAINER FOR ANY PURPOSE. Rinse empty container thoroughly by using an integrated pressure rinsing device, or by manually rinsing five times. Add washings to the spray tank at time of filling.



IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Crops	Maximum individual dose	Maximum number of treatments	Latest time of application
Strawberry	0.9 litres/ha	2 (per crop)	3 days before harvest

Other specific restrictions:

- 1) This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years (UK only).
- 2) Non-returnable containers must not be re-used for any purpose

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Crops

FRUPICA SC can be used on outdoor and protected strawberries.

Mixing

Shake well before use. Fill the tank with half the required volume of water. Measure the appropriate amount of FRUPICA SC into the tank and make up to the required volume of water. Agitate thoroughly before and during application.

Application

FRUPICA SC should be applied as a preventative spray at a rate of 0.68 L/ha under conditions where *Botrytis* could develop. This rate can be increased to 0.9 L/ha under conditions of severe disease pressure. The product should be applied in 500 to 1000 litres of water/ha to ensure complete coverage of the strawberry plant.

Method of application

Tractor mounted sprayer/ knapsack/ rotary atomiser.

Timing

The first application should be made from the start of flowering, with a second application 7–10 days later or as required in a programme of treatments. A maximum of two applications should be made per crop. To maintain control of *Botrytis cinerea*, FRUPICA SC must be used as part of a programme with other fungicides that provide effective control of grey mould. (See also Other information).

Rate of use

Maximum individual dose: 0.9 L/ha

Tank washing procedure

Immediately after application of FRUPICA SC, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with water and a commercially available cleaning agent. Rinse inside of tank with clean water and cleaning agent and flush through boom and hoses using at least one tenth of the spray tank volume. Drain completely. Do not contaminate surface waters or ditches with chemical or used container.

Dispose of spray washings in accordance with the Code of Practice for the Safe Use of Pesticides on Farms and Holdings (DEFRA).

Storage

Store above 5°C.
Protect from frost.

Other information

Resistance of *Botrytis* species can occur to groups of chemicals if they are over used. There is currently no known resistance to FRUPICA SC. However, as a precaution, it is advisable to use other products from different chemical groups as part of a mixed spray programme. Do not use more than two applications per crop of FRUPICA SC or other anilinopyrimidine containing products, such as pyrimethanil (e.g. Scala). FRUPICA SC is an anilinopyrimidine fungicide.

Do not allow direct spray from crop sprayers to fall within 5 m of the top of a static or flowing water body or within 1 m of the top of a ditch that is dry at the time of application. Consult processor before using on crops for processing.

PROTECT FROM FROST

This section is not part of the Product Label under the Control of Pesticides Regulations 1986 or the Plant Protection Products Regulations 1995 (as appropriate) and that it provides additional advice on product use at the discretion of the applicant.

Company Advisory Information

CONDITIONS OF SUPPLY

The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade or otherwise and whether as to condition, quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort, for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumers statutory rights are not affected.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:

WEBLINK to specific data sheet on manufacturer's website:



https://certisbelchim.co.uk/pdf/MSDS/Frupica_SC_MSDS.pdf

Alternatively, contact your supplier.

Specimen-
2023 to date