



Swift[®] SC

1L e

GROUP 11 FUNGICIDE

For use only as an agricultural fungicide for the control of foliar diseases in winter and spring wheat and winter and spring barley.

A suspension concentrate formulation containing 500 g/L (45.1% w/w) trifloxystrobin.

For Professional use only



Authorisation holder
Bayer CropScience Ltd.
230 Cambridge Science Park
Milton Road, Cambridge, CB4 0WB

Marketing company
Bayer CropScience Ltd
Bayer Ltd, 1st Floor,
The Grange Offices, The Grange,
Brewery Road, Stillorgan, Co. Dublin
A94 H2K7

For 24 hour emergency information contact Bayer CropScience Ltd
Telephone: 00800 0120 3333

F

Safety Information

SWIFT SC

UFI: EUA0-X0R2-700C-5WGH
Contains 500 g/L (45.09 % w/w) trifloxystrobin



Warning

Very toxic to aquatic life with long lasting effects.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains trifloxystrobin, 1,2-Benzisothiazolin-3-one and 5-Chloro-2-methyl-isothiazol-3-one/2-Methyl-isothiazol-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use

PCS No. 02145

SAFETY PRECAUTIONS

Operator Protection

Avoid contact with skin and eyes

Wear suitable protective clothing and gloves

In case of contact with eyes, rinse immediately with plenty of water and seek medical advice.

If swallowed, seek medical advice immediately and show this container or label.

Environmental Protection

Do not contaminate ponds, waterways or ditches with chemical or used container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

Do not allow direct spray from ground crop sprayers to fall within 5 metres of the top of the bank of a static or flowing waterbody or within 1 metre of the top of a ditch that is dry

at the time of application. Direct spray away from water.

Storage and Disposal

Keep out of reach of children.

Do not re-use container for any other purpose and dispose of safely.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. Triple rinsed containers should be punctured to prevent re-use and may be disposed of by an authorised contractor or at a municipal waste recycling site

PROTECT FROM FROST

STORE IN A COOL DRY PLACE

READ ALL INSTRUCTIONS CAREFULLY BEFORE USE



swiftscsds

To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.bayercropscience.ie/sds/swiftsc.pdf

or alternatively contact your supplier

IE87323528b rA5d

Bayer

DIRECTIONS FOR USE

Swift SC is a suspension concentrate containing the strobilurin fungicide trifloxystrobin. It is recommended for control of a range of foliar diseases on winter and spring wheat and winter and spring barley. Swift SC may be used on all commercial varieties of winter and spring wheat and barley.

Treatment at early stages of disease development will ensure best results. Apply at first signs of disease. A second application may be needed if disease attack is prolonged.

RATE OF USE

| Crop | Maximum individual dose: | Maximum number of applications | Latest time of application |
|--------------------------|--------------------------------|--------------------------------|---|
| Winter and spring wheat | 0.5 litres product per hectare | 2 per season | At any stage before caryopsis watery ripe stage (GS 71) |
| Winter and spring barley | 0.5 litres product per hectare | 2 per season | At any stage before caryopsis watery ripe stage (GS 71) |

Method of application: Tractor mounted sprayer

A spray pressure of 2-3 bar is recommended. Apply Swift SC in 200-300 litres per hectare water.

Apply as a medium spray quality.

DISEASES CONTROLLED

When applied as recommended Swift SC controls the following diseases when applied preventatively before disease is established in the crop:

Winter and Spring Wheat

Moderate control of Brown rust

Winter & Spring Barley

Rhynchosporium

Net blotch

Moderate control of Brown rust

Appropriate disease risk assessment or decision support systems must be used when deciding whether to treat with Swift SC.

Management Strategy to Minimise the Likelihood of Resistance

Trifloxystrobin is a strobilurin fungicide.

Apply STAR (Strobilurin Type Action and Resistance) fungicides according to manufacturer's recommendations for the target disease at the specific crop growth stage indicated. Effective disease management is a critical parameter in delaying the build-up of resistance pathogen populations.

Apply a maximum of two STAR containing sprays per cereal crop. Limiting the number of sprays is an important factor in delaying the build-up of resistant pathogen populations.

Swift SC must always be used in mixture with another product, containing an active ingredient from a different cross resistance group, that is recommended for control of the same target disease applied at a rate that will give robust control.

TO AVOID BUILD-UP OF RESISTANCE DO NOT APPLY THIS OR ANY OTHER PRODUCT CONTAINING STROBILURINS MORE THAN TWO TIMES PER CROP PER YEAR.

MIXING AND APPLICATION

Make sure the sprayer is set to give an even application at the correct volume. Fill the spray tank with half the volume of clean water and start agitation. Add required amount of Swift SC and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying.

COMPATIBILITY

Swift SC may be applied as a tank-mix with a range of products. Contact Bayer Crop Science for compatibility information on specific tank-mixes. Full manufacturer's instructions must be followed for each tank-mix component.