

RHINO[®]

MAPP 14311/ PCS 04400

GROUP FRAC 7 FUNGICIDE

For use as a potato seed treatment for the control of black scurf and stem canker.

A soluble concentrate containing 460 g/L (40.7% w/w) flutolanil.

RHINO

A soluble concentrate containing 460 g/L (40.7% w/w) flutolanil.



WARNING

CAUSES SERIOUS EYE IRRITATION
TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS

UFI: ANVO-30SW-H007-UVJX
Avoid breathing mist/spray.
IF ON SKIN: Wash with plenty of soap and water.
IF IN EYES: Rinse continuously with water for several minutes.
Remove contact lenses, if present and easy to do continue rinsing.
If eye irritation persists: Get medical advice/attention.
Wash contaminated clothing before reuse.
Avoid release to the environment.
Collect spillage.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

Allergic risks to human health and the environment, comply with the instructions for use.

MAPP 14311/ PCS 04400

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL SEED TREATMENT

Crops	Maximum individual dose	Maximum number of treatments	Latest time of application	Other specific restrictions
See DIRECTIONS FOR USE within the leaflet attached to the pack				

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

FUNGICIDE

Net content

1L



READ CAREFULLY BEFORE USE

PROTECT FROM FROST

Production date / Batch number:
see packaging

For advice on medical emergencies, fires, spillages or chemical hazards ONLY - phone +44 (0)870 190 6777

For technical and non-emergency calls
phone +44 (0)845 373 0305

Authorisation Holder:
Nichino Europe Co. Limited
5 Pioneer Court, Vision Park
Histon, Cambridge CB24 9PT

® Registered trademark of
Certis Belchim B.V.

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK – UK only.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces, freshly treated material or applying the product.
WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.
However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection – UK only.
When using, do not eat, drink or smoke.
Causes serious eye irritation.
IF IN EYES: Rinse continuously with water for several minutes. Remove contact lenses, if present and easy to do continue rinsing.
If eye irritation persists: Get medical advice/attention.

Environmental protection

Do not contaminate water with the product or its container.
Do not clean application equipment near surface water.
Avoid contamination via drains from farmyards and roads.

Storage and disposal

Keep out of reach of children.
Keep away from food, drink and animal feeding stuffs.
KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
WASH OUT CONTAINER THOROUGHLY, and dispose of safely.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL SEED TREATMENT

Crops:	Seed potatoes
Maximum individual dose:	200 ml of product per tonne of potatoes
Maximum number of treatments:	One per batch
Latest time of application:	At planting

Other specific restriction:

This product must only be applied by conventional canopied, hydraulic or spinning disc equipment (with or without electrostatics) mounted on a rolling conveyor or table.

Treated seed must not be used for food or feed.

Sacks containing treated seed must not be re-used for food or feed.

The product must be diluted with a minimum of 0.2 L water, up to a maximum of 2.0 L water per tonne of tubers to ensure good coverage of the seeds.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS/WARNINGS

- RHINO acts against the tuber borne inoculum of *Rhizoctonia solani* which independent sources have indicated is the most important source of infection. However, where the soil borne inoculum level is high, treatment of tubers with RHINO may not give full protection against infection of stems and daughter tubers.
- For best results the tubers should be clean and free from soil (less than 5% of surface). Disease under areas of soil/dust will not be exposed to spray and will not be controlled.
- Seed tubers should be of good quality and vigour (free from bacterial rots, physical damage, virus infection etc), and should not be sprouted to such an extent that mechanical damage to the shoots will occur during treatment or planting.
- Check with your processor before using RHINO.

DISEASE CONTROL

RHINO is a seed potato treatment which controls stem canker and black scurf (*Rhizoctonia solani*) when applied to the seed prior to planting. It controls black scurf on seed tubers and protects the progeny against black scurf and stem canker.

CROPS

RHINO can be used on seed potatoes for the seed, ware or processing market. It can be used on the seed tubers of all varieties of early, second early and main crop potatoes grown for seed, ware or processing.

TIMING

RHINO should be applied to the seed tubers before chitting, prior to planting or at planting. Only apply once per batch.

APPLICATION

Apply 200 ml of RHINO per tonne of seed potatoes. It must be diluted with a minimum of 0.2 L water, up to a maximum of 2.0 L water per tonne of tubers to ensure good coverage of the seeds. Apply RHINO only by canopied, hydraulic or spinning disc equipment (with or without electrostatics) mounted on a rolling conveyor or table.

MIXING

Fill the tank with half the required volume of water. Measure the appropriate amount of RHINO into the tank and make up to the required volume of water. Agitate thoroughly before and during application.

Storage

Store above 5°C

Protect from frost

Company Advisory Information

This section is not part of the Product Label under Regulation 1107/2009. It provides additional advice on product use at the discretion of the applicant.

CONDITIONS OF SUPPLY

The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade or otherwise and whether as to condition, quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort, for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumers statutory rights are not affected.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:

WEBLINK to specific data sheet on manufacturer's website:



https://certisbelchim.co.uk/pdf/MSDS/Rhino_MSDS.pdf

Alternatively, contact your supplier.