

DIMIX 500 SC

FUNGICIDE

Contains 500 g/L dimethomorph as a Suspension Concentrate (SC)
Fungicide for the control of foliar late blight (*Phytophthora infestans*) in potato

Risk & Safety Information

Contains 1,2-benzisothiazol-3(2H)-one.
May produce an allergic reaction

Warning

Harmful if swallowed
Very toxic to aquatic life with long lasting effects.

Keep out of reach of children.
Wash thoroughly after handling.
Do not eat, drink or smoke when using this product.
IF SWALLOWED: Call a POISON Centre or doctor/
physician if you feel unwell.
Rinse mouth.
Dispose of contents/container to a licensed
hazardous-waste disposal contractor or
collection site except for triple rinsed
empty containers which can be disposed
of as non-hazardous waste.

To avoid risks to human health and the
environment, comply with the instructions for use.

PCS no. 05191



Marketing Company and
Approval holder:
Arysta LifeScience Benelux sprl
Rue de Renory 26/1
B-4102, Ougrée
Belgium
Tel: +32 4 385 97 11

24-Hour emergency telephone
number: +44 (0)1235 239670

Manufacturing date and Batch No.:
See packaging

Important Information

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE.

<u>Crops</u>	<u>Max. single dose</u> <u>(L product/ha)</u>	<u>Max no. of</u> <u>treatments</u> <u>(per crop)</u>	<u>Max. total dose</u> <u>(L product/ha)</u>	<u>Latest time of</u> <u>application</u>
Seed and ware potatoes (early and maincrop)	0.36	5	1.8	7 days before harvest

Other specific restrictions:

Dimix 500 SC must always be applied in a tank mixture with an approved mancozeb formulation at an equivalent dose rate of a minimum of 1200g mancozeb/ha.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

Operator protection

WASH SPLASHES from skin or eyes immediately.
WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental Protection

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies. Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

DISEASES CONTROLLED

Dimix 500 SC is a systemic fungicide containing 500 g/L dimethomorph. It should be used in tank mixtures with an approved mancozeb formulation for the preventative control of potato blight (*Phytophthora infestans*) on potato.

CROP-SPECIFIC INFORMATION

Dose Rate

Apply Dimix 500 SC at 0.36 L/ha (equivalent to 180 g dimethomorph/ha) in tank mixture with an approved mancozeb formulation at an equivalent dose rate of 1200 g mancozeb/ha when infection is high and curative action is a priority. Alternatively, where there is low infection, apply Dimix 500 SC at 0.3 L/ha (equivalent to 150 g dimethomorph/ha) with an approved mancozeb formulation at equivalent 1340 g mancozeb/ha.

Timing

The first spray should be applied as soon as there is a risk of blight infection or when a disease warning is given. Subsequent applications should be made every 7 days (up to 5 applications in total).

APPLICATION

To be applied via tractor mounted/trailed sprayer only.
Add the required quantity of Dimix 500 SC to a half-filled spray tank with continuous agitation, then add the remaining volume of water. Spray immediately after mixing and continue agitation during spraying. Rinse all equipment well immediately after use. Follow any specific washing instructions on the label of products applied in mixture with Dimix 500 SC.

Tank-Mix recommendations: Dimix 500 SC must always be used in tank mixture with an approved mancozeb formulation. Apply using standard field sprayers as a MEDIUM spray. Always follow the instructions given on the label of the tank-mix partner.

RESISTANCE

Any fungicide resistance advice or restrictions on the label of the partner product should be complied with.

CONDITIONS OF SALE

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

IE-DMX

5L

 Arysta
LifeScience