



TOWER®



ADAMA



HERBICIDES

Suspension concentrate

PCS 05425

FOR PROFESSIONAL USE ONLY



Ingredients

● Chlorotoluron	250 g/l (21.7 % w/w)
● Diflufenican	40 g/l (3.5 % w/w)
● Pendimethalin	300 g/l (26.1% w/w)



Crops

A herbicide for use against a range of broad-leaved weeds and certain grass weeds in winter and spring wheat, winter and spring barley, winter and spring rye and winter and spring triticale.




SAFETY INFORMATION

Warning

Suspected of causing cancer.
 Suspected of damaging the unborn child.
 Very toxic to aquatic life with long lasting effects.
 Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.
 Keep out of reach of children.
 Obtain special instructions before use.
 Collect spillage.
 Wear protective gloves.
 Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.
 To protect aquatic organisms, respect an unsprayed buffer zone of 9 metres to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.



PCS 05425



Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/409Imfr>
Alternatively, contact your supplier.



R 3.0.8 Apr 23

LJJ1671IPFR_10

10Le





TOWER[®]



HERBICIDES

Suspension concentrate

PCS 05425



PROTECT FROM FROST

SHAKE WELL BEFORE USE

GROUP 5 | 12 | 3 HERBICIDES

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate and when handling contaminated surfaces.

TAKE OFF IMMEDIATELY all contaminated clothing.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

TOWER[®] may be subject to reduced buffer zones using STRIPE - Surface water Tool for Reducing the Impact of Pesticides in the Environment.

It is a legal requirement to maintain records and, to facilitate this, DAFM has produced an illustrative Plant Protection Product application

record sheet for professional users.

Be aware that when using nozzles producing coarser droplet sizes in conjunction with low water volumes and reduced product application rates, the efficacy of some products may be reduced depending on the situation. Consult ADAMA Agricultural Solutions UK Ltd.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and Disposal
KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

THIS MATERIAL AND ITS CONTAINER must be disposed of in a safe way.

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0)1865 407333 (24 hours, 7 days a week).

IMPORTANT INFORMATION FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE	
Crops:	Winter & spring wheat, winter & spring barley, winter and spring rye, winter and spring triticale
Maximum individual dose:	2 litres per ha
Maximum total dose:	-
Maximum no. of treatments:	1 per crop
Latest time of application:	Before pseudostem (leaf sheath) erect stage
Aquatic buffer zone distance:	9 metres
Other specific restrictions:	This product must not be applied via hand-held equipment. Substances containing the chlorotoluron active agent may not be applied more than once per year on the same area.

Authorised to & marketed by:
 ADAMA Agricultural Solutions UK Ltd
 Third Floor East, 1410 Arlington Business Park,
 Theale, Reading, RG7 4SA
 Tel: +44 (0)1635 860555
 Technical Helpline: +44 (0)1635 876622
 www.adama.com
 Email: ukenquiries@adama.com

Distributed in Ireland by: Croplink (Ireland) Ltd
 Springhill, Carrigtwohill, Co. Cork, T45 F685 Tel: 021 4882806

UFI: MA94-D1EF-W002-5D3M



5 056325 501093

R 3.0.8 Apr 23

L1167JIPBB_10



SAFETY INFORMATION



Warning

Suspected of causing cancer.

Suspected of damaging the unborn child.

Very toxic to aquatic life with long lasting effects.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.

Obtain special instructions before use.

Collect spillage.

Wear protective gloves.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To protect aquatic organisms, respect an unsprayed buffer zone of 9 metres to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 05425

This leaflet is part of the approved label.

TOWER[®] suspension concentrate formulation containing 250 g/l (21.7% w/w) chlorotoluron, 40 g/l (3.5% w/w) diflufenican and 300 g/l (26.1% w/w) pendimethalin. It is a herbicide for use against a range of broad-leaved weeds and certain grass weeds in winter and spring wheat, winter and spring barley, winter and spring rye and triticale.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS OR WARNINGS

Do not use if it is frosty.

Apply to crops that are healthy and not suffering stress from effects such as frost damage, if it is very dry or waterlogged, or a lack of nutrients.

Avoid overlapping spray swaths as this may lead to crop damage and reduced yields in unfavourable conditions.

DO NOT apply to soils with greater than 10% organic matter.

Prepare a firm, fine seedbed for best results with clods no more than fist size. The crop should be sown to the normal depth of 25 mm. It is important to ensure that the seed is well covered. Good weed control depends on the burying and dispersing of any trash or burnt straw before or during seedbed preparation.

For best results apply to a moist seedbed. Do not apply to dry, cracked or waterlogged soils.

Do not use on sands (Soil Texture 85 System) or very stony or gravelly soils due to risk of crop damage.

Do not allow spray drift onto neighbouring crops.

Crops should not be re-entered until spray residues are dry.

RESISTANCE MANAGEMENT

TOWER contains chlorotoluron a group 5 herbicide, diflufenican a group 12 herbicide and pendimethalin a Group 3 herbicide based on the mode of action classification system of the Herbicide Resistance Action Committee.

Strains of some annual grasses, e.g. black-grass, wild oats, and Italian rye-grass, have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the AHBD, CropLife UK, your distributor, crop adviser or product manufacturer.

Growers are advised to apply resistance management strategies which may include rotating and/or tank mixing with products having different modes of action against the target weeds.

Use the recommended rate of TOWER and the correct application timing for the hardest to control weed species present in the field. The above should be used in conjunction with effective cropping rotation and cultivation techniques e.g. stale seedbed, cultivation and use of non-selective herbicide prior to drilling.

WEEDS CONTROLLED

TOWER is a suspension concentrate formulation used as a herbicide against annual dicotyledonous weeds and grasses in winter and spring wheat, winter and spring rye, winter and spring barley and winter and spring triticale. It is absorbed by the roots, shoots and leaves. The soil should be moist at the time of application to give improved efficacy.

TOWER will control the following weeds up to the 2 leaf stage when applied early post-emergence to the crop:

Susceptible

Annual meadowgrass

Loose silky bent

Cleavers

Charlock

Common chickweed

Common fumitory

Common poppy

Cranesbill spp

Dead-nettle spp

Field pansy

Field pennycress

Forget-me-not

Mayweed spp

Runch

Shepherd's purse

Speedwell spp

Volunteer oilseed rape

Moderately susceptible

Rough stalked meadow grass

Moderately resistant

Black-grass

Resistant

Perennial weeds

Couch grass (Scutch)

Wild oats

CROP SPECIFIC INFORMATION

Timing

Apply TOWER pre or post-emergence up to before BBCH GS30 (before leaf sheath erect stage) to winter wheat, winter barley, winter rye and winter triticale.

Apply TOWER pre or post-emergence up to before BBCH GS30 (before leaf sheath erect stage) to spring wheat, spring barley, spring rye and spring triticale.

RATE OF USE

Apply TOWER at 2.0 L/ha in 150-300 L water/ha.

FOLLOWING CROPS

Following normal harvest, there are no restrictions. In the event of crop failure, plough before drilling the following crop in the spring. Only winter wheat can be re-drilled in the same autumn following crop failure.

Where diflufenican containing products are applied to successive cereal crops, levels of diflufenican will build up in the soil. Even with ploughing to 150mm and thorough mixing of the soil, there may still be a risk of damage to following crops of onions, leeks, other allium crops and clover. As a precaution, users who rent out their land to growers of these crops should not use diflufenican containing products in successive years before renting out that land.

MIXING AND SPRAYING

For use by tractor mounted/trailed sprayer only.

Ensure the sprayer is clean and in good working order. Calibrate according to the sprayer manufacturers recommendations.

Half fill the spray tank with clean water and begin agitation. Add the required quantity of TOWER to the tank and complete filling. Continue agitation until spraying is complete.

Do not leave the sprayer standing with the spray mixture in it for a long period of time.

Wash out the sprayer thoroughly after use, using a wetting agent or a proprietary tank cleaner.

Apply as a MEDIUM spray as defined by BCPC. Use a water volume of 150-300 L/ha.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorised to and marketed by:
ADAMA Agricultural Solutions UK Ltd
Third Floor East
1410 Arlington Business Park
Theale
Reading
RG7 4SA

Tel: +44 (0)1635 860555
Technical Helpline: +44 (0)1635 876622

Distributed in Ireland by:
Croplink (Ireland) Ltd
Springhill
Carrigtwohill
Co. Cork
T45 F685
Ireland

Tel: 021 4882806
www.adama.com
Email: ukenquiries@adama.com

TOWER® is a registered trademark of a company of the ADAMA Group.
© ADAMA Agricultural Solutions UK Ltd

Specimen-
2023 to date