

500g

tribe 75 wg

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

A water dispersible granule formulation containing tribenuron-methyl 750 g/kg (75%w/w) for use as a broad spectrum agricultural herbicide for post-emergence application in wheat, barley and oats, winter rye and winter triticale against annual broad leaf weeds and Creeping Thistle (*Cirsium arvense*).

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE.
THIS PRODUCT IS FOR PROFESSIONAL USE ONLY.

SAFETY INFORMATION



WARNING

May cause an allergic skin reaction.
May cause damage to organs through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects.

Do not breathe dust/mist/spray

Wear protective gloves/protective clothing

Get medical advice/attention if you feel unwell

IF ON SKIN: Wash with plenty of water.

Collect spillage.

Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste

Do not contaminate water with the product or its container. Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads.

Tractor mounted/trailed horizontal boom sprayer: to protect aquatic organisms, respect an unsprayed buffer zone of 5m to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use
PCS No: 05696

UFI: NJK4-F2JF-H80C-V15F

Batch number: see container

Registration holder: Sharda Cropchem España S.L.

Edificio Atalayas Business Center, Carril Condomina nº 3, 12th Floor, 30006 Murcia, Spain, Phone no.: +34 868 12 75 89

Distributed by: Sharda Cropchem España S.L.

Edificio Atalayas Business Center, Carril Condomina nº 3, 12th Floor, 30006 Murcia, Spain, Phone no.: +34 868 12 75 89

Emergency number: In the event of emergency call the National Poisons Information Centre, Beaumont Hospital at 01-8092166 or 01-8379964
Registered logos of Sharda Cropchem Ltd.



Sharda Cropchem

SHARDA CROPICHEM LIMITED

Domnic Holm 29th Road, Bandra (West)
Mumbai - 400050, India, Tel.: +91 22 6678 2800

READ THE LABEL CAREFULLY BEFORE USE

TRIBE 75 WG is a herbicide for the control of annual dicotyledonous weeds and Creeping Thistle (*Cirsium arvense*). Product contains tribenuron methyl, a herbicide from the sulphonyl urea class. It acts through inhibition of the enzyme aceto lactate synthase (ALS) involved in the synthesis of branched-chain amino acids. Inhibition of ALS leads to the cessation of cell division and subsequent growth processes in plants. Tribenuron methyl is classified by the Herbicide Resistance Action Committee (HRAC) as belonging to Group B.

Instruction for use:

Crop	Max single dose	Max no. Of applications	Max. Total dose	Latest time of application
Wheat, barley, oat, rye (winter) & triticale	20 g/ha	-	20 g/ha	Third Node Detectable GS 33
Setaside	20 g/ha	-	20 g/ha	N/A

Specific weeds controlled: chamomile weeds, corn chamomile, common chickweed, field nettles, hemp nettles, wild radish, charlock, field pennycress, shepherd's purse, field poppy, common lambsquarters, perennial sow thistle, creeping mistle.

Application Time – pre-harvest interval is determined by the time between the application and harvest.

Apply in 200-400 L water/ha using tractor mounted/trailed horizontal boom sprayer.
Max.number of applications: 1

Susceptible: *Chamomilla*, *Anthemis arvensis*, *Stellaria media agg.*, *Lamium*, *Galeopsis*, *Raphanus raphanistrum*, *Sinapis arvensis*, *Thlaspi arvense*, *Capsella bursapastoris*, *Papaver rhoeas*, *Chenopodium*, *Sonchus arvensis*, *Cirsium arvense*
Moderately susceptible: *Galium aparine*, *Fumaria officinalis*

Succeeding crops:

In the event of a crop failure for any reason, sow only a cereal crop within three months of application of TRIBE 75WG. After 3 months, field beans and oilseed rape may also be sown in the same year as application.

The product should not intervene surrounding vegetation or seeded land or land for sowing.

Risk of resistance and strategy to prevent the emergence of resistance

In order to reduce the development of resistance to tribenuron methyl it is important not to exceed the maximum number of applications per season as provided on the label.

Use only as part of a resistance management strategy that includes cultural method of control and does not use ALS inhibitors as the sole chemical method.

Mixing

Half fill the spray tank with clean water and start agitation. Pour the required quantity of product into the spray tank. Top up the spray tank with water to the required level. Continue agitation until the tank is sprayed out. Spray immediately after mixing.

Cleaning of application equipment

To avoid subsequent damage to crops other than cereals, immediately after spraying TRIBE 75 WG thoroughly clean all spray equipment including inside and outside of lid (spray tank, boom, hoses, filters and nozzles) using a proprietary sprayer cleaner (e.g. All Clear Extra®) according to the label instructions for that product.