



# HAKSAR<sup>®</sup>

## COMPLEX 260 EW

An emulsion in water formulation containing 200 g/l (18.23% w/w) of MCPA (present as potassium salt), 20 g/l (1.82% w/w) of clopyralid (present as monoethanolamine salt) and 40 g/l (3.65% w/w) of fluroxypyr (present as 1-methylheptyl ester).

A herbicide in the form of liquid to prepare an aqueous emulsion, foliar applied and designed for control of many dicotyledonous weeds in winter wheat and amenity grasslands.

**Causes skin irritation.**

**Causes serious eye damage.**

**Toxic to aquatic life with long lasting effects.**

Wear protective gloves/eye protection/face protection.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER/doctor.

If skin irritation occurs: Get medical advice/attention.

**Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).**

**To avoid risks to human health and the environment, comply with the instructions for use.**

**PROTECT FROM FROST. FOR PROFESSIONAL USE ONLY**

**EMERGENCY 24-HOUR CONTACT: 01 8092166**



UPL. E 6U-XOS0-U/0W D65U



5l

**Approval Holder**  
**CIECH Sarzyna S.A.**

ul. Chemików 1, 37-310 Nowa Sarzyna POLAND  
tel.: +48 17 24 07 111, fax: +48 17 24 07 122



# HAKSAR<sup>®</sup>

## COMPLEX 260 EW


**FOR USE ONLY AS AN AGRICULTURAL AND AMENITY HERBICIDE**

Crop	Max. Single Dose	Max. Total Dose	Latest Timing
Winter wheat	3.0 L/ha	3.0 L/ha	Before 2 <sup>nd</sup> node detectable (GS 32)
Grassland (amenity)	3.5 L/ha	7.0 L/ha	N/A

### Other specific restriction

**To protect aquatic organisms respect an unsprayed buffer zone of 10m to surface water bodies.**

For use in tractor mounted/trailed sprayers. Do not apply if rain is forecast within 48 hours. Do not apply to saturated soils. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.



**Distributor:**

**DHM Agrochemicals Ltd.**  
Castle Lodge, Kilgobbin Road,  
Dublin 18 IRELAND,  
Tel: +353 (01) 2952377



# HAKSAR COMPLEX 260 EW

## SAFETY PRECAUTIONS

### Operator protection

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work. WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

### Environmental Protection

To protect aquatic organisms respect an unsprayed buffer zone of 10m to surface water bodies.

To protect non-target plants and arthropods respect an unsprayed buffer zone of: 15m to non-agricultural land or 5m to non-agricultural land with the simultaneous application of techniques to reduce drifting of the spray liquid during the treatment by 50%.

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area. Avoid release to the environment.

### Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

### DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

## RESTRICTIONS

- DO NOT mow the lawn up to 3 days prior to the application of the product (so that weeds were the biggest) and for at least 3 days after treatment (to allow product transport to the roots).
- DO NOT use treated crops for feed.
- When spraying lawns near other crops, trees and shrubs, prevent the product from contact with crops, treetops and shrubs (especially on the green parts of plants). Minimise the possibility of contact of the liquid with the soil near shrubs and under trees.
- DO NOT use on newly established grass with poorly rooted plants.
- DO NOT use the grass from three successive mowings of the treated lawn for compost or mulching of any crops, ornamental plants, shrubs and trees.
- DO NOT spray in windy weather, avoid drift on to neighbouring crops, especially dicotyledonous.
- DO NOT apply in volumes less than 200 liters of water per hectare.
- DO NOT spray if frost is expected.
- DO NOT apply during drought.



Expiry date 3 years

Net content:

**5L**

Batch: On the packaging

Production date: On the packaging



**WEEDS CONTROLLED****Winter wheat**

SUSCEPTIBILITY	WEED SPECIES	DOSE	TIMING
Susceptible	Common Chickweed ( <i>Stellaria media</i> ) Red Dead-Nettle ( <i>Lamium purpureum</i> ) Scentless Mayweed ( <i>Matricaria inodora</i> ) Common Poppy ( <i>Papaver rhoeas</i> ) Creeping Thistle ( <i>Cirsium arvense</i> ) Common Speedwell ( <i>Veronica arvensis</i> ) Cleavers ( <i>Galium aparine</i> ) Volunteer Oilseed Rape ( <i>Brassica napus</i> ) Corn Chamomile ( <i>Anthemis arvensis</i> ) Scented Mayweed ( <i>Chamomilla recutita</i> ) Shepherd's-Purse ( <i>Capsella bursa-pastoris</i> ) Cornflower ( <i>Cyanus segetum</i> ) Fat Hen ( <i>Chenopodium album</i> ) Black Bindweed ( <i>Fallopia convolvulus</i> )	3.0 L/ha	Before 2 <sup>nd</sup> node detectable GS 21-32
Moderately Susceptible	Field Violet ( <i>Viola arvensis</i> )		

**Grassland - amenity**

SUSCEPTIBILITY	WEED SPECIES	DOSE	TIMING
Susceptible	Narrow Leaved Plantain ( <i>Plantago lanceolata</i> ) Common Chickweed ( <i>Stellaria media</i> ) Satin Flower ( <i>Stellaria holostea</i> ) Scentless Mayweed ( <i>Matricaria inodora</i> ) Common Sandelson ( <i>Taraxacum officinale</i> ) Greater Stitchwort ( <i>Stellaria holostea</i> ) Field Buttercup ( <i>Ranunculus acris</i> ) House-Ear Hawkweed ( <i>Lactuca pilosella</i> ) White Clover ( <i>Trifolium repens</i> ) Shamrock ( <i>Trifolium dubium</i> ) Red Clover ( <i>Trifolium pratense</i> ) Creeping Thistle ( <i>Cirsium arvense</i> ) Smooth Hawkbeard ( <i>Crepis capillaris</i> ) Common Fat's-Ear ( <i>Hypochoeris radicata</i> ) Hogweed ( <i>Erigeron canadensis</i> ) Cleavers ( <i>Galium aparine</i> ) Common Mouse-Ear ( <i>Cerastium pontanum</i> ) Common Horsetail ( <i>Equisetum arvense</i> ) Curled Dock ( <i>Rumex crispus</i> ) Sheeps Sorrel ( <i>Rumex acetosella</i> ) Broadleaf Dock ( <i>Rumex obtusifolius</i> ) Common Sorrel ( <i>Rumex acetosa</i> ) Shepherds Purse ( <i>Capsella bursa pastoris</i> ) Narrow Leaf Vetch ( <i>Vicia angustifolia</i> )	2.5 L/ha	from spring to early autumn

SUSCEPTIBILITY	WEED SPECIES	DOSE	TIMING
Susceptible	Broadleaf Plantain ( <i>Plantago major</i> ) Wall Speedwell ( <i>Veronica arvensis</i> ) Red Dead Nettle ( <i>Lamium purpureum</i> ) Common Nettle ( <i>Urtica dioica</i> ) Mugwort ( <i>Artemisia vulgaris</i> ) Creeping Buttercup ( <i>Ranunculus repens</i> )	3.5 L/ha	from spring to early autumn
Moderately Susceptible (susceptible with 2 treatments)	Cranesbill ( <i>Geranium Pusillum</i> ) Field Bindweed ( <i>Convolvulus arvensis</i> ) Ground Ivy ( <i>Glechoma Hederacea</i> )	3.5 L/ha	
Moderately Resistant	Persian Speedwell ( <i>Veronica persica</i> ) Yarrow ( <i>Achillea millefolium</i> )	3.5 L/ha	

### CROP SPECIFIC INFORMATION

The product should be applied to the entire area or to weeded areas only.

#### Winter wheat

Recommended Application Dose: 2.0 to 3.0 L product/ha

Maximum Application Dose: 3.0 L product/ha

Maximum Total Dose: 3.0 L product/ha

Maximum Number of Treatments: 1

Apply in spring after the vegetation starts from the beginning of tillering to 2nd node detectable (GS 21-32).

Apply in 200-300 L water/ha.

#### Grassland – amenity

Recommended Application Dose: 2.0 to 3.0 L product/ha

Maximum Application Dose: 3.0 L product/ha

Maximum Total Dose: 7.0 L product/ha

Maximum Number of Treatments: 2

Interval Between Applications: 60 days

Do not use HAKSAR COMPLEX 260 EW on new grassland with poorly rooted plants.

Apply HAKSAR COMPLEX 260 EW from spring to early autumn when growing conditions are favourable and the air temperature is above 10°C and below 25°C.

Apply in 200-500 L water/ha.

#### MIXING AND SPRAYING

Half-fill the tank with clean water and add the required quantity of HAKSAR COMPLEX 260 EW, and add the remainder of the clean water with gently agitation which should continue until the contents are thoroughly mixed. Spray using a MEDIUM/COARSE spray (BCPC).

#### TANK CLEANING

WASH EQUIPMENT thoroughly immediately after use. Rinse with water three times, drain and allow to dry. Traces of herbicide left

in the sprayer may damage susceptible crops if these are subsequently sprayed using the same equipment.

#### COMPATIBILITY

HAKSAR COMPLEX 260 EW can be tank-mixed with other pesticides, please consult your distributor.

#### RESISTANCE MANAGEMENT

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered to be resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

#### TERMS AND CONDITIONS OF SUPPLY, SALE OR USE

All goods supplied by CIECH Sarzyna S.A. are high grade and we believe them to be suitable for the purpose for which we expressly supply them: but as we cannot exercise any control over their mixing, use or application which may affect the performance of the goods all conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us or our Associate Companies for any damage or injury whatsoever arising from their storage, handling, re-application or use. These conditions cannot be varied by our staff, our agents or the re-sellers of the product whether or not they supervise or assist in the use of such goods.