

# LEGADO®

## FUNGICIDE

A suspension concentrate formulation for the control of cereal diseases



Contains 250 g/litre (22.93% w/w) azoxystrobin.

### SAFETY INFORMATION

#### WARNING

**Harmful if inhaled.  
Very toxic to aquatic life  
with long lasting effects.**

Wear protective gloves / protective clothing / eye protection / face protection. Use only outdoors or in a well-ventilated area.

Avoid breathing dust / fume / gas / mist / vapours / spray.

IF INHALED: Remove victim to fresh air and keep it rest in a position comfortable for breathing.

Call a POISON CENTRE or doctor / physician if you feel unwell.

Collect spillage.


Dispose of contents / container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains 1, 2-benzisothiazol-2(2H)-one. May produce an allergic reaction.

**To avoid risks to human health and the environment, comply with the instructions for use.**

Do not contaminate water with the product or its container (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

FOR PROFESSIONAL USE ONLY.



### IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

**Crops:** Wheat, barley, rye, triticale and oats.  
**Maximum single dose:** 1.0 L/ha.

**Maximum number of treatments:** Two per crop.

**Maximum total dose:** 2.0 L/ha.

**Latest time of application:** Before grain watery ripe stage (before GS 71).

**Other specific restrictions:** To reduce the risk of resistance developing in target diseases the total number of applications of product containing QoI fungicides made to any cereal crop must not exceed two.

**READ ALL OTHER SAFETY PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USE**  
USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE USING PLANT PROTECTION



8 435354 817356

ET-218e-IE-018-R00

PCS No. 06202

**PROTECT FROM FROST - SHAKE WELL BEFORE USE**

© Registered trademark of INDUSTRIAS AFRASA

Batch number: See  
Manufacture date: packaging  
Expiry date:



Hanseatic Trade Center, Kehrwieider 11  
20457 Hamburg, Germany

Content **5L**

Holder: **INDUSTRIAS AFRASA S.A.** Ciudad de Sevilla, 53 - 46988 Pol. Ind. Fuente del Jarro, Paterna (Valencia), SPAIN - Tel: (+34)961321700 - [afrasa@afrasa.es](mailto:afrasa@afrasa.es) - [www.afrasa.es](http://www.afrasa.es)

## SAFETY PRECAUTIONS

### Operator Protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the inside of gloves.

WASH SPLASHES from skin immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

### Environmental Protection:

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To protect aquatic organism, respect a 5 m buffer zone from surface water bodies.

### Storage and Disposal:

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

## DIRECTIONS FOR USE

**IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.**

LEGADO is a broad spectrum fungicide from strobilurin group, with systemic, trans-laminar and protectant properties, for the control of several diseases in cereals (wheat and barley).

Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stages of disease development or as a protectant treatment following a disease risk assessment or the use of appropriate decision support systems.

## CROPS

Crop	Max. single dose	Max. number of applications	Max. total dose	Latest time of application
Wheat, rye and triticale	1 L/ha	1	2 L/ha	Before grain watery ripe stage (GS 71)
Barley and oats	1 L/ha	2	2 L/ha	Before grain watery ripe stage (GS 71)

### Rate of use, Timing, interval of applications

Apply by foliar spray, at 1.0 L/ha, with a water volume 200-400 L water/ha, preferable in a preventive way at 14 days interval. It is recommended make two consecutive applications, first application during stem elongation (GS 31-39) and a second application before end of flowering (GS 69).

Latest Time of application: before grain watery ripe stage (before GS 71).

### Diseases controlled

**Wheat** (winter / spring): Control - Yellow rust (*Puccinia striiformis*), Brown rust (*Puccinia recondita*), Glume blotch (*Phaeosphaeria nodorum*), Cladosporium, Alternaria. Reduction - in severity of 'take-all'.

**Barley** (winter / spring): Control - Brown rust (*Puccinia horde*), Moderate control - Net blotch (*Pyrenophora teres*). Reduction - Leaf blotch (*Rhynchosporium commune*), and in the severity of 'take-all'.

**Rye:** Control - Brown rust (*Puccinia recondita*). Moderate control - Powdery mildew.

**Triticale:** Control - Yellow rust (*Puccinia striiformis*), Brown rust (*Puccinia recondita*), stoochy moulds. Moderate control - Powdery mildew.

### Processing

Where a crop is destined for processing, consult your processor before treating with LEGADO.

### Resistance Management

LEGADO contains azoxystrobin a member of the QoI cross resistance group. LEGADO should be used preventatively and should not be relied on for its curative potential.

Use LEGADO as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

Disease control may be reduced if strains of other pathogens less sensitive to azoxystrobin develop.

To reduce the risk of resistance developing in target diseases the total number of applications of product containing QoI fungicides made to any cereal crop must exceed two.

On cereal crops, LEGADO must always be used in mixture with another product, recommended for control of the same target disease that contains a fungicide from a different cross resistance group and is applied at a dose that will give robust control. Users should refer to current FRAG-UK guidelines for QoI compounds.

### Application and disposal details

Apply using a medium quality spray (BCPC) at a pressure of at least 1 bar. Apply through conventional crop spray equipment calibrated to give an even application at the current volume.

### Tank mixing

On cereal crops, LEGADO must always be used in mixture with another product, recommended for control of the same target disease that contains a fungicide from a different cross resistance group and is applied at a dose that will give robust control.

The following tank mixes have been tested for physical compatibility with LEGADO at recommended rates of use and will mix in the sprayer tank. No tests have been undertaken on crop safety or product performance therefore, use is at the user's own risk. Industrias Afrasa, S.A., will support the physical compatibility of 2 way tank mixes of LEGADO with any of the fungicides: BRAVO and TEBUCUR 250 EW; in case of mixtures with BRAVO, please use an anti-foaming agent for positive tank mixes. For further information on compatibilities, contact Industrias Afrasa, S.A.

Before using any tank mixture, consult and comply with the recommendations of the partner products. Each product should be added separately to the bulk of the water in the spray tank and thoroughly mixed before adding the next chemical. Always use constant agitation of the sprayer tank during mixing, transportation and application. Spray immediately.

Wash out containers thoroughly, preferably using an integrated pressure rinsing device, or manually rinsing three times. Add washings to the sprayer at time of filling.

### NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.