

Difenofin[®]



Emulsifiable Concentrate (EC) formulation containing 250 g/L difenoconazole
FOR USE ONLY AS AN AGRICULTURE FUNGICIDE. Please see inside for DIRECTIONS FOR USE

DIFENOFIN is a fungicide used to control a range of diseases in brussels sprouts and head cabbage.

FOR PROFESSIONAL USE ONLY

Risk and Safety Information

DANGER

May be fatal if swallowed and enters airways.

Causes serious eye irritation.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/ protective clothing/ eye protection/face protection.

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician.

IF IN EYES: Rinse cautiously for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.

DO NOT induce vomiting.

Collect spillage.

Store locked up.

Wash hands thoroughly after handling.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty triple rinsed clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

difenoconazole, 250 g/L

Authorisation holder:

Finchimica S.p.A.

Via Lazio, 13, 25025 Manerbio (BS), Italy

(Registered Company number: IT00573160983)

Technical Enq. +39 (0) 2 3569961 (business hours)

Emergency No. In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital at 01 809 2166 or 01 837 9964

www.finchimica.it
info@finchimica.it

© Finchimica S.p.A. registered mark

PROTECT FROM FROST

 **FINCHIMICA**
trust in agriscience

1 Litre e

Batch number: see packaging

SHAKE WELL BEFORE USE

PCS No. 06218

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY

Crop	Maximum individual dose	Maximum number of application	Maximum total dose	Latest timing of application
Brussels sprouts, head Cabbage	0.5 L/ha	-	1 L/ha	21 days before harvest

Other specific restrictions:

A minimum interval of 14 days must be observed between applications to brassica crops.

Applications to brassica must only be made to developed canopy and not before growth stage BBCH 41.

Risk Mitigation Measures:

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACE SHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces during spray application.

WASH SPLASHES from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

FOR USE VIA TRACTOR MOUNTED/TRAILED SPRAYERS ONLY.

Environmental Protection

Do not contaminate water with this product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains and farmyard roads).

Storage and disposal

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty triple rinsed clean containers which can be disposed of as non-hazardous waste.

DIRECTIONS FOR USE:

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

DIFENOFIN is a DMI fungicide with contact and systemic activity against a range of diseases

DISEASES CONTROLLED

Crop	Disease controlled
Brussels sprouts, cabbage	Leaf spot, Ring spot

CROP SPECIFIC INFORMATION

Apply preventatively or at the disease onset, to the crop having a minimum of 80% ground cover at time of first application.

Dose rate: 0.5 litres per hectare.

Apply in a recommended 200 litres of water per hectare (MEDIUM SPRAY).

MIXING, APPLICATION AND CLEANING

Make sure the sprayer is set at the correct volume. Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of DIFENOFIN and continue agitation while adding the rest of the water.

Agitate the mixture thoroughly before use and continue agitation during spraying.

Thoroughly wash all spray equipment with water immediately after use.

Method of Application: Tractor mounted/trailed sprayer.

RESISTANCE MANAGEMENT

DIFENOFIN is a DMI fungicide. Strains of Light Leaf Spot resistant to DMI fungicides are known to exist.

To avoid development of resistance, apply product protectively in response to disease forecasts.

Where possible, when Light Leaf Spot is present, avoid the use of DMI based fungicides when targeting other disease such as Sclerotinia at mid-flowering.

Finchimica S.p.A., as the seller, shall be under no liability (except for liability for death or personal injury resulting from the negligence of the seller) whether in contract or in tort or in respect of any loss or damage resulting from an arising out of the mixing or sequential use, of the goods with any other goods (whether the Seller or any third-party) otherwise than in accordance with the Seller's Recommendations for Use, or resulting from or arising out of the use of the goods in or before abnormal weather conditions or in unusual soil conditions notwithstanding that such conditions may be known or may have been known to the Seller or on plant varieties not known to the Seller to be abnormally susceptible to damage by the goods.

DIFENOFIN® is a registered mark of Finchimica S.p.A., Italy.