



ULTRALEGEND™



An emulsifiable concentrate formulation containing 125 g/L (12.8%) prothioconazole and 125g/L (12.8%) tebuconazole.

For use as an agricultural fungicide for the control of stem-base, foliar and ear diseases in winter and spring wheat (also reduction of the mycotoxin deoxynivalenol), winter rye, triticale and winter and spring barley, winter and spring oats and for disease control in winter oilseed rape.

FOR PROFESSIONAL USE ONLY.

ULTRALEGEND

Contains 125 g/L (12.8%) prothioconazole, 125 g/L (12.8%) tebuconazole and N,N-Dimethyl decanamide.

SAFETY INFORMATION

Warning

Causes skin irritation.

Causes serious eye irritation.

May cause respiratory irritation.

May cause an allergic skin reaction.

Suspected of damaging the unborn child.

Very toxic to aquatic life with long lasting effects.

Do not get in eyes, on skin or on clothing.

Avoid release to the environment.

Wear protective gloves/protective clothing/eye protection/face protection.

IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell.

Collect spillage.

Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

Contains 2-[2-(1-chlorocyclopropyl)-2-hydroxy-3-phenylpropyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use

PCS 06471

**PROTECT FROM FROST
SHAKE WELL BEFORE USE**

Transport Information 04/20
UN 3082 ADR/RID Class: 9 Packaging Group: III

Marketing Company Authorisation Holder
Crophthetics Ltd JT Agro Ltd
126-134, Baker Street, London W1U 6UE
Telephone: +44 (0)1628 421890
www.jtagro-crophthetics.com
Email: info@jtagro-crophthetics.com


crophthetics

net contents
5 litres e



IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops/situations	Maximum individual dose (Litres product/ha)	Maximum total dose (Litres product/ season)	Latest time of application
Wheat, triticale and winter rye	1.0	2.0	End of flowering (GS 69)
Barley and oats	1.0	2.0	Beginning of flowering (GS 61)
Oilseed rape (winter and spring)	1.0	2.0	56 days before harvest

Other specific restrictions:

Do not apply by hand held equipment.

Method of application: Tractor mounted sprayer.

A spray pressure of 2-3 bar is recommended. Apply ULTRALEGEND in 200-300 litres per hectare water.

Apply as a medium spray quality.

A minimum of 21 days must be observed between applications to the above crops.

SAFETY PRECAUTIONS

Operator Protection

Wash any contamination from eyes immediately.
Wash hands and exposed skin before meals and after work.

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms, respect an unsprayed buffer zone of 5M to surface water bodies. Spray must be aimed away from water

Storage and Disposal

Do not re-use container for any other purpose and dispose of safely.

Keep away from food, drink and animal feedingstuffs.

Keep out of reach of children

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

PROTECT FROM FROST.

STORE IN A COOL DRY PLACE.

READ ALL INSTRUCTIONS CAREFULLY BEFORE USE.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

ULTRALEGEND™ is a mixture of a triazolinthione fungicide and a triazole fungicide recommended for control of a wide range of diseases on winter and spring barley, winter and spring wheat, triticale, winter rye, winter and spring oats and for disease control in winter oilseed rape.

DISEASES CONTROLLED

CEREALS

	Wheat	Barley	Oats	Rye
Eyespot (<i>Oculimacula</i> spp.)	R	R	R	R
Septoria leaf blotch (<i>Mycosphaerella Graminicola</i>)	MC	-	-	-
Glume blotch (<i>Stagnospora nodorum</i>)	MC	-	-	-
Powdery mildew (<i>Blumeria graminis</i>)	MC	C	C	C
Yellow rust	C	C	-	-
Brown rust	C	C	-	C
Crown rust	-	-	C	-
Tan spot	MC	-	-	-
<i>Fusarium</i> ear blight	R	MC	-	-
<i>Sooty moulds</i>	R	R	-	-
<i>Rhynchosporium</i> leaf blotch	-	MC	-	C
Net blotch	-	C	-	-

C = Control MC = Moderate control R = Reduction

Eyespot (*Oculimacula* spp.)

ULTRALEGEND reduces the incidence and severity of eyespot. Spray in the spring at the first sign of disease, from when the leaf sheaths begin to become erect until the 2nd node is detectable (GS 30-32).

Septoria Leaf Spot and Glume Blotch (*Septoria tritici* and *Leptosphaeria nodorum*)

Apply before disease is established in the crop. To protect the upper leaves and ear apply ULTRALEGEND at full flag leaf emergence (GS 37) up to mid-flowering (GS 65). Where disease pressure remains high application may be repeated.

Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective.

Powdery Mildew (*Blumeria graminis*)

Apply ULTRALEGEND at the first signs of disease. Where disease pressure remains high application may be repeated.

Yellow Rust

Apply ULTRALEGEND at the first signs of disease. ULTRALEGEND controls yellow rust (*Puccinia striiformis*) in wheat. A second application may be made 3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

Brown Rust

Apply ULTRALEGEND at the first signs of disease. ULTRALEGEND controls brown rust in barley (*Puccinia hordei*), rye (*P. recondita*) and wheat (*P. recondita*).

A second application may be made 3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

Crown Rust (*Puccinia coronata*)

Apply ULTRALEGEND at the first signs of disease. ULTRALEGEND controls crown rust in winter and spring oats.

A second application may be made 3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

Tan Spot (*Pyrenophora tritici-repensis*)

Apply ULTRALEGEND at the first signs of disease in spring/early summer. Where disease pressure remains high application may be repeated.

Ear Disease Complex

Apply ULTRALEGEND soon after ear emergence until the end of flowering (GS 59-69) for control of *Fusarium* ear blight and reduction of sooty moulds. Control of ear diseases can result in cleaner, brighter ears.

Through the reduction of ear blight, ULTRALEGEND effectively reduces the level of the *Fusarium* mycotoxin deoxynivalenol (DON) in wheat grain. However, where *Fusarium* levels are high, the reduction achieved may not always be sufficient to ensure that DON levels fall below the statutory limit.

Leaf Blotch (*Rhynchosporium secalis*)

ULTRALEGEND gives high levels of *Rhynchosporium* control. Apply ULTRALEGEND in spring at the first signs of disease. For severe infections a second application may be necessary 3 weeks later.

Net Blotch (*Pyrenophora teres*)

Apply ULTRALEGEND at the first signs of disease in spring/early summer. For severe infections, a second application 3 weeks later will give most effective control when conditions remain favourable for disease development.

WINTER OILSEED RAPE

Light leaf spot	MC
<i>Phoma</i> leaf spot/stem canker	C
<i>Sclerotinia</i> stem rot	C

C = Control MC = Moderate control

Light Leaf Spot

Apply ULTRALEGEND in autumn/winter (usually late October to early December) protectively. Follow up spray(s) may be required in early spring from the onset of stem elongation, depending on disease development.

Phoma Leaf spot/Stem Canker

Apply ULTRALEGEND in autumn at the first sign of disease. Repeat application in late autumn/winter, if disease symptoms recur.

***Sclerotinia* stem rot (*Sclerotinia sclerotiorum*)**

Apply ULTRALEGEND at early to full flower.

RESISTANCE MANAGEMENT

Repeated application of ULTRALEGEND alone should not be used on the same crop against a high risk pathogen such as cereal powdery mildew. Tank-mixtures or alternation with fungicides having a different mode of action (e.g. morpholines) have been shown to protect against the development of resistant forms of disease.

CAUTION: The possible development of disease strains resistant to ULTRALEGEND cannot be excluded or predicted. Where such resistant strains occur, ULTRALEGEND is unlikely to give satisfactory control.

CROP SPECIFIC INFORMATION

ULTRALEGEND may be used on all commercial varieties of winter and spring barley, winter and spring wheat, triticale, winter rye, winter and spring oats and winter oilseed rape.

Mixing

Thoroughly shake the pack before use.

Add the required quantity of ULTRALEGEND to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

General

Sprayers should be thoroughly cleaned before use, and filters and jets checked for damage and blockages.

Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop. In dense crops, at later growth stages, higher water volumes should be used.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

Marketing Company

Crophthetics Ltd

126-134, Baker Street, London W1U 6UE

Telephone: +44 (0)1628 421890

www.jtagro-crophthetics.com

Email: info@jtagro-crophthetics.com

Authorisation Holder

JT Agro Ltd

ULTRALEGEND™ is a trademark of companies of JT Agro Ltd.