

Difure[®] Solo

**A systemic fungicide for use
on all varieties of sugar beet
and fodder beet.**



FUNGICIDE

**Emulsifiable concentrate
containing 250 g/L
of difenoconazole
(23.6 % w/w).**

RISK AND SAFETY INFORMATION

**Harmful if swallowed.
May be fatal if swallowed and enters airways.
Causes serious eye irritation.
May cause damage to organs through pro-
longed or repeated exposure.
Very Toxic to aquatic life with long lasting
effects.**

Keep out of reach of children.
Avoid release to the environment.
Wear eye protection/face protection.
IF SWALLOWED: Immediately call a POISON
CENTER or doctor/physician.

IF IN EYES: Rinse continuously with water for
several minutes. Remove contact lenses if pres-
ent and easy to do – continue rinsing.
Do NOT induce vomiting.

If eye irritation persists: Get medical advice/
attention.

Dispose of contents/container to a
licensed hazardous-waste disposal
contractor or collection site except
for empty triple rinsed contain-
ers which can be disposed of
as non-hazardous waste.

**Repeated exposure may
cause skin dryness or cracking.
To avoid risks to human health
and the environment, comply with
the instructions for use.**



DANGER

FOR PROFESSIONAL USE ONLY

PCS no. 06537

Difure[®] is a registered trademark for Globachem N.V.

PROTECT FROM FROST
SHAKE WELL BEFORE USE
BATCH NUMBER: SEE BOTTLE

CONTENT: 1 L

For 24-hour
emergency information contact
Globachem n.v.
Telephone: +32 11 78 57 17

Registration holder:

GLOBACHEM NV
Brustem Industriepark
Lichtenberglaan 2 • 19
B-3800 Sint-Truiden • Belgium
Tel: +32 11 78 57 17
Fax: +32 11 68 15 65
globachem@globachem.com
www.globachem.com


Globachem
Your Crop Counts

**SAFETY PRECAUTIONS
OPERATOR PROTECTION**

Engineering control of operator expo-
sure must be used where reasonably
practical in addition to the following
personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTH-
ING (COVERALLS), GLOVES AND FACE
PROTECTION (FACESHIELD), when
handling the concentrate.

WASH HANDS AND EXPOSED SKIN
before meals and after work.
WHEN USING, DO NOT DRINK, EAT OR
SMOKE.

IF YOU FEEL UNWELL, seek medical
advice immediately and show this
container or label.

IN CASE OF CONTACT WITH EYES,
RINSE IMMEDIATELY with plenty of
water and seek medical attention.

ENVIRONMENTAL PROTECTION

To protect aquatic organisms respect
an unsprayed buffer zone of 5m to
surface water bodies.

DO NOT CONTAMINATE WATER with
the product or its container. Do not
clean application equipment near
surface water. Avoid contamination
via drains from farmyards and roads.

STORAGE AND DISPOSAL

KEEP IN ORIGINAL CONTAINER, tightly
closed, in a safe place.

WASH OUT CONTAINER THOROUGH-
LY, empty washings into spray tank
and dispose of safely.

KEEP OUT OF REACH OF CHILDREN.
Do not re-use container for any pur-
pose.

KEEP AWAY FROM FOOD, DRINK AND
ANIMAL FEEDING STUFF.

THIS MATERIAL AND ITS CONTAINER
must be disposed of in a safe way.
Store in a cool, dry place.

DIRECTIONS FOR USE

IMPORTANT: This information is ap-
proved as part of the product label. All
instructions within this section must
be read carefully in order to obtain
safe and successful use of this product.

IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

CROPS	MAXIMUM INDIVIDUAL DOSE (L PRODUCT PER HECTARE)	MAX NUMBER OF APPLICATIONS (PER CROP)	LATEST TIME OF APPLICATION
Sugar beet & Fodder beet	0.5	2	21 days before harvest

Other specific restrictions: A minimum interval of 14 days must be observed between applications.
Do not apply before BBCH 39.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.

GENERAL INFORMATION

DIFURE SOLO is a systemic fungicide
containing difenoconazole which is
an active ingredient belonging to the
chemical family of triazoles (FRAC
code 3) for use on fodder and sugar
beet. DIFURE SOLO is aimed at con-
trolling rust (*Uromyces betae*).

DIFURE SOLO quickly penetrates into
the beet leaves and protects them for
several weeks.

CROP SPECIFIC INFORMATION

Apply DIFURE SOLO at 0.5 L/ha at the
first signs of the disease, before the
disease becomes established. DIFURE
SOLO gives prolonged protection
from re-infection, but a second ap-
plication can be applied where crops
are at risk from later attacks. Respect
a minimum application interval of 14
days between applications.

Do not apply before BBCH 39 (Crop
over complete: leaves cover 90% of
ground).

MIXING AND SPRAYING

Make sure the sprayer is clean and
set to give an even application at the
correct volume.

Fill the spray tank with half the re-
quired volume of clean water and
start agitation. Add the required
amount of DIFURE SOLO and continue
agitation while adding the rest of the
water. Agitate the mixture thorough-
ly before use and continue agitation
during spraying.

Do not leave the spray liquid in the
sprayer for long periods (such as dur-
ing meal breaks or overnight).

Thoroughly wash all spray equipment
with water immediately after use.

WATER VOLUME

Apply DIFURE SOLO in a recommended
100-1000 litres of water per hectare.
A MEDIUM quality spray (BCPC) is
preferred for application of DIFURE
SOLO. A spray pressure of 2-3 bars is
recommended.

COMPATIBILITY

The mixtures should be performed in
accordance with the relevant regula-
tions and the official recommenda-
tions of good agricultural practices.
For up to date details of compatible
tank-mixes contact your manufactur-
er/distributor.

RESISTANCE

DIFURE SOLO contains difenocona-
zole, a DMI fungicide (FRAC code 3).
Avoid repeated use of fungicides
having the same mode of action.
Strains of Light Leaf Spot resistant
to DMI fungicides are known to exist.
To avoid development of resistance,
apply this product protectively in re-
sponse to disease forecasts.

For some further advice on resistance
management in DMI's contact your
distributor, crop adviser or product
manufacturer.

COMPANY INFORMATION

All goods supplied by us are of high
grade and we believe them to be suit-
able but, as we cannot exercise con-
trol over their storage, handling, mix-
ing or use, or the weather conditions
before, during or after application
which may affect the performance
of the goods, all conditions and war-
ranties, statutory or otherwise, as
to the quality or fitness for any pur-
pose of our goods are excluded. No
responsibility will be accepted by us
or re-sellers for any failure in perfor-
mance, damage or injury whatsoever
arising from their storage, handling,
application or use. These conditions
cannot be varied by our staff or
agents whether or not they supervise
or assist in the use of such goods.

Our company is not liable for any
consequences resulting from the mix-
tures with other products.



ET13877-ED1-V1



IE