

### Risk and safety information

Contains ipconazole and imazalil

FOR PROFESSIONAL USE ONLY

#### WARNING:

**VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS.  
SUSPECTED OF CAUSING CANCER.**

OBTAIN SPECIAL INSTRUCTIONS BEFORE USE. DO NOT HANDLE UNTIL ALL SAFETY PRECAUTIONS HAVE BEEN READ AND UNDERSTOOD.

AVOID RELEASE TO THE ENVIRONMENT.

WEAR PROTECTIVE GLOVES/ PROTECTIVE CLOTHING/ EYE PROTECTION/ FACE PROTECTION.

IF EXPOSED OR CONCERNED: GET MEDICAL ADVICE/ATTENTION.

COLLECT SPILLAGE.

STORE LOCKED UP.

DISPOSE OF CONTENTS/CONTAINER TO A LICENSED HAZARDOUS-WASTE DISPOSAL CONTRACTOR OR COLLECTION SITE EXCEPT FOR TRIPLE-RINSED EMPTY CLEAN CONTAINERS WHICH CAN BE DISPOSED OF AS NON-HAZARDOUS WASTE

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farm yards and roads.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 06588



### IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL SEED TREATMENT.

#### Crops:

Barley (seed)  
Wheat (seed)

**Maximum individual dose:** 1.0 L product per 1000 kg seed

**Maximum number of treatments:** One per batch

**Latest time of application:** Pre-drilling

**Other specific restrictions:** Returnable containers must not be re-used for any other purpose. Returnable containers must be returned to the supplier. Treated seed must not be used for food or feed. Sacks containing treated seed must not be re-used for food or feed. Treated seed must not be applied from the air.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE PRINCIPLES OF GOOD PLANT PROTECTION PRACTICE.**

### A micro-emulsion seed treatment for the control of seed borne diseases in wheat and barley

Micro-emulsion formulation for seed treatment containing 20 g/L ipconazole and 50 g/L imazalil

AGITATE WELL BEFORE USE

### Authorisation Holder and Marketing Company:

Arysta LifeScience Great Britain Ltd  
3-5 Melville Street, Edinburgh EH3 7PE  
Tel: 01924 888151

### Distributing Company:

Arysta LifeScience Great Britain Ltd.  
3-5 Melville Street, Edinburgh  
EH3 7PE  
United Kingdom  
Tel: 01924 888151

**In case of toxic or transport emergency,  
ring +44 (0) 1235 239 670 (24 hour).**

Batch No. See Container

### Resistance Management:

At the time of printing, there are no cases of resistance to Conima known to Arysta LifeScience. The possible development of diseases resistant to Conima cannot be excluded or predicted. Conima contains ipconazole (a triazole fungicide) and imazalil (an imidazole fungicide), both of which belong to the DMI-fungicide group (SBI: Class I). Disease control may be reduced accordingly if strains of fungi resistant to ipconazole or imazalil develop.

The use of Conima should conform to resistance management strategies. Such strategies should include subsequent foliar applications being made with products containing an active substance with a different mode of action. For further advice on resistance management in DMI's consult your agronomist or specialist advisor, and visit the Fungicide Resistance Action Group (FRAG-UK) website.

### SAFETY PRECAUTIONS (when handling the product)

#### Operator exposure

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate, contaminated surfaces or treated seed.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when bagging treated seed.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WASH CONCENTRATE from skin or eyes immediately.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

WHEN USING DO NOT EAT, DRINK OR SMOKE.

#### Environmental protection

To protect birds/wild mammals, treated seed should not be left on the soil surface. Bury or remove spillages.

#### Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY and dispose of safely.

DO NOT RE-USE CONTAINER FOR ANY OTHER PURPOSE.

### SAFETY PRECAUTIONS (when handling treated seed).

**LABEL TREATED SEED with the appropriate precautions using printed racks, labels or bag tags.**

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling treated seed.

DO NOT HANDLE seed unnecessarily.

DO NOT USE TREATED SEED as food or feed.

KEEP TREATED SEED SECURE from people, domestic stock, pets and wildlife at all times during storage and use.

To protect birds/wild mammals, treated seed should not be left on the soil surface. Bury or remove spillages.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE BEEN USED FOR TREATED SEED for food or feed.

WASH HANDS AND EXPOSED SKIN before meals and after work.

DO NOT APPLY TREATED SEED FROM THE AIR.

### DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

### RESTRICTIONS

Do not treat seed with a moisture content exceeding 16%.

Do not apply Conima to cracked, split or sprouted seed.

Seed treated with Conima should be drilled within six months of treatment.

Do not store seed treated with Conima from one season to the next. Arysta LifeScience accepts no liability for the performance of stored, treated seed.

Flow of seed through drills may be affected by Conima. Always re-calibrate your seed drill with treated seed before drilling.

### Disease control

Conima is a micro-emulsion seed treatment for the control of seed borne diseases in wheat and barley.

### Application rates

Barley: 1.0 L product per 1000 kg of seed

Wheat: 1.0 L product per 1000 kg of seed

CROP	TARGET/CLAIM	RATE OF Conima PER 1000 KG SEED*
Barley	Control of loose smut ( <i>Ustilago nuda</i> ) in winter and spring barley. Control of leaf stripe ( <i>Pyrenophora graminea</i> ) in spring barley. Conima improves crop establishment by giving a reduction in seedling blight and foot rot caused by <i>Fusarium</i> species and <i>Microdochium nivale</i>	1.0 L
Wheat	Control of seed and soil-borne bunt ( <i>Tilletia caries</i> ). Reduction of seedling blight caused by <i>Fusarium</i> species and <i>Microdochium nivale</i>	1.0 L

\* Conima can be applied neat but co-application with 3L water may improve coverage

### Application

Apply Conima through a conventional seed treater suitable for handling liquid products. Co-application of Conima with water is likely to improve the evenness of coverage of the seed with Conima.

Agitate well before use.

### Cleaning of application equipment

Do not transfer pumps from one treatment to another without cleaning. When changing from Conima, wash out the pump and pipeline with clean water using a double rinse procedure before use.

### Compatibility

Conima should not be pre-mixed with any other seed treatment product.

We accept responsibility for this product meeting the declared specification and also for our written directions for use and warnings. Users however should beware that no responsibility can be accepted for use which disregards our written instructions, including directions, warnings and precautions, nor for any advice given by any person whether verbal or written, about storage, mixing and application, or about any use which is not strictly in accordance with our written instructions.