

sitara

A suspension emulsion formulation containing 100 g/litre fluroxypyr and 2.5 g/litre florasulam. A post-emergence herbicide for use on all varieties of WINTER AND SPRING WHEAT, BARLEY, OATS, RYE and TRITICALE for the control of a range of BROAD LEAVED WEEDS

5L

UFI : M3D6-6KPA-8G02-AVFS



**FOR USE ONLY AS AN AGRICULTURAL HERBICIDE
FOR PROFESSIONAL USE ONLY**

SAFETY INFORMATION



DANGER

May be fatal if swallowed and enters airways.
Causes skin irritation.
May cause an allergic skin reaction.
Causes serious eye irritation.
May cause respiratory irritation.
May cause drowsiness or dizziness.
Very toxic to aquatic life with long lasting effects.

Wear protective gloves/clothing/eye/face protection.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: Get medical advice/attention.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician.

Do NOT induce vomiting.

Collect Spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads). To protect aquatic organisms, respect an unsprayed buffer zone of 5m to surface water bodies.

Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No: 06627

Safety Precautions

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental protection:

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from yards and roads.

Batch no.: See container

Registration holder: Sharda Cropchem España S.L.

Edificio Atalayas Business Center,
Carril Condomina nº 3, 12th Floor, 30006
Murcia, Spain

Phone no.: +34 868 12 75 89

Distributed by: Sharda Cropchem España S.L.

Emergency no.: In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital at 01 809 2166 or 01 837 9964
Registered logos of Sharda Cropchem Ltd.

Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

SHAKE WELL BEFORE USE. PROTECT FROM FROST.

If exposed or if you feel unwell seek medical advice immediately (show the label where possible).

No specific antidote is available. Treat symptomatically.

In case of accident, call the 24-hour National Poisons Information Centre:

Tel.: +353 (0)1 809 2566 or +353 (0) 1 837 9964.

General information:

SITARA herbicide is mainly absorbed through the foliage of weeds. The ideal timing for application is when the weeds are small and actively growing.

FOR PROFESSIONAL USE ONLY AS FOR USE ONLY AS AN POST-EMERGENCE HERBICIDE				
Crop	Max. single dose	Max. no. of applications	Max. total dose	Latest time of application
Winter wheat, winter barley	1 L/ha	1 per year	1 L/ha	Before GS 39
Spring wheat, Durum wheat, Spring barley, Winter rye, Spring rye, Triticale, Winter oats, Spring oats	1 L/ha	1 per year	1 L/ha	Before GS 32

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Specific restrictions:

For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1st in the year of harvest.

The total amount of florasulam applied to a cereal crop must not exceed 7.5 g

Method of application: SITARA may be applied through tractor-mounted hydraulic sprayers providing they are in good working order and have been calibrated according to the manufacturers' recommendations. Do not apply through CDA applicators.

Important information:

Do not spray when crops are under stress from cold, drought, pest damage, nutrient deficiency etc.

Do not roll or harrow 7 days before or after application.

Take extreme care to avoid drift onto crops and non-target plants outside the target area.

Mixing, application and cleaning procedures:

Mixing information: Half fill the spray tank with water and add the required amount of SITARA. Fill up the spray tank, agitating continuously to ensure thorough mixing, and maintain agitation until spraying is complete. Use only clean water for mixing. For overall application apply SITARA in 150 to 250 litres of water per hectare. The lower volume must only be used in open crops on small weeds.

Following crops: For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1st in the year of harvest.

Joint application: A joint application is the use of a product in tank mixture or sequence with another product. Joint applications should only be made within the label recommendations of every product in the application. Only one other product with an ALS-inhibitor mode of action may be applied to a cereal crop treated with SITARA. However, a further application of SITARA or another product containing florasulam may also be made providing the maximum total dose of florasulam is not exceeded.

SITARA may be applied in joint application to the same cereal crop with one of the following ALS products:

Ally Express	Harmony M SX
Ally Max SX	Harmony SX
Ally SX	Presite SX
Biplay SX	Quantum
Broadway Star ¹	Spitfire ¹
Calibre SX	Starane XL ¹
Finish SX	Taxi
Galaxy	

¹The maximum total dose of florasulam applied to the crop must not exceed 7.5 g. For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1st in the year of harvest. Apart from these specific joint applications SITARA must NOT be applied with any other product containing an ALS-inhibitor, for example amidosulfuron or triasulfuron.

Tank mixtures: Where tank mixes are used, and unless directed otherwise, the preferred order of addition of product to the spray tank is as follows: water dispersible granules, wettable powders, suspension concentrates (flowable), emulsifiable concentrate, solution concentrates. Each product should be added to the half-full sprayer and be fully dispersed before the addition of the next product.

Tank Cleaning: To avoid subsequent injury to crops other than cereals, all spraying equipment must be thoroughly cleaned both inside and out, using All Clear Extra spray cleaner as follows:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water and add All Clear Extra at the recommended rate. Agitate and then briefly flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full and allow to stand for 10 minutes with agitation. Flush the boom and hoses and drain tank completely.
4. Nozzles and filters should be removed and cleaned separately with All Clear Extra solution containing 50 ml of All Clear Extra per 10 litres of water.
5. Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume. Drain tank completely.
6. For disposal of washings, follow Code of Practice for Using Plant Protection Products.

Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

Specific pest controlled:

Weed	Size controlled
Cleavers	Flowering
Chickweed	Flowering
Mayweed	Flowering