

pastisha

An emulsifiable concentrate containing 240 g/L Triclopyr (present as 334 g/L of Triclopyr butyl) and 60 g/L Clopyralid for use as an agricultural herbicide on established grassland and amenity grassland

1 L

**FOR USE ONLY AS AN AGRICULTURAL HERBICIDE
THIS PRODUCT IS FOR PROFESSIONAL USE ONLY**

SAFETY INFORMATION



DANGER

- Highly flammable liquid and vapour.
- May be fatal if swallowed and enters airways.
- Causes skin irritation.
- May cause an allergic skin reaction.
- Causes serious eye irritation.
- May cause respiratory irritation.
- May cause drowsiness or dizziness.
- May cause damage to aquatic life through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects

Keep away from heat, hot surfaces, sparks, open flames and other ignition sources.

No smoking.

Avoid breathing dust/fume/gas/mist/vapours/spray.

Wear protective gloves/protective clothing/eye protection.

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing.

Do NOT induce vomiting.

Collect spillage.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean triple rinsed containers which can be disposed of as non-hazardous waste.

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water, avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No: 06629

Precautions

If exposed or if you feel unwell seek medical advice immediately (show the label where possible).

No specific antidote is available. Treat symptomatically.

In case of accident, call the 24-hour National Poisons Information Centre:

Tel.: +353 (0)1 809 2566 or +353 (0)1 837 9964.

Batch no.: See container

Registration holder:

Sharda Cropchem España S.L. Edificio Atalayas Business Center, Carril Condomina nº 3, 12th Floor, 30006, Murcia, Spain

Phone no.: +34 868 12 75 89

Distributed by: Sharda Cropchem España S.L.

Emergency no.: In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital at 01 809 2166 or 01 837 9964

General Information

PASTISHA is a foliar acting herbicide for the control of NETTLES, DOCKS, THISTLES, BRAMBLES, BROOM and GORSE in established grassland and amenity grassland.

Safety precautions

Operator protection

- Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
- Operators must wear SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES, RUBBER BOOTS and FACE PROTECTION (FACESHIELD) when handling the concentrate.
- Operators must wear SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES and RUBBER BOOTS when applying by hand-held equipment.
- Operators must wear SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.
- WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.
- DO NOT APPLY by hand-held rotary atomiser equipment.
- When using DO NOT EAT, DRINK OR SMOKE.

Consumer protection

NOT TO BE USED ON FOOD CROPS.

Protection of environment

- Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.
- To protect aquatic organisms, respect an unsprayed buffer zone of 5m to surface water bodies.
- Livestock must be kept out of treated areas for at least 7 days following treatment. If ragwort is present, the guidance in the 'DIRECTIONS FOR USE' must be followed.
- DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards or roads.
- Applications must be made outside the period of 1st March to 31st of October.

Storage and disposal

- KEEP OUT OF REACH OF CHILDREN.
- KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. RE-USE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.
- DO NOT RE-USE CONTAINER for any purpose.
- PROTECT FROM FROST.

Instructions for use

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Max. single dose	Max. no. of applications	Max. total dose	Latest time of application
Grassland (established) Grassland (amenity)	1.2 L/ha	-	1.2 L/ha	-

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Other specific restrictions:

- Livestock must be kept out of treated areas for at least 7 days following treatment. If ragwort is present, the guidance in the 'DIRECTIONS FOR USE' must be followed.
- Do not apply through hand-held rotary atomiser equipment.
- Applications must not be made outside the period of 1st March to 31st October.
- The maximum concentration must not exceed 60 ml product per 10 litres of water (6 ml product per litre of water).

Method of Application:

- Tractor mounted/trailed horizontal boom sprayer or knapsack sprayer.
- DO NOT APPLY through hand-held rotary atomizer equipment.
- Apply PASTISHA using a handheld sprayer. Ensure the equipment is in good working order and has been calibrated according to the manufacturers' recommendations.
- The weeds should be thoroughly wetted with the spray solution but spraying until "run-off" will decrease activity.
- The use of flood jets is recommended to prevent drift. Care should be taken to avoid local overdosing. The maximum concentration must not exceed 60 ml product per 10 litres of water (6 ml product per litre of water).

Application timing:

- If the grass has been cut for hay or silage or grazed leave for 2-3 weeks to allow sufficient re-growth to occur before spraying. Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass.

- To allow for maximum translocation of PASTISHA to the roots, do not cut grass for 28 days after application.

Cleaning procedure:

- Wash out spray equipment thoroughly with water and detergent immediately after use. Traces of PASTISHA could cause harm to susceptible crops sprayed later.

Warnings:

Read all warnings before applying PASTISHA.

- Do not roll or harrow grass for 10 days before or 7 days after application. Do not spray in drought, very hot or very cold weather.

- PASTISHA is safe to grass. Very occasionally some yellowing of the sward may occur; this is transient and quickly outgrown.

- Clover will be killed or severely checked by application of PASTISHA.

- DO NOT sow kale, swedes, turnips or grass mixtures containing clover by direct drilling or minimum cultivation techniques within 6 weeks of applying PASTISHA.

- PASTISHA residues in plant tissues (including manure and digestate) which have not completely decayed may affect succeeding susceptible crops (e.g. peas, beans and other legumes, carrots and umbelliferae, potatoes and tomatoes, lettuce and other compositae. Do not plant susceptible autumn-sown crops (e.g. winter beans) in the same year as treatment with PASTISHA. Where susceptible beans are to be planted in the spring do not apply PASTISHA later than the end of July of the previous year. Following good agricultural practice ensure that plant remains have completely decayed before planting susceptible crops.

- Do not use any plant material treated with PASTISHA for composting or mulching.

- Do not use manure from animals fed on crops treated with PASTISHA for composting.

- In hot conditions, vapour drift may occur, making it particularly important to assess the risk to neighbouring vegetation.

- Take extreme care to avoid drift onto crops and non-target plants outside the target area.

- Do not apply directly to, or allow spray drift to come into contact with, agricultural or horticultural crops, amenity plantings or gardens, ponds, lakes or water courses.

- All conifers, especially pine and larch, are very sensitive to spray drift.

Specific weeds controlled:

- Common nettle: before flowering (normally up to mid-June)
- Curled dock & Broad-leaved dock: Rosette stage up to 25 cm high. On large well-established docks, or where there is a high reservoir of seed in the soil, a second application the following year may be required.
- Creeping thistle: Before flowering spikes are 15 cm high.
- Bramble, broom and gorse: Spray in June-August when actively growing but before plants begin to senesce in the autumn. It is essential that, particularly with large bushes, all the foliage is thoroughly wetted or incomplete kill may result.



Sharda Cropchem

SHARDA CROPCHEM LIMITED
Domnic Holm 29th Road, Bandra (West)
Mumbai - 400050, India, Tel. +39 02 66101029