



STEMPER®



ADAMA



PLANT GROWTH REGULATOR

Emulsifiable concentrate

PCS 06676

FOR PROFESSIONAL USE ONLY



Ingredients

- Trinexapac-ethyl 175 g/L (17.2% w/w)



Crops


A growth regulator for reduction of crop height, which can lead to control of lodging and yield protection in winter and spring wheat, winter and spring barley, winter and spring oats, durum wheat, rye, triticale, and ryegrass seed crops.



SAFETY INFORMATION

Warning

Causes skin irritation.
 May cause an allergic skin reaction.
 Causes serious eye irritation.
 May cause damage to organs through prolonged or repeated exposure.
 Toxic to aquatic life with long lasting effects.
 Keep out of reach of children.
 Wear protective gloves, protective clothing, eye protection & face protection.
 IF ON SKIN: Wash with plenty of soap and water
 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
 Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.
 To avoid risks to human health and the environment, comply with the instructions for use.



PCS 06676



Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/3SEYRhp>
Alternatively, contact your supplier.



R 3.0 Mar 23

LJJ1556IPFR_01



STEMPER[®]



PLANT GROWTH REGULATOR

Emulsifiable
concentrate

PCS 06676

PROTECT FROM FROST

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0)1865 407333 (24 hours, 7 days a week).

IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL PLANT GROWTH REGULATOR

Crops: Winter & spring wheat; winter & spring barley; winter & spring oats; rye, triticale, durum wheat; grassland (seed crop)

Maximum individual dose; Maximum total dose; Latest time of application and Other specific restriction:

Full details are given in the information box within the leaflet attached to the product.

L11556IPBB_01

Authorised to & marketed by:
ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA

Tel: +44 (0)1635 860555

Technical Helpline: +44 (0)1635 876622

www.adama.com

Email: ukenquiries@adama.com

Distributed in Ireland by: Croplink (Ireland) Ltd, Springhill,
Carrigtwohill, Co. Cork, T45 F685 Tel: 021 4882806

R 3.0 Mar 23

UFI: U31H-R3YK-F00U-M7M9



5 056325 502489

Open here 



Directions For
Use

SAFETY INFORMATION

Warning

Causes skin irritation.

May cause an allergic skin reaction.

Causes serious eye irritation.

May cause damage to organs through prolonged or repeated exposure.

Toxic to aquatic life with long lasting effects.

Keep out of reach of children.

Wear protective gloves, protective clothing, eye protection & face protection.

IF ON SKIN: Wash with plenty of soap and water

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



PCS 06676

This booklet is part of the approved label.

An emulsifiable concentrate formulation containing 175 g/l (17.2% w/w) trinexapac-ethyl.

A growth regulator for reduction of crop height, which can lead to control of lodging and yield protection in winter and spring wheat, winter and spring barley, winter and spring oats, durum wheat, rye, triticale, and eye grass seed crops.

IMPORTANT INFORMATION FOR USE ONLY AS A PROFESSIONAL PLANT GROWTH REGULATOR

Crop	Maximum individual dose (L product/ha)	Maximum total dose (L product/ha)	Latest time of application*
Winter wheat	0.6	0.6	Before flag leaf sheath extending stage (GS 41)
Spring wheat	0.6	0.6	Before third node detectable (GS 33)
Winter barley	0.9	0.9	Before flag leaf sheath extending stage (GS 41)
Spring barley	0.7	0.7	Before third node detectable (GS 33)
Winter & spring oats	0.6	0.6	Before second node detectable (GS 32)
Rye, triticale, durum wheat	0.6	0.6	Before third node detectable (GS 33)
Grassland (seed crop)	1.1	1.1	Before second node detectable (GS 32)
Other Specific Restriction			
This product must not be used on grass seed crops that will be grazed by livestock or cut for fodder.			

*See the CROP SPECIFIC INFORMATION section for details of the earliest and latest time of application for each recommendation.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

STEMPER® is a growth regulator for crop height reduction, which can lead to control of lodging and yield protection in all varieties of winter and spring wheat, winter and spring barley, winter and spring oats, rye, triticale, durum wheat and ryegrass seed crops.

RESTRICTIONS

Avoid spray drift onto neighbouring crops.

Only use on crops at risk of lodging.

Do not apply if rain or frost is expected or if the crop is wet.

Do not spray crops under stress or to crops suffering from water logging, pest attack, disease or frost.

Take particular care to avoid overlapping of sprayed strips.

Treatment may lead to cereal ears remaining erect through to harvest.

Crops should not be re-entered until spray residues are dry.

CROP SPECIFIC INFORMATION

Crop	Rate of use (L/ha)	Timing
Winter wheat	0.6	Between tillers erect stage and the flag leaf fully emerged stage (GS 30-39).
Spring wheat	0.6	Between tillers erect stage and the second node detectable stage (GS 30-32).
Winter barley	0.6 OR 0.9	At 0.6 litres product/ha Between tillers erect stage and second node detectable stage (GS 30-32). At 0.9 litres product/ha Between flag leaf just visible and flag leaf fully emerged stages (GS 37-39).
Spring barley	0.7	Between tillers erect stage and 2 nd node detectable (GS 30-32).
Winter and spring oats	0.6	Between tillers erect stage and 1 st node detectable (GS 30-31).
Durum wheat, rye and triticale	0.6	Between tillers erect stage and 2 nd node detectable (GS 30-32).
Ryegrass seed crops	1.1	Between tillers erect stage and 1 st node detectable (GS 30-31).

MIXING AND APPLICATION

For use by tractor mounted/trailed sprayers only.

Water volume

Apply STEMPER in 200-400 L/ha water. The higher water volume may be used if the crop is dense and increased penetration of spray is required. Apply as a MEDIUM (BCPC definition) quality spray.

Mixing and spraying

For use by tractor mounted/trailed sprayers only.

Half fill the spray tank with water and begin agitation. Add the required quantity of STEMPER. Add the remainder of the water and continue agitation until spraying is complete.

Spray tank cleaning

Thoroughly wash all spraying equipment and measuring equipment with water and a wetting agent using an integrated pressure rinsing device or manually rinsing three times immediately after use.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorised to and marketed by:

ADAMA Agricultural Solutions UK Ltd

Third Floor East

1410 Arlington Business Park

Theale

Reading

RG7 4SA

Tel: +44 (0)1635 860555

Technical Helpline: +44 (0)1635 876622

www.adama.com

Email: ukenquiries@adama.com

Distributed in Ireland by:

Croplink (Ireland) Ltd

Springhill

Carrigtwohill

Co. Cork

T45 F685

Ireland

Tel: 021 4882806

STEMPER® is a registered trademark of a company of the ADAMA Group.

© ADAMA Agricultural Solutions UK Ltd

Specimen-
2023 to date