

Production date / Batch number: see packaging

24 hour emergency number: 0032 14 58 45 45



008731E_02

PROMINO® 300 EC

Contains: 300 g/L Prothioconazole as emulsifiable concentrate (EC) formulation.

Contents: 5 L

Harmful if swallowed.

Causes skin irritation.

Causes serious eye damage.

Very toxic to aquatic life with long lasting effects.

Wash thoroughly after handling.

Do not eat, drink or smoke when using this product.

Wear protective gloves/ protective clothing/ eye protection/ face protection.

IF SWALLOWED: Call a POISON CENTER/ doctor if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTER or doctor/ physician. Rinse mouth.

If skin irritation occurs: Get medical advice/ attention.

Take off contaminated clothing and wash before reuse. Store locked up.

Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste. Triple rinsed containers should be punctured to prevent re-use.

To avoid risks human health and the environment, comply with the instructions for use.

Authorisation Holder:

CAC Chemical GmbH · Böttgerstrasse 12 · D-20148 Hamburg, Germany

Marketing Company:

Helm Great Britain Ltd · The Harlequin Building · 65 Southwark Street · London · SE1 0HR
Tel: +44 (0) 207 620 0756

PCS No.: 06764



SPECIMEN
2022 to date



DANGER



Open here

SAFETY PRECAUTIONS

Operator protection

Wear protective gloves/ protective clothing/ eye protection/ face protection.

Do not eat, drink or smoke when using this product.

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. To protect aquatic organisms respect an unsprayed buffer zone of 10 m to surface water bodies.

Storage and disposal

DO NOT RE-USE CONTAINER FOR ANY OTHER PURPOSE AND DISPOSE OF SAFELY.

KEEP AWAY FROM FOOD, DRINK, AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

Do not contaminate water with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads)

IMPORTANT INFORMATION

| For use only as an agricultural fungicide. | | | | |
|---|------------------|-------------------------|-----------------|--|
| Crop | Max. single dose | Max.no. of applications | Max. total dose | Latest time of application |
| Barley, Wheat, Durum wheat, Triticale | 0,65 L/ha | – | 1,3 L/ha | Before end of flowering stage (GS69) |
| Rye | 0,65 L/ha | – | 1,3 L/ha | Before beginning of flowering stage (GS61) |
| Oilseed rape | 0,6 L/ha | – | 1,2 L/ha | Before end of flowering (GS69) |
| Method of application: Tractor mounted/trailed sprayer | | | | |
| READ LABEL BEFORE USE: USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE: FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS | | | | |

Resistance strategy

Repeated application of PROMINO® alone should not be used on the same crop against a high-risk pathogen such as cereal powdery mildew. Tank-mixtures or alternation with fungicides having a different mode of action (e.g. morpholines) have been shown to protect against the development of resistant forms of disease.

Take all precautionary measures to reduce the selection pressure for insensitive Septoria strains (e.g. tank-mix with product having a different mode of action which is active against Septoria). Consult your advisor for up to date guidance regarding current resistance status and a strategy for preventing and managing resistance in the cereal and oilseed rape pathogens listed on the label.

The Fungicide Resistance Action Committee (FRAC) produces recommendations that may be consulted for additional information.

Strains of light leaf spot resistant to azole fungicides are known to exist. To avoid development of resistance apply product protectively in response to disease forecasts. Where possible, when light leaf spot is present, avoid the use of azole based fungicides when targeting other diseases such as Sclerotinia at mid flowering.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

PROMINO® is a triazolothione fungicide recommended for control of a wide range of diseases on winter and spring barley, winter and spring wheat, durum wheat, triticale, winter rye, and for disease control in winter oilseed rape.

Diseases controlled: Cereals

| | Wheat | Barley | Spelt | Triticale | Rye |
|--|-------|--------|-------|-----------|-----|
| Eyespot (<i>Oculimacula spp</i>) | R | R | R | R | R |
| Septoria Leaf Blotch (<i>Mycosphaerella graminicola</i>) | MC | – | MC | MC | MC |
| Glume Blotch (<i>Stagonospora nodorum</i>) | MC | – | MC | – | MC |
| Powdery Mildew (<i>Blumeria graminis</i>) | C | C | C | C | C |
| Yellow Rust | C | C | C | C | |
| Brown Rust | MC | C | MC | C | C |
| Crown Rust | – | – | – | – | – |
| Tan Spot | MC | – | MC | – | – |
| Fusarium Ear blight | MC | MC | MC | – | – |
| Leaf Blotch (<i>Rhynchosporium secalis</i>) | – | C | – | C | C |
| Net Blotch (<i>Pyrenophora teres</i>) | – | C | – | – | – |

Disease controlled: Oil seed rape

| | Oil seed rape |
|-----------------------------|---------------|
| Phoma leaf spot/stem canker | C |
| <i>Sclerotinia</i> stem rot | MC |
| Light leaf spot | MC |

C = control, MC = moderate control, R = reduction

Water volume

Apply PROMINO® in 200 – 400 liters of water per hectare. The higher spray volumes are recommended where the crop is dense or disease pressure / risk is high to ensure good penetration to the lower leaves and stem bases. Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve.

Crops

PROMINO may be used on all commercial varieties of winter and spring barley, winter and spring wheat, durum wheat, triticale, winter rye and winter oilseed rape.

Mixing

Half fill the spray tank with clean water and start the re-circulation system. Pour the required quantity of product into the spray tank and continue agitation. Top up the spray tank with water to the required level. Continue agitation until spraying is completed. Spray immediately after mixing.

SPRAY TANK CLEANING

It is important that the spray tank, boom, hoses filters and nozzles are thoroughly washed out to remove all traces of PROMINO® immediately after spraying and before the sprayer is used again:

1. Drain sprayer completely, then wash out tank boom and hoses with clean water. Drain again.
2. Fill the tank with clean water. Flush through boom and hoses then leave sprayer for 10 minutes with agitation on. Then drain completely.
3. Flush the tank, boom and hoses with clean water twice. Ensure any spray contamination on the outside of the sprayer is removed by washing in clean water.

CONDITIONS OF SUPPLY

All goods supplied by us are of a high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or

re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use contrary to label instructions. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

PROMINO® is a registered trademark of CAC.

SPECIMEN -
2022 to date

Production date / Batch number: see packaging

24 hour emergency number: 0032 14 58 45 45



008731E_02

PROMINO® 300 EC

Contains: 300 g/L Prothioconazole as emulsifiable concentrate (EC) formulation.

Contents: 5L

Harmful if swallowed.

Causes skin irritation.

Causes serious eye damage.

Very toxic to aquatic life with long lasting effects.

Wash thoroughly after handling.

Do not eat, drink or smoke when using this product.

Wear protective gloves/ protective clothing/ eye protection/ face protection.

IF SWALLOWED: Call a POISON CENTER/ doctor if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.

Immediately call a POISON CENTER or doctor/ physician.

Rinse mouth.

If skin irritation occurs: Get medical advice/ attention.

Take off contaminated clothing and wash before re-use.

Store locked up.

Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Triple rinsed containers should be punctured to prevent re-use.

To avoid risks human health and the environment, comply with the instructions for use.

Authorisation Holder:

CAC Chemical GmbH · Böttgerstrasse 12 · D-20148 Hamburg, Germany

Marketing Company:

Helm Great Britain Ltd · The Harlequin Building · 65 Southwark Street · London · SE1 0HR
Tel: +44 (0) 207 620 0756

PCS No.: 06764

