

## **Chapter 7**

**List of off label approvals which have been granted up to 1/1/19 for use on food crops**



---

**Off-Label Extension of Authorisation for Minor uses (EAMU) - Food Crops as of 01 March 2019**

**Warning: Off-Label EAMU Approvals:**

*The use of any product in accordance with an off-label EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

*Instruction sheets for off-label EAMU approvals can be obtained from your Teagasc office, or Bord Bia, or local advisor or on the PCS website.*

**Disclaimer**

*The Department of Agriculture Food and the Marine maintains its websites to enhance public access to information about the Department and its activities and to provide services to its customers. Every effort has been made to ensure that the information provided on the Department's websites is accurate and up-to-date. If you notice any errors or omissions please let the Department know as soon as possible. The Department gives no guarantees, undertakings or warranties concerning the accuracy, completeness or up-to-date nature of information provided on its websites and does not accept any liability whatsoever arising from any errors or omissions.*

*The Department reserves the right to change all or any of the information provided at any time, but shall not be responsible for or liable in respect of any such change in information. The Department has no liability for any loss occasioned from the use of information appearing on these websites. Any links from these websites to external websites are provided as a matter of convenience only, and should not be taken as an endorsement of the contents or practices of those external websites.*

*Unless otherwise stated reproduction is authorised except for commercial purposes, provided the source is acknowledged.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Apples</i>											
	Cerone	03933	Bayer CropScience Limited	Growth Regulator			0.75 l/ha	0.75 l/ha	-	10 days before harvest	10 days
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide	<i>Ethephon</i>	<i>480 g/l</i>	2.0 kg/ha	8.0 kg/ha	-	Before GS 07	
	Fathom	04688	FMC Agro Ltd	Fungicide	<i>Copper oxychloride</i>	<i>87.8% w/w</i>	0.437 l/ha	0.437 l/ha	1	GS 20	
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide	<i>Tebuconazole</i>	<i>200 g/l</i>	0.25 l/ha	1.0 l/ha	-	28 days before harvest	28 days
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide	<i>Tebuconazole</i>	<i>200 g/l</i>	0.25 l/ha	1.0 l/ha		28 days before harvest	28 days
	Switch	03761	Syngenta UK Ltd	Fungicide	<i>Difenoconazole</i>	<i>250 g/l</i>	0.8 kg/ha	2.4 kg/ha	3	3 days before harvest	
	Talius	04604	DuPont (UK) Ltd	Fungicide	<i>Cyprodinil</i>	<i>37.5 % w/w</i>	0.25 l/ha	0.5 l/ha	2 per year	49 days before harvest	49 days
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Proquinazid</i>	<i>200 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Asparagus</i>											
	Amistar Opti	05068	Syngenta UK Ltd	Fungicide			2.5 l/ha	2.5 l/ha	-	Before senescence	
					<i>Chlorothalonil</i>	<i>400 g/l</i>					
					<i>Azoxystrobin</i>	<i>80 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	Before senescence	
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	Before senescence	
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	Before senescence	
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	Before senescence	
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Lentagran WP	03082	Belchim Crop Protection NV	Herbicide			2.0 kg/ha	4.0 kg/ha	2 - 1 to growing crop. 1 - post harvest underneath	28 days before harvest	
					<i>Pyridate</i>	<i>45 % w/w</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	After harvest and before end of September	
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	PDM 330 EC	03582	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>330 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		After harvest and before end of September	
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Sencorex Flow	04361	Bayer CropScience Limited	Herbicide			0.9 l/ha	0.9 l/ha	-	Before end of dormancy	
					<i>Metribuzin</i>	<i>600 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Aubergines - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.45 l/ha	0.9 l/ha	2 per crop	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	FLIPPER	05027	Alpha Biopesticides Ltd	Insecticide			16.0 l/ha	80.0 l/ha	5	N/A	
					<i>Fatty Acid - C7-C20</i>	<i>479.8 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Microthiol Special	90548	UPL Europe Ltd	Fungicide			0.01kgs/5 litres water	-	-	-	0 days
					<i>Sulphur</i>	<i>80 %w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	2.0 l/ha	4 per crop	14 days before harvest	14 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.06 kgs/100 litres water	0.18 kgs/ha	3 / crop	3 days before harvest	3 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product</i>					<i>Max Ind.</i>	<i>Max. Total</i>	<i>Max No of</i>	<i>Latest Time</i>	
<i>Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Dose:</i>	<i>Dose:</i>	<i>Applic's:</i>	<i>of Applic:</i>	<i>PHI:</i>
<i>Aubergines - protected</i>										
Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3.0 l/ha	12.0 l/ha	-	3 days before harvest	
				<i>Propamocarb</i>	<i>530 g/l</i>					
				<i>Propamocarb</i>	<i>530 g/l</i>					
				<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
				<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
Revus	05154	Syngenta UK Ltd	Fungicide			0.6 l/ha	2.4 l/ha	4 per crop	3 days before harvest	3 days
				<i>Mandipropamid</i>	<i>250 g/l</i>					
Scala	03916	BASF plc	Fungicide			0.5 l/ha	1.0 l/ha	2 per crop	3 days before harvest	3 days
				<i>Pyrimethanil</i>	<i>400 g/l</i>					
Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	4.5 kgs/ha	3	1 day before harvest	1 day
				<i>Fenhexamid</i>	<i>50 %w/w</i>					
Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Baby leaf production</i>											
	Acrobat MZ	04059	BASF plc	Fungicide			2.0 kg/ha	6.0 kg/ha	-	21 days before harvest	21 days
					<i>Dimethomorph</i>	<i>75 g/kg</i>					
					<i>Mancozeb</i>	<i>667 g/kg</i>					
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Bonalan	05763	Gowan Crop Protection Ltd	Herbicide			8.0 l/ha	8.0 l/ha	-	Incorporation pre sowing/planting	
					<i>Benfluralin</i>	<i>180 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	<i>220 g/kg</i>					
					<i>GHA / ATCC 74040</i>						
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.25 l/ha	1 per crop	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	CIPC	03066	Arysta LifeScience Benelux SPRL	Herbicide			1.9 l/ha	3.8 l/ha	2 per crop	2 weeks after planting out	
					<i>Chlorpropham</i>	<i>400 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha*	1.26 l/ha*	3 per crop*	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Devrinol	05234	UPL Europe Ltd	Herbicide			0.85 l/ha	0.85 l/ha	1 per crop	Pre-planting	
					<i>Napropamide</i>	<i>450 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Baby leaf production</i>											
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide	<i>Mancozeb</i>	<i>75% w/w</i>	75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	250 g/ha	500 g/ha	2	3 days pre harvest	3 days
	Globaztar SC	05529	Globachem nv	Fungicide	<i>Acetamiprid</i>	<i>20 % w/w</i>	1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
	Infinito	04309	Bayer CropScience Limited	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.6 l/ha	1.6 l/ha	1 per crop	Before GS 49	14 days
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide	<i>Propamocarb</i>	<i>625 g/l</i>	75 mls/ha	150 mls/ha	4	7 days pre harvest	7 days
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide	<i>Fluopicolide</i>	<i>62.5 g/l</i>	75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
	Kerb Flo	02830	Dow AgroSciences Ltd	Herbicide	<i>Propamocarb</i>	<i>625 g/l</i>	1.875 l/ha	1.875 l/ha	1	24 days before harvest	24 days
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide	<i>Fluopicolide</i>	<i>62.5 g/l</i>	75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	1.0 kg/ha	3.0 kg/ha	-	N/A	
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide	<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>	75 mls/ha	150 mls/ha	4	7 days before harvest	7 days
	Markate 50	05001	Globachem nv	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Baby leaf production</i>											
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide	<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>	0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
	Movento	05349	Bayer CropScience Limited	Insecticide	<i>Spirotetramat</i>	<i>150 g/l</i>	0.5 l/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 mls/ha	150 mls/ha	4	7 days before harvest	7 days
	Paraat	03614	BASF plc	Fungicide	<i>Dimethomorph</i>	<i>500 g/kg</i>	0.36 kg/ha	0.72 kg/ha	2 per crop	14 days before harvest	14 days
	PDM 330 EC	03582	BASF plc	Herbicide	<i>Pendimethalin</i>	<i>330 g/l</i>	4.5 l/ha	4.5 l/ha	1	6 weeks pre harvest	
	Pendifin 400 SC	05731	Finchimica S.p.A	Herbicide	<i>Pendimethalin</i>	<i>400 g/l</i>	3.75 l/ha	3.75 l/ha	1 per crop	6 weeks before harvest	6 weeks
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide	<i>Pymetrozine</i>	<i>50 %w/w</i>	0.4 kgs/ha	1.2 kgs / ha	3	7 days	
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide	<i>Propamocarb</i>	<i>530 g/l</i>	2.5 l/ha	5.0 l/ha	-	21 days before harvest	21 days
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide	<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>	75 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide	<i>Mancozeb</i>	<i>64 %w/w</i>	1.9 kg/ha	5.7 kg/ha*		35 days before harvest	35 days
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	8 l/ha	48 l/ha	6		0 days

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Baby leaf production</i>											
	Signum	03799	BASF plc	Fungicide			1.5 kgs/ha	3.0 kgs/ha	-	14 days before harvest	
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Stomp 400 SC	02879	BASF plc	Herbicide			3.75 l/ha	3.75 l/ha	1 / crop	6 weeks pre harvest	6 weeks
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1 / crop	6 weeks pre harvest	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			0.7 kg/ha	1.4 kg/ha	2	7 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha	2	3 days before harvest	3 days
					<i>Fenhexamid</i>	<i>50 %w/w</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha	600 ml/ha	3 per crop	3 days before harvest	3 days
					<i>Spinosad</i>	<i>480 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Basil</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
					<i>Cycloxydim</i>	100 g/l					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	50 %w/w					
<i>Beans - broad</i>											
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	125 g/l					
					<i>Azoxystrobin</i>	200 g/l					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			1.0 l/ha	1.0 l/ha	-	30 days before harvest	30 days
					<i>Clethodim</i>	120 g/l					
	Claw 100	05874	DHM Agrochemicals Ltd.,	Herbicide			1.5 l/ha	1.5 l/ha	1 per crop	Before flower buds visible stage	7 weeks
					<i>Propaquizafop</i>	100 g/l					
	Farmco Zealot	05299	Farmco Agritrading Ltd	Herbicide			1.5 l/ha	1.5 l/ha	1 per crop	Before flower buds visible stage	7 weeks
					<i>Propaquizafop</i>	100 g/l					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	375 g/kg					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l					
	Tepeki	02683	ISK Biosciences Europe N.V.	Insecticide			0.14 kg/ha	0.14 kg/ha	1 per crop	14 days before harvest	14 days
					<i>Flonicamid</i>	500 g/kg					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beans - field</i>											
Anthem	04761	Adama Agricultural Solutions UK Ltd	Herbicide				3.3 l/ha	3.3 l/ha		Pre crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide				2.1 kg/ha	2.1 kg/ha	1 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
Biscaya	03602	Bayer CropScience Limited	Insecticide				0.4 l/ha	0.8 l/ha	2 per crop	7 days before harvest	7 days
					<i>Thiacloprid</i>	<i>240 g/l</i>					
Defy	04277	Syngenta UK Ltd	Herbicide				5.0 l/ha	5.0 l/ha	1 per crop	Pre-emergence	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
Dithane 945	03705	Indofil Industries (Netherlands) B.V	Fungicide				2.0 kg/ha	2.0 kg/ha	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>80 % w/w</i>					
Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide				2.1 kg/ha	2.1 kg/ha	1 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
PDM 330 EC	03582	BASF plc	Herbicide				4.0 l/ha	4.0 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>330 g/l</i>					
Pendifin 400 SC	05731	Finchimica S.p.A	Herbicide				3.3 l/ha	3.3 l/ha	1 per crop	Pre-crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide				2.0 kg/ha	2.0 kg/ha	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>64 %w/w</i>					
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
Roxy 800 EC	05739	Globachem nv	Herbicide				5.0 l/ha	5.0 l/ha	1 per crop	Pre-emergence	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide				8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
Stomp 400 SC	02879	BASF plc	Herbicide				3.3 l/ha	3.3 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
Stomp Aqua	03750	BASF plc	Herbicide				2.9 l/ha	2.9 l/ha	1 / crop	Pre crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beans - field</i>											
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
	Wakil XL	05471	Syngenta UK Ltd	Seed Dressing	<i>Esfenvalerate</i>	<i>25 g/l</i>					
					<i>Cymoxanil</i>	<i>100 g/kg</i>	2.0 kg/tonne	2.0 kg/tonne	1 per batch	Before drilling	
					<i>Fludioxonil</i>	<i>50 g/kg</i>					
					<i>Metalaxyl M</i>	<i>175 g/kg</i>					
<i>Beans - french</i>											
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			1.0 l/ha	1.0 l/ha	-	30 days before harvest	30 days
	Claw 100	05874	DHM Agrochemicals Ltd.,	Herbicide	<i>Clethodim</i>	<i>120 g/l</i>	1.5 l/ha	1.5 l/ha	1 per crop	Before flower buds visible stage	7 weeks
	Dual Gold	04192	Syngenta UK Ltd	Herbicide	<i>Propaquizafop</i>	<i>100 g/l</i>	1.0 l/ha	1.0 l/ha	-	Immediately after sowing	
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide	<i>S-Metolachlor</i>	<i>960 g/l</i>	1.0 l/ha	1.0 l/ha	-	Immediately after sowing	
	Farmco Zealot	05299	Farmco Agritrading Ltd	Herbicide	<i>S-Metolachlor</i>	<i>960 g/l</i>	1.5 l/ha	1.5 l/ha	1 per crop	Before flower buds visible stage	7 weeks
	Fathom	04688	FMC Agro Ltd	Fungicide	<i>Propaquizafop</i>	<i>100 g/l</i>	0.95 l/ha	0.95 l/ha	1	GS 20	
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Tebuconazole</i>	<i>200 g/l</i>	8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
<i>Beans - navy</i>											
	Claw 100	05874	DHM Agrochemicals Ltd.,	Herbicide	<i>Propaquizafop</i>	<i>100 g/l</i>	1.5 l/ha	1.5 l/ha	1 per crop	Before flower buds visible stage	7 weeks
	Farmco Zealot	05299	Farmco Agritrading Ltd	Herbicide	<i>Propaquizafop</i>	<i>100 g/l</i>	1.5 l/ha	1.5 l/ha	1 per crop	Before flower buds visible stage	7 weeks

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beans - runner</i>											
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			1.0 l/ha	1.0 l/ha	-	30 days before harvest	30 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.0 l/ha	1.0 l/ha	-	Immediately after sowing	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.0 l/ha	1.0 l/ha	-	Immediately after sowing	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Fathom	04688	FMC Agro Ltd	Fungicide			0.95 l/ha	0.95 l/ha	1	GS 20	
					<i>Tebuconazole</i>	<i>200 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beet - fodder</i>											
	Deca 2.5% EC	04952	Sharda Europe b.v.b.a	Insecticide			300 mls/ha	300 mls/ha	-	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	300 mls/ha		30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	500 mls/ha	-	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Fortress	02364	Dow AgroSciences Ltd	Fungicide			0.3 l/ha	0.4 l/ha	2	28 days before harvest	28 days
					<i>Quinoxifen</i>	<i>500 g/l</i>					
	GAT Decline 2.5 E	04422	FMC Chemical s.p.r.l.	Insecticide			300 mls/ha	300 mls/ha	-	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha	-	28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	NEMGuard DE	05368	ECOspray Ltd	Nematicide			10 kg/ha	10 kg/ha	1	At drilling	
					<i>Garlic Extract</i>	<i>450 g/kg</i>					
	Polux	05946	Galenika-Fitofarmacija	Insecticide			0.3 l/ha	0.3 l/ha	-	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha		28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beet - red</i>											
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	NEMGuard DE	05368	ECOSpray Ltd	Nematicide			10 kg/ha	10 kg/ha	1	At drilling	
					<i>Garlic Extract</i>	<i>450 g/kg</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
<i>Beet - sugar</i>											
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha	-	28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha		28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beetroot</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>240 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			1.0 l/ha	1.0 l/ha	-	Before row closure	56 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Escolta	05681	Bayer CropScience Limited	Fungicide			0.35 l/ha	0.7 l/ha	2 per year	21 days before harvest	21 days
					<i>Trifloxystrobin</i>	<i>375 g/l</i>					
					<i>Cyproconazole</i>	<i>160 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beetroot</i>											
	Fortress	02364	Dow AgroSciences Ltd	Fungicide			0.3 l/ha	0.4 l/ha	2	28 days pre harvest	28 days
					<i>Quinoxifen</i>	<i>500 g/l</i>					
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	8 weeks before harvest	56 days
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha	-	28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Microthiol Special	90548	UPL Europe Ltd	Fungicide			10 kg/ha	20 kgs/ha	2	Last application before September	-
					<i>Sulphur</i>	<i>80 %w/w</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Beetroot</i>											
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha		28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	8 weeks before harvest	8 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
	Venzar 500 SC	04805	DuPont (UK) Ltd	Herbicide			0.4 l/ha	0.4 l/ha	-	Pre-emergence	
					<i>Lenacil</i>	<i>500 g/l</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blackberries - field</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
	Amistar Top	04582	Syngenta UK Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	Apollo 50SC	04610	Adama Agricultural Solutions UK Ltd	Acaricide			0.4 l/ha	0.4 l/ha	1 per year	9 days before harvest	9 days
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide	<i>Clofentezine</i>	<i>500 g/l</i>	0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 mls/ha	2.49 l/ha	3	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	Before flowering or after harvest	
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 g/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kgs/ha	1.2 kgs/ha	3	28 days before harvest	28 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
	Scala	03916	BASF plc	Fungicide			2.0 l/ha	4.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha	-	21 days before harvest	
					<i>Carfentrazone-ethyl</i>	<i>60 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blackberries - field</i>											
Switch	03761	Syngenta UK Ltd	Fungicide				1.0 kg/ha	3.0 kg/ha	3	10 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
Tracer	02649	Dow AgroSciences Ltd	Insecticide				200 ml/ha in 1000L water	400 ml/ha	2 per year	1 day before harvest	1 day
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blackberries - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Apollo 50SC	04610	Adama Agricultural Solutions UK Ltd	Acaricide			0.4 l/ha	0.4 l/ha	1 per year	9 days before harvest	9 days
					<i>Clofentezine</i>	<i>500 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.75 l/ha	3 per crop	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 g/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
	Scala	03916	BASF plc	Fungicide			2.0 l/ha	4.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha	-	21 days before harvest	21 days
					<i>Carfentrazone-ethyl</i>	<i>60 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	14 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blackberries - protected</i>											
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha in 1000L water	600 ml/ha	3 per year	1 day before harvest	1 day
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blackcurrants - field</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.75 l/ha	3 per year	3 days before harvest	3 days
					<i>Thiacloprid</i>	480 g/l					
	Chikara Weed Cont	04203	ISK Biosciences Europe N.V.	Herbicide			120 g/ha	120 g/ha	1 per year	Before bud burst	
					<i>Flazasulfuron</i>	25 % w/w					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			400 ml/ha	800 ml/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	15 g/l					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			100 mls/ha	200 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	100 g/l					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	375 g/kg					
	Manfil 75 WG	03598	Indofil Industries Limited	Fungicide			1.5 kg per 1000 litres water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	75 % w/w					
	Manfil 80 WP	03563	Indofil Industries Limited	Fungicide			1.4 kg per 1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	800 g/kg					
	Masai	03585	BASF plc	Insecticide			1.5 kgs/ha	1.5 kgs/ha	-	7 days before harvest	
					<i>Tebufenpyrad</i>	200 g/kg					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			100 mls/ha	200 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	100 g/l					
	Penncozeb 80 WP	03609	UPL Europe Ltd	Fungicide			1.4 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	800 g/kg					
	Penncozeb WDG	03608	UPL Europe Ltd	Fungicide			1.5 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	750 g/kg					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blackcurrants - field</i>											
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kgs/ha	1.2 kgs/ha	3	28 days before harvest	28 days
	Scala	03916	BASF plc	Fungicide	<i>Pymetrozine</i>	<i>50 %w/w</i>	2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Pyrimethanil</i>	<i>400 g/l</i>	8 l/ha	48 l/ha	6		0 days
	Tracer	02649	Dow AgroSciences Ltd	Insecticide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	200 ml/ha in 250L water	400 ml/ha	2 per year	7 days before harvest	7 days
					<i>Spinosad</i>	<i>480 g/l</i>					
<i>Blackcurrants - protected</i>											
	AQ 10	05801	CBC (Europe) S.r.l.	Fungicide	<i>Ampelomyces Quisqualis</i>	<i>58 %w/w</i>	70 g/ha	-	12	-	
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide	<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>	0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
	Scala	03916	BASF plc	Fungicide	<i>Pyrimethanil</i>	<i>400 g/l</i>	2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	8 l/ha	48 l/ha	6		0 days
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide	<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>	0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide	<i>Spinosad</i>	<i>480 g/l</i>	200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blueberries - field</i>											
Batavia	05905	Bayer CropScience Limited	Insecticide				0.75 l/ha	1.5 l/ha	2	Post harvest application	
					<i>Spirotetramat</i>	<i>100 g/l</i>					
Botanigard WP	06264	Mycotech Europe Ltd	Insecticide				0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
CaLypso	02441	Bayer CropScience Limited	Insecticide				0.25 l/ha	0.75 l/ha	3 per year	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
Chikara Weed Cont	04203	ISK Biosciences Europe N.V.	Herbicide				120 g/ha	120 g/ha	1 per year	Before bud burst	
					<i>Flazasulfuron</i>	<i>25 % w/w</i>					
Fusilade Max	01472	Syngenta UK Ltd	Herbicide				3.0 l/ha	3.0 l/ha	1	Before flowering or after harvest	
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide				1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
Manfil 75 WG	03598	Indofil Industries Limited	Fungicide				1.5 kg per 1000 litres water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75 % w/w</i>					
Manfil 80 WP	03563	Indofil Industries Limited	Fungicide				1.4 kg per 1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
Masai	03585	BASF plc	Insecticide				1.5 kgs/ha	1.5 kgs/ha	-	7 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
Penncozeb 80 WP	03609	UPL Europe Ltd	Fungicide				1.4 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
Penncozeb WDG	03608	UPL Europe Ltd	Fungicide				1.5 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>750 g/kg</i>					
Scala	03916	BASF plc	Fungicide				2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blueberries - field</i>											
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.5 kgs/ha	4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest)	-	3 days before harvest, N/A for post harvest	
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Blueberries - protected</i>											
AQ 10	05801	CBC (Europe) S.r.l.	Fungicide		<i>Ampelomyces Quisqualis</i>	58 %w/w	70 g/ha	-	12	-	
Botanigard WP	06264	Mycotech Europe Ltd	Insecticide		<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg	0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide		<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	375 g/kg	1.0 kg/ha	3.0 kg/ha	-	N/A	
Serenade ASO	03847	Bayer CropScience Limited	Fungicide		<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l	8 l/ha	48 l/ha	6		0 days
Tracer	02649	Dow AgroSciences Ltd	Insecticide		<i>Spinosad</i>	480 g/l	200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Broccoli</i>											
	Bigalo	06250	Globachem nv	Fungicide			1.0 kg/ha	3.0 kg/ha	-	14 days before harvest	14 days
					<i>Pyraclostrobin</i>	<i>6.7% w/w</i>					
					<i>Boscalid</i>	<i>26.7% w/w</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Gamit 36 CS	05357	FMC Chemical s.p.l.	Herbicide			0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
					<i>Clomazone</i>	<i>360 g/l</i>					
	Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	3.2 l/ha	2 per crop*	Before GS 49	14 days
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Lentagran WP	03082	Belchim Crop Protection NV	Herbicide			2.0 kg/ha	2.0 kg/ha	1	42 days before harvest	
					<i>Pyridate</i>	<i>45 % w/w</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide			2.0 kgs/ha	4.0 kgs/ha		30 days pre harvest	30 days
					<i>Mancozeb</i>	<i>64 %w/w</i>					
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Broccoli</i>											
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide	<i>Esfenvalerate</i>	<i>25 g/l</i>	0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	7 days after transplanting	
					<i>Pendimethalin</i>	<i>250g/l</i>					
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Brussel sprouts</i>											
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	6.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	28 days before harvest	28 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide			0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
					<i>Clomazone</i>	<i>360 g/l</i>					
	Infinito	02532	Bayer CropScience Limited	Fungicide			1.6 l/ha	3.2 l/ha	2 per crop*	Before GS 49	14 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Insyst	03249	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	250 g/ha	1	21 days before harvest	21 days
					<i>Acetamiprid</i>	<i>20 %w/w</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	6.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Brussel sprouts</i>											
	Steward	02603	DuPont (UK) Ltd	Insecticide			85 g/ha	255 g/ha	3 / crop	1 day before harvest	
					<i>Indoxacarb</i>	<i>300 g/kg</i>					
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
					<i>Esfenvalerate</i>	<i>25 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	7 days after transplanting	
					<i>Pendimethalin</i>	<i>250g/l</i>					
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					
<i>Bush/Cane fruit plantations</i>											
	Digital	03147	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha		365 days before first harvest	
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Fastnet	06253	Sipcam U.K. Ltd	Herbicide			3.75 l/ha	3.75 l/ha	1 per year	N/A	
					<i>Pendimethalin</i>	<i>365 g/l</i>					
	Guillotine	03146	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	365 days before first harvest	
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Most Micro	04837	Sipcam U.K. Ltd	Herbicide			3.75 l/ha	3.75 l/ha	1 per year	N/A	
					<i>Pendimethalin</i>	<i>365 g/l</i>					
	Roundup Biactive	02313	Monsanto UK Ltd	Herbicide			4 l/ha	10 l/ha		28 days pre harvest	28 days
					<i>Glyphosate</i>	<i>360 g/l</i>					
	Sumimax	02764	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	365 days before first harvest	
					<i>Flumioxazin</i>	<i>300 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cabbage - chinese</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	7 days before harvest	7 days
					<i>Thiacloprid</i>	<i>240 g/l</i>					
	Butisan S	04631	BASF plc	Herbicide			1.0 l/ha	1.0 l/ha	-	Post emergence GS 16-18. 6-8 days after planting	
					<i>Metazachlor</i>	<i>500 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Devrinol	05234	UPL Europe Ltd	Herbicide			2.8 l/ha	2.8 l/ha	-	Before drilling or transplanting	
					<i>Napropamide</i>	<i>450 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide			0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
					<i>Clomazone</i>	<i>360 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cabbage - chinese</i>											
	Globazar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	21 days before harvest	21 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	1.0 l/ha	2 per crop	3 days before harvest	3 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kgs/ha	1.2 kgs/ha	3	14 days before harvest	14 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		21 days before harvest	21 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
					<i>Esfenvalerate</i>	<i>25 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m <sup>2</sup> *	0.5 g/m <sup>2</sup>	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cabbage - head</i>											
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	28 days before harvest	28 days
	Dual Gold	04192	Syngenta UK Ltd	Herbicide	<i>Clethodim</i>	<i>120 g/l</i>	1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide	<i>S-Metolachlor</i>	<i>960 g/l</i>	1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
	Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide	<i>S-Metolachlor</i>	<i>960 g/l</i>	0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
	Infito	04309	Bayer CropScience Limited	Fungicide	<i>Clomazone</i>	<i>360 g/l</i>	1.6 l/ha	3.2 l/ha	2 per crop*	Before GS 49	14 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide	<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>	0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide	<i>Mancozeb</i>	<i>64 %w/w</i>	1.5 kgs/ha	4.5 kgs/ha		14 days pre harvest	14 days
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	8 l/ha	48 l/ha	6		0 days
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide	<i>Esfenvalerate</i>	<i>25 g/l</i>	165 mls/ha	330 mls/ha	2	7 days pre harvest	

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cabbage - head</i>											
T34	Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
Teppeki		02683	ISK Biosciences Europe N.V.	Insecticide			0.14 kg/ha	0.28 kg/ha	2 per crop	14 days before harvest	14 days
					<i>Flonicamid</i>	<i>500 g/kg</i>					
Wing P		03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	Before planting	
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					
					<i>Pendimethalin</i>	<i>250g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Calabrese</i>											
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide	<i>S-Metolachlor</i>	<i>960 g/l</i>	1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
	Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide	<i>S-Metolachlor</i>	<i>960 g/l</i>	0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
	Infinito	02532	Bayer CropScience Limited	Fungicide	<i>Clomazone</i>	<i>360 g/l</i>	1.6 l/ha	3.2 l/ha	2 per crop*	Before GS 49	14 days
	Lentagran WP	03082	Belchim Crop Protection NV	Herbicide	<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide	<i>Pyridate</i>	<i>45 % w/w</i>	0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST</i>	<i>2% w/w</i>					
					<i>BIPESCO 5/F52</i>						
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide			2.0 kgs/ha	4.0 kgs/ha		30 days pre harvest	30 days
					<i>Mancozeb</i>	<i>64 %w/w</i>					
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
					<i>Esfenvalerate</i>	<i>25 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Calabrese</i>											
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	7 days after transplanting	
					<i>Pendimethalin</i>	<i>250g/l</i>					
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Carrots</i>											
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Corbel	90438	BASF plc	Fungicide			1.0 l/ha	3.0 l/ha	3	28 days before harvest	28 days
					<i>Fenpropimorph</i>	<i>750 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Digital	03147	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha		28 days before harvest	28 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Emerger	06190	Bayer CropScience Limited	Herbicide			2.5 l/ha	2.5 l/ha	-	GS 12	
					<i>Aclonifen</i>	<i>600 g/l</i>					
	Fazor Star	03729	Arysta LifeScience Great Britain Ltd	Growth Regulator			5.0 kg/ha	5.0 kg/ha	1 per crop	21 days before harvest	
					<i>Maleic hydrazide</i>	<i>600 g/kg</i>					
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide			0.15 l/ha	0.15 l/ha	1 per year	Pre-emergence	
					<i>Isoxaben</i>	<i>500 g/l</i>					
	Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide			0.083l/ha	0.25 l/ha*		Before 4th trye leaf stage & 55 days before harvest	55 days
					<i>Clomazone</i>	<i>360 g/l</i>					
	Guillotine	03146	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	28 days before harvest	28 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Carrots</i>											
	Hurricane SC	04357	Adama Agricultural Solutions UK Ltd	Herbicide			0.24 l/ha	0.24 l/ha	-	Pre-emergence	
					<i>Diflufenican</i>	<i>500 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	40 days before harvest	40 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Sencorex Flow	04361	Bayer CropScience Limited	Herbicide			0.255 l/ha	0.51 l/ha	3	Before GS 16	28 days
					<i>Metribuzin</i>	<i>600 g/l</i>					
	Sumimax	02764	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	28 days before harvest	28 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cauliflowers</i>											
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Gamit 36 CS	05357	FMC Chemical s.p.l.	Herbicide			0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
					<i>Clomazone</i>	<i>360 g/l</i>					
	Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	3.2 l/ha	2 per crop*	Before GS 49	14 days
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Lentagran WP	03082	Belchim Crop Protection NV	Herbicide			2.0 kg/ha	2.0 kg/ha	1	42 days before harvest	
					<i>Pyridate</i>	<i>45 % w/w</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide			2.0 kgs/ha	4.0 kgs/ha		30 days pre harvest	30 days
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
					<i>Mancozeb</i>	<i>64 %w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cauliflowers</i>										
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
				<i>Esfenvalerate</i>	<i>25 g/l</i>					
T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
				<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
				<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	7 days after transplanting	
				<i>Pendimethalin</i>	<i>250g/l</i>					
				<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Celeriac</i>										
Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>200 g/l</i>					
				<i>Difenoconazole</i>	<i>125 g/l</i>					
AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
				<i>Mancozeb</i>	<i>75% w/w</i>					
Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	7 days before harvest	7 days
				<i>Thiacloprid</i>	<i>240 g/l</i>					
Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
				<i>Clethodim</i>	<i>120 g/l</i>					
Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	14 days pre harvest	14 days
				<i>Acetamiprid</i>	<i>20% w/w</i>					
Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
				<i>Deltamethrin</i>	<i>25 g/l</i>					
Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
				<i>Deltamethrin</i>	<i>15 g/l</i>					
Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	
				<i>Prosulfocarb</i>	<i>800 g/l</i>					
Devrinol	05234	UPL Europe Ltd	Herbicide			1.25 l/ha	1.25 l/ha	-	Pre crop emergence	
				<i>Napropamide</i>	<i>450 g/l</i>					
Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
				<i>Mancozeb</i>	<i>75% w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Celeriac</i>											
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	8 weeks before harvest	56 days
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2 per crop	14 days before harvest	14 days
					<i>Acetamiprid</i>	<i>20 % w/w</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.3 kg/ha	0.6 kg/ha	* 2 per crop	21 days before harvest	21 days
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
					<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kg/ha	1.2 kg/ha	3 per crop	14 days before harvest	14 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Celeriac</i>											
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Sencorex Flow	04361	Bayer CropScience Limited	Herbicide			0.255 l/ha	0.51 l/ha	3	Before GS 16	28 days
					<i>Metribuzin</i>	<i>600 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.0 kg/ha	2.0 kg/ha	2 per crop	14 days before harvest	14 days
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	8 weeks before harvest	8 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Venzar 500 SC	04805	DuPont (UK) Ltd	Herbicide			0.4 l/ha	0.4 l/ha	-	Pre-emergence	
					<i>Lenacil</i>	<i>500 g/l</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Celery - field</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	1 per crop	Before GS11	50 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			50 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			50 mls/ha	150 mls/ha	4	7 days pre harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			50 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			50 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			50 mls/ha	150 mls/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			50 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Celery - field</i>										
Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.3 kg/ha	0.9 kg/ha	* 3 per crop	21 days before harvest	21 days
				<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
				<i>Tebuconazole</i>	<i>500 g/kg</i>					
Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			50 mls/ha	150 mls/ha	4	7 days before harvest	7 days
				<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
PDM 330 EC	03582	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1	Before crop emergence	
				<i>Pendimethalin</i>	<i>330 g/l</i>					
Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kg/ha	1.2 kg/ha	3 per crop	14 days before harvest	14 days
				<i>Pymetrozine</i>	<i>50 %w/w</i>					
Ravane	05536	Globachem nv	Insecticide			50 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
				<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	50 days
				<i>Prosulfocarb</i>	<i>800 g/l</i>					
Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
				<i>Difenoconazole</i>	<i>250 g/l</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
Sparviero	04469	Sipcam Oxon SpA	Insecticide			50 ml/ha	150 ml/ha	4	7 days before harvest	7 days
				<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
				<i>Pendimethalin</i>	<i>455 g/l</i>					
Switch	03761	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
				<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
				<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Celery - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6	-	0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cherries</i>											
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.0 kg/ha	8.0 kg/ha	4 per year	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.312 l/ha	0.625 l/ha	2 per year	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide			2.0 kg/ha	8.0 kg/ha	-	Before GS 09	
					<i>Copper oxychloride</i>	<i>87.8% w/w</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.0 kg/ha	8.0 kg/ha	4 per year	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			0.75 kg/ha	2.25 kg/ha		7 days before harvest	7 days
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
	Systhane 20 EW	04944	Dow AgroSciences Ltd	Fungicide			0.45 l/ha	0.9 l/ha	2	14 days before harvest	14 days
					<i>Myclobutanil</i>	<i>200 g/l</i>					
<i>Chicory</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide			0.2 l/ha	0.2 l/ha	-	Before GS 16	
					<i>Isoxaben</i>	<i>500 g/l</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	1.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Chives</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.2 l/ha	0.6 l/ha	3 per year	14 days before harvest	14 days
					<i>Thiacloprid</i>	480 g/l					
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide			1.5 kgs/ha	4.5 kgs/ha		14 days pre harvest	14 days
					<i>Mancozeb</i>	64 %w/w					
					<i>Metalaxyl M</i>	4.0 %w/w					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
					<i>Cycloxydim</i>	100 g/l					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	50 %w/w					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha	800 ml/ha	4 per crop	7 days before harvest	7 days
					<i>Spinosad</i>	480 g/l					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Collards</i>										
Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	21 days before harvest	21 days
				<i>Thiacloprid</i>	<i>240 g/l</i>					
Devrinol	05234	UPL Europe Ltd	Herbicide			2.8 l/ha	2.8 l/ha	-	Before drilling or transplanting	
				<i>Napropamide</i>	<i>450 g/l</i>					
Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
				<i>S-Metolachlor</i>	<i>960 g/l</i>					
Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
				<i>S-Metolachlor</i>	<i>960 g/l</i>					
Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide			0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
				<i>Clomazone</i>	<i>360 g/l</i>					
Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	3.2 l/ha	2 per crop*	Before GS 49	14 days
				<i>Fluopicolide</i>	<i>62.5 g/l</i>					
				<i>Fluopicolide</i>	<i>62.5 g/l</i>					
				<i>Propamocarb</i>	<i>625 g/l</i>					
				<i>Propamocarb</i>	<i>625 g/l</i>					
Lentagran WP	03082	Belchim Crop Protection NV	Herbicide			2.0 kg/ha	2.0 kg/ha	1	42 days before harvest	
				<i>Pyridate</i>	<i>45 % w/w</i>					
Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
				<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
				<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Collards</i>										
T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
				<i>Trichoderma</i>	<i>12% w/w</i>					
				<i>Asperellum (strain T34)</i>						
				<i>Trichoderma</i>	<i>12% w/w</i>					
				<i>Asperellum (strain T34)</i>						
<i>Coriander</i>										
Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
				<i>Beauveria bassiana</i>	<i>220 g/kg</i>					
				<i>GHA / ATCC 74040</i>						
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus</i>	<i>1015.1 g/l</i>					
				<i>amyloliquefaciens str.</i>						
				<i>QST 713</i>						
Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
				<i>Fenhexamid</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Courgettes - field</i>											
Amistar	05072	Syngenta UK Ltd	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
Amistar Top	04582	Syngenta UK Ltd	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
Azoxystar	05534	Life Scientific Limited	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
Chamane	05002	UPL Europe Ltd	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide				2.1 kg/ha	8.4 kg/ha	4 per crop	3 days before harvest	3 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide				0.25 l/ha	0.25 l/ha	-	Before 2 true leaf stage	
					<i>Clomazone</i>	<i>360 g/l</i>					
Globaztar SC	05529	Globachem nv	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
Signum	03799	BASF plc	Fungicide				1.5 kg/ha	3.0 kg/ha	2 per crop	1 day before harvest	1 day
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Courgettes - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Amistar Top	04582	Syngenta UK Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide	<i>Difenoconazole</i> <i>Azoxystrobin</i>	<i>125 g/l</i> <i>200 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Azoxystar	05534	Life Scientific Limited	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	CaLypso	02441	Bayer CropScience Limited	Insecticide	<i>Thiacloprid</i>	<i>480 g/l</i>	0.45 l/ha	0.9 l/ha	2 per crop	3 days before harvest	3 days
	Chamane	05002	UPL Europe Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Decis Protech	05269	Bayer CropScience Limited	Insecticide	<i>Deltamethrin</i>	<i>15 g/l</i>	830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
	Globaztar SC	05529	Globachem nv	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Karma	05116	Certis Europe BV	Fungicide	<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>	3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide	<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>	1.0 kg/ha	3.0 kg/ha	-	N/A	
	Movento	05349	Bayer CropScience Limited	Insecticide	<i>Spirotetramat</i>	<i>150 g/l</i>	0.5 l/ha	2.0 l/ha	4 per crop	14 days before harvest	14 days
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide	<i>Pymetrozine</i>	<i>50 %w/w</i>	0.4 kgs / ha	1,2 kgs/ha	3	3 days before harvest	3 days

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Courgettes - protected</i>											
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3.0 l/ha	6.0 l/ha	-	3 days before harvest	3 days
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
	Revus	05154	Syngenta UK Ltd	Fungicide			0.6 l/ha	2.4 l/ha	4 per crop	3 days before harvest	3 days
					<i>Mandipropamid</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			0.8 kg/ha	2.4 kg/ha	3	3 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	4.5 kgs/ha	3 / crop	1 day before harvest	1 day
					<i>Fenhexamid</i>	<i>50 %w/w</i>					
	Vivando	05685	BASF plc	Fungicide			0.2 l/ha	0.4 l/ha	-	3 days before harvest	3 days
					<i>Metrafenone</i>	<i>500 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cranberries - field</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis</i>	375 g/kg					
					<i>subspecies kurstaki EG</i>						
					<i>2348</i>						
	Masai	03585	BASF plc	Insecticide			1.5 kgs/ha	1.5 kgs/ha	-	7 days before harvest	
					<i>Tebufenpyrad</i>	200 g/kg					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium</i>	2% w/w					
					<i>anisoplice - ST</i>						
					<i>BIPESCO 5/F52</i>						
	Scala	03916	BASF plc	Fungicide			2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Pyrimethanil</i>	400 g/l					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus</i>	1015.1 g/l					
					<i>amyloliquefaciens str.</i>						
					<i>QST 713</i>						
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days
					<i>Spinosad</i>	480 g/l					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cranberries - protected</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kgs/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	375 g/kg					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>			Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	2% w/w					
	Scala	03916	BASF plc	Fungicide			2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Pyrimethanil</i>	400 g/l					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m <sup>2</sup> *	0.5 g/m <sup>2</sup>	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	12% w/w					
					<i>Trichoderma Asperellum (strain T34)</i>	12% w/w					
<i>Cress - field</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	1.26 l/ha	3	7 days before harvest	7 days
					<i>Deltamethrin</i>	15 g/l					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cucumbers - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Amistar Top	04582	Syngenta UK Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Apollo 50SC	04610	Adama Agricultural Solutions UK Ltd	Acaricide			0.4 l/ha	0.4 l/ha	1 per year	9 days before harvest	9 days
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide	<i>Clofentezine</i>	<i>500 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Azoxystar	05534	Life Scientific Limited	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	CaLypso	02441	Bayer CropScience Limited	Insecticide	<i>Thiacloprid</i>	<i>480 g/l</i>	0.45 l/ha	0.9 l/ha	2 per crop	3 days before harvest	3 days
	Chamane	05002	UPL Europe Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Decis Protech	05269	Bayer CropScience Limited	Insecticide	<i>Deltamethrin</i>	<i>15 g/l</i>	830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
	Globaztar SC	05529	Globachem nv	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Karma	05116	Certis Europe BV	Fungicide	<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>	3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
	Luna Privilege	05916	Bayer CropScience Limited	Fungicide	<i>Fluopyram</i>	<i>500 g/l</i>	0.6 l/ha	1.2 l/ha	2	1 day before harvest	1 day
	Microthiol Special	90548	UPL Europe Ltd	Fungicide	<i>Sulphur</i>	<i>80 %w/w</i>	0.01kgs/5 litres water	-	-	-	0 days
	Movento	05349	Bayer CropScience Limited	Insecticide	<i>Spirotetramat</i>	<i>150 g/l</i>	0.5 l/ha	2.0 l/ha	4 per crop	14 days before harvest	14 days

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Cucumbers - protected</i>											
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kg/ha	1.2 kgs/ha	3	3 days before harvest	3 days
	Revus	05154	Syngenta UK Ltd	Fungicide	<i>Pymetrozine</i>	<i>50 %w/w</i>	0.6 l/ha	2.4 l/ha	4 per crop	3 days before harvest	3 days
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Mandipropamid</i>	<i>250 g/l</i>	8 l/ha	48 l/ha	6		0 days
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	10 g/m3	10 g/m3	1	Pre planting / sowing	
	Teldor	02003	Bayer CropScience Limited	Fungicide	<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>	1.5 kgs/ha	4.5 kgs/ha	3 / crop	1 day before harvest	1 day
	Vivando	05685	BASF plc	Fungicide	<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>	0.2 l/ha	0.4 l/ha	-	3 days before harvest	3 days
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide	<i>Fenhexamid</i>	<i>50 %w/w</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Metrafenone</i>	<i>500 g/l</i>					
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
<i>Damsons</i>											
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	8 l/ha	48 l/ha	6		0 days

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Dill</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Emerger	06190	Bayer CropScience Limited	Herbicide			3.0 l/ha	3.0 l/ha	1 per crop	Before crop germination	
					<i>Acifluorfen</i>	600 g/l					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquifaciens str. QST 713</i>	1015.1 g/l					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	50 %w/w					
<i>Elderberries</i>											
	Batavia	05905	Bayer CropScience Limited	Insecticide			0.75 l/ha	1.5 l/ha	2	Post harvest application	
					<i>Spirotetramat</i>	100 g/l					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Endives</i>										
Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
				<i>Acetamiprid</i>	<i>20% w/w</i>					
Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	1.26 l/ha	3	7 days before harvest	7 days
				<i>Deltamethrin</i>	<i>15 g/l</i>					
Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2	3 days pre harvest	3 days
				<i>Acetamiprid</i>	<i>20 % w/w</i>					
Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
				<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
				<i>Spirotetramat</i>	<i>150 g/l</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
<i>Fennel</i>										
Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
				<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
				<i>Fenhexamid</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Garlic</i>											
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Basagran SG	05192	BASF plc	Herbicide			1.65 kg/ha	1.65 kg/ha	-	21 days before harvest	21 days
					<i>Bentazone</i>	<i>870 g/kg</i>					
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	56 days before harvest	56 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	CIPC	03066	Arysta LifeScience Benelux SPRL	Herbicide			1.8 l/ha	5.4 l/ha	3 per crop	GS 14	
					<i>Chlorpropham</i>	<i>400 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before 5th leaf visible	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Emerger	06190	Bayer CropScience Limited	Herbicide			2.5 l/ha	2.5 l/ha	2 per crop	GS 12	
					<i>Aclonifen</i>	<i>600 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Garlic</i>											
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide			0.2 l/ha	0.2 l/ha	1 per year	After transplanting or beginning of root formation	
	Folio Gold	02363	Syngenta UK Ltd	Fungicide	<i>Isoxaben</i>	<i>500 g/l</i>	2.0 l/ha	6.0 l/ha	3	14 days before harvest	
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide	<i>Chlorothalonil</i>	<i>500 g/l</i>					
					<i>Metalaxyl M</i>	<i>37.5 g/l</i>	3.0 l/ha	3.0 l/ha	1	4 weeks before harvest	28 days
	Infinito	04309	Bayer CropScience Limited	Fungicide	<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>	1.6 l/ha	1.6 l/ha	1 per crop	7 days before harvest	7 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			75 mls/ha	225 mls/ha	3 per crop	21 days before harvest	21 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Lentagran WP	03082	Belchim Crop Protection NV	Herbicide			2.0 kg/ha	2.0 kg/ha	1	Before 7 leaf stage	
					<i>Pyridate</i>	<i>45 % w/w</i>					
	NEMGuard DE	05368	ECOspray Ltd	Nematicide			20 kg/ha	20 kg/ha	1	At drilling	
					<i>Garlic Extract</i>	<i>450 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			75 mls/ha	225 mls/ha	3	21 days before harvest	21 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 15	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Starane Hi-Load H	04549	Dow AgroSciences Ltd	Herbicide			0.3 l/ha	0.3 l/ha		Before GS 15	
					<i>Fluroxypyr</i>	<i>333 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Garlic</i>											
	Stomp 400 SC	02879	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	Pre-emergence / Pre-transplanting	
					<i>Pendimethalin</i>	<i>250g/l</i>					
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Gherkin - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	FLiPPER	05027	Alpha Biopesticides Ltd	Insecticide			16.0 l/ha	80.0 l/ha	5	N/A	
					<i>Fatty Acid - C7-C20</i>	<i>479.8 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3 mls per sq m	6 mls per sq m	-	Before transplanting	
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
	Revus	05154	Syngenta UK Ltd	Fungicide			0.6 l/ha	2.4 l/ha	4 per crop	3 days before harvest	3 days
					<i>Mandipropamid</i>	<i>250 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Gherkin - protected</i>											
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	4.5 kgs/ha	3	1 day before harvest	1 day
					<i>Fenhexamid</i>	<i>50 %w/w</i>					
	Vivando	05685	BASF plc	Fungicide			0.2 l/ha	0.4 l/ha	-	3 days before harvest	3 days
					<i>Metrafenone</i>	<i>500 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Gooseberries - field</i>											
Batavia	05905	Bayer CropScience Limited	Insecticide				0.75 l/ha	1.5 l/ha	2	Post harvest application	
					<i>Spirotetramat</i>	<i>100 g/l</i>					
Botanigard WP	06264	Mycotech Europe Ltd	Insecticide				0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
CaLypso	02441	Bayer CropScience Limited	Insecticide				0.25 l/ha	0.75 l/ha	3 per year	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
Chikara Weed Cont	04203	ISK Biosciences Europe N.V.	Herbicide				120 g/ha	120 g/ha	1 per year	Before bud burst	
					<i>Flazasulfuron</i>	<i>25 % w/w</i>					
Decis Protech	05269	Bayer CropScience Limited	Insecticide				400 ml/ha	800 ml/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
Karate Zeon	04084	Syngenta UK Ltd	Insecticide				100 mls/ha	200 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide				1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
Manfil 75 WG	03598	Indofil Industries Limited	Fungicide				1.5 kg per 1000 litres water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75 % w/w</i>					
Manfil 80 WP	03563	Indofil Industries Limited	Fungicide				1.4 kg per 1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
Masai	03585	BASF plc	Insecticide				1.5 kgs/ha	1.5 kgs/ha	-	7 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
Microthiol Special	90548	UPL Europe Ltd	Fungicide				0.25 kgs/100 litres water	-	3	Up to and including fruit swell	
					<i>Sulphur</i>	<i>80 %w/w</i>					
Ninja 10CS	05178	Syngenta UK Ltd	Insecticide				100 mls/ha	200 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Gooseberries - field</i>											
	Penncozeb 80 WP	03609	UPL Europe Ltd	Fungicide			1.4 kg/1000L water	-	-	28 days before harvest	28 days
	Penncozeb WDG	03608	UPL Europe Ltd	Fungicide	<i>Mancozeb</i>	<i>800 g/kg</i>	1.5 kg/1000L water	-	-	28 days before harvest	28 days
	Radspor 400	01797	Arysta LifeScience Benelux SPRL	Fungicide	<i>Mancozeb</i>	<i>750 g/kg</i>	73 mls/100 litres water	-	5		
	Scala	03916	BASF plc	Fungicide	<i>Dodine</i>	<i>40 %w/w</i>	2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Pyrimethanil</i>	<i>400 g/l</i>	8 l/ha	48 l/ha	6		0 days
	Signum	03799	BASF plc	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	1.5 kgs/ha	4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest)	-	3 days before harvest, N/A for post harvest	
	Stroby WG	05092	BASF plc	Fungicide	<i>Boscalid</i>	<i>26.7 % w/w</i>	0.2 kg/ha	0.6 kg/ha	-	14 days before harvest	14 days
	Switch	03761	Syngenta UK Ltd	Fungicide	<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>	1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
	Tracer	02649	Dow AgroSciences Ltd	Insecticide	<i>Kresoxim Methyl</i>	<i>50% w/w</i>	200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Gooseberries - protected</i>											
AQ 10	05801	CBC (Europe) S.r.l.	Fungicide	<i>Ampelomyces Quisqualis</i>	58 %w/w	70 g/ha	-	-	12	-	
Botanigard WP	06264	Mycotech Europe Ltd	Insecticide	<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg	0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour		
Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide	<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	375 g/kg	1.0 kgs/ha	3.0 kgs/ha	-	N/A		
Scala	03916	BASF plc	Fungicide	<i>Pyrimethanil</i>	400 g/l	2.0 l/ha	4.0 l/ha	2	21 days before harvest	21 days	
Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l	8 l/ha	48 l/ha	6		0 days	
T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide	<i>Trichoderma Asperellum (strain T34)</i> <i>Trichoderma Asperellum (strain T34)</i>	12% w/w 12% w/w	0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings		
Tracer	02649	Dow AgroSciences Ltd	Insecticide	<i>Spinosad</i>	480 g/l	200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days	
<i>Grassland - seed crops</i>											
Decis Protech	05269	Bayer CropScience Limited	Insecticide	<i>Deltamethrin</i>	15 g/l	420 mls/ha	1.26 l/ha	3	N/A		

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Herbs - field</i>											
	Acrobat MZ	04059	BASF plc	Fungicide			2.0 kg/ha	6.0 kg/ha	-	21 days before harvest	21 days
					<i>Dimethomorph</i>	<i>75 g/kg</i>					
					<i>Mancozeb</i>	<i>667 g/kg</i>					
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Bonalan	05763	Gowan Crop Protection Ltd	Herbicide			8.0 l/ha	8.0 l/ha	-	Incorporation pre sowing/planting	
					<i>Benfluralin</i>	<i>180 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	<i>220 g/kg</i>					
					<i>GHA / ATCC 74040</i>						
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	CIPC	03066	Arysta LifeScience Benelux SPRL	Herbicide			1.9 l/ha	3.8 l/ha	2 per crop	2 weeks after planting out	
					<i>Chlorpropham</i>	<i>400 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 ml/ha	1.26 l/ha	3 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Devrinol	05234	UPL Europe Ltd	Herbicide			0.85 l/ha	0.85 l/ha	-	Pre crop emergence	
					<i>Napropamide</i>	<i>450 g/l</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Herbs - field</i>											
	Globazar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	1.6 l/ha	1 per crop	Before GS 49	14 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			75 mls/ha	150 mls/ha	4	7 days pre harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Kerb Flo	02830	Dow AgroSciences Ltd	Herbicide			1.875 l/ha	1.875 l/ha	1	24 days before harvest	24 days
					<i>Propyzamide</i>	<i>400 g/l</i>					
					<i>Propyzamide</i>	<i>400 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			75 mls/ha	150 mls/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	PDM 330 EC	03582	BASF plc	Herbicide			4.5 l/ha	4.5 l/ha	1	6 weeks pre harvest	
					<i>Pendimethalin</i>	<i>330 g/l</i>					
	Pendifin 400 SC	05731	Finchimica S.p.A	Herbicide			3.75 l/ha	3.75 l/ha	1 per crop	6 weeks before harvest	6 weeks
					<i>Pendimethalin</i>	<i>400 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Herbs - field</i>										
Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kgs/ha	1.2 kgs/ha	3	14 days before harvest	14 days
				<i>Pymetrozine</i>	<i>50 %w/w</i>					
Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
				<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
Scala	03916	BASF plc	Fungicide			1.33 l/ha	4.0 l/ha	-	14 days before harvest	14 days
				<i>Pyrimethanil</i>	<i>400 g/l</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
Signum	03799	BASF plc	Fungicide			1.5 kgs/ha	3.0 kgs/ha	-	14 days before harvest	
				<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
				<i>Boscalid</i>	<i>26.7 % w/w</i>					
Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
				<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
Stomp 400 SC	02879	BASF plc	Herbicide			3.75 l/ha	3.75 l/ha	1 / crop	6 weeks pre harvest	6 weeks
				<i>Pendimethalin</i>	<i>400 g/l</i>					
Stomp Aqua	03750	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1 / crop	6 weeks pre harvest	
				<i>Pendimethalin</i>	<i>455 g/l</i>					
Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
				<i>Cycloxydim</i>	<i>100 g/l</i>					
Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha	2	3 days before harvest	3 days
				<i>Fenhexamid</i>	<i>50 %w/w</i>					
Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha	600 ml/ha	3 per year	3 days before harvest	3 days
				<i>Spinosad</i>	<i>480 g/l</i>					
Wing P	03590	BASF plc	Herbicide			1.25 l/ha	1.25 l/ha	-	Before transplanting	35 days
				<i>Pendimethalin</i>	<i>250g/l</i>					
				<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					
Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Herbs - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
	AQ 10	05801	CBC (Europe) S.r.l.	Fungicide	<i>Azoxystrobin</i>	250 g/l	70 g/ha	-	12	-	
					<i>Ampelomyces Quisqualis</i>	58 %w/w					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	250 g/l					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	250 g/l					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.25 l/ha	1 per crop	14 days before harvest	14 days
					<i>Thiacloprid</i>	480 g/l					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	250 g/l					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	1.66 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	15 g/l					
	Devrinol	05234	UPL Europe Ltd	Herbicide			0.85 l/ha	0.85 l/ha	-	Pre crop emergence	
					<i>Napropamide</i>	450 g/l					
	Dynamec	02640	Syngenta UK Ltd	Acaricide/Insecticide			50 mls/product/100 litres water	-	4	14 days before harvest	14 days
					<i>Abamectin</i>	18 g/l					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	250 g/l					
	Kerb Flo	02830	Dow AgroSciences Ltd	Herbicide			1.875 l/ha	1.875 l/ha	1	24 days pre harvest	24 days
					<i>Propyzamide</i>	400 g/l					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	2% w/w					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Herbs - protected</i>											
	Microthiol Special	90548	UPL Europe Ltd	Fungicide			0.01kgs/5 litres water	-	2	-	0 days
	Movento	05349	Bayer CropScience Limited	Insecticide	<i>Sulphur</i>	<i>80 %w/w</i>	0.5 l/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
	Paraat	03614	BASF plc	Fungicide	<i>Spirotetramat</i>	<i>150 g/l</i>	0.36 kg/ha	0.72 kg/ha	2 per crop	14 days before harvest	14 days
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide	<i>Dimethomorph</i>	<i>500 g/kg</i>	0.4 kg/ha	1.2 kg/ha	3	14 days before harvest	14 days
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide	<i>Pymetrozine</i>	<i>50 %w/w</i>	2.5 l/ha	5.0 l/ha	-	21 days before harvest	21 days
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide			1.9 kg/ha	5.7 kg/ha*		35 days before harvest	35 days
					<i>Mancozeb</i>	<i>64 %w/w</i>					
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
	Scala	03916	BASF plc	Fungicide			1.33 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.5 kgs/ha	3.0 kgs/ha	-	14 days before harvest	
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha	600 ml/ha	3 per year	3 days before harvest	3 days
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product</i>					<i>Max Ind.</i>	<i>Max. Total</i>	<i>Max No of</i>	<i>Latest Time</i>	
<i>Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Dose:</i>	<i>Dose:</i>	<i>Applic's:</i>	<i>of Applic:</i>	<i>PHI:</i>
<i>Herbs - protected</i>										
Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Horseradish</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>240 g/l</i>					
	Butisan S	04631	BASF plc	Herbicide			1.0 l/ha	1.0 l/ha	-	Pre-emergence	
					<i>Metazachlor</i>	<i>500 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	14 days pre harvest	14 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Corbel	90438	BASF plc	Fungicide			1.0 l/ha	3.0 l/ha	3	28 days before harvest	28 days
					<i>Fenpropimorph</i>	<i>750 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Horseradish</i>											
	Devrinol	05234	UPL Europe Ltd	Herbicide			1.25 l/ha	1.25 l/ha	-	Pre crop emergence	
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide	<i>Napropamide</i>	<i>450 g/l</i>	2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide	<i>Mancozeb</i>	<i>75% w/w</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	0.15 l/ha	0.15 l/ha	1 per year	Pre-emergence	
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide	<i>Isoxaben</i>	<i>500 g/l</i>	3.0 l/ha	3.0 l/ha	1	8 weeks before harvest	56 days
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide	<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>	250 g/ha	500 g/ha	2 per crop	14 days before harvest	14 days
	Globaztar SC	05529	Globachem nv	Fungicide	<i>Acetamiprid</i>	<i>20 % w/w</i>	1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
	Goltix 70 SC	04668	Adama Agricultural Solutions UK Ltd	Herbicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	2.0 l/ha	5.0 l/ha	-	Pre crop emergence	
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide	<i>Metamitron</i>	<i>700 g/l</i>	150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
	Markate 50	05001	Globachem nv	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide	<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>	0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Horseradish</i>											
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.3 kg/ha	0.9 kg/ha	* 3 per crop	21 days before harvest	21 days
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
					<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	PDM 330 EC	03582	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>330 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Rapsan 500SC	05886	Globachem nv	Herbicide			1.0 l/ha	1.0 l/ha	-	Pre-emergence	
					<i>Metazachlor</i>	<i>500 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.0 kg/ha	2.0 kgs/ha	-	14 days before harvest	
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
	SL567A	03106	Syngenta UK Ltd	Fungicide			1.3 l/ha	1.3 l/ha	1	30 days before harvest	30 days
					<i>Metalaxyl M</i>	<i>465.2 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Horseradish</i>											
	Stomp 400 SC	02879	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	8 weeks before harvest	8 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	Venzar 500 SC	04805	DuPont (UK) Ltd	Herbicide			0.4 l/ha	0.4 l/ha	-	Pre-emergence	
					<i>Lenacil</i>	<i>500 g/l</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Jerusalem Artichoke</i>											
	Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	8 weeks before harvest	8 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Kale</i>											
Anthem	04761	Adama Agricultural Solutions UK Ltd	Herbicide				3.3 l/ha	3.3 l/ha	-		
					<i>Pendimethalin</i>	<i>400 g/l</i>					
Biscaya	03602	Bayer CropScience Limited	Insecticide				0.4 l/ha	0.8 l/ha	2	21 days before harvest	21 days
					<i>Thiacloprid</i>	<i>240 g/l</i>					
Dual Gold	04192	Syngenta UK Ltd	Herbicide				1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
Efica 960 EC	05942	Syngenta UK Ltd	Herbicide				1.4 l/ha	1.4 l/ha	-	Drilled crops- pre-emergence. Or within 5 days transplanting	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide				0.25 l/ha	0.25 l/ha	-	Before 6 true leaf stage	
					<i>Clomazone</i>	<i>360 g/l</i>					
Infinito	02532	Bayer CropScience Limited	Fungicide				1.6 l/ha	3.2 l/ha	2 per crop*	Before GS 49	14 days
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
Lentagran WP	03082	Belchim Crop Protection NV	Herbicide				2.0 kg/ha	2.0 kg/ha	1	42 days before harvest	
					<i>Pyridate</i>	<i>45 % w/w</i>					
Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide				0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide				0.4 kg/ha	1.2 kg/ha	3	14 days before harvest	14 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide				8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Kale</i>											
Signum	03799	BASF plc		Fungicide			1.0 kg/ha	3.0 kg/ha	3 per crop	14 days before harvest	14 days
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS		Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
					<i>Esfenvalerate</i>	<i>25 g/l</i>					
T34 Biocontrol	05620	Biocontrol Technologies S.L.		Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
Tracer	02649	Dow AgroSciences Ltd		Insecticide			12 mls/product/plant	-	1	Pre planting - 6 leaf stage	
					<i>Spinosad</i>	<i>480 g/l</i>					
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Kohl-rabi</i>											
	Butisan S	04631	BASF plc	Herbicide			1.0 l/ha	1.0 l/ha	-	Pre-emergence	
					<i>Metazachlor</i>	<i>500 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	1.0 l/ha	2 per crop	3 days before harvest	3 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			165 mls/ha	330 mls/ha	2	7 days pre harvest	
					<i>Esfenvalerate</i>	<i>25 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m <sup>3</sup>	10 g/m <sup>3</sup>	1	Pre planting / sowing	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Leeks</i>											
	Acrobat MZ	04059	BASF plc	Fungicide			2.0 kg/ha	6.0 kg/ha	-	21 days before harvest	21 days
					<i>Dimethomorph</i>	<i>75 g/kg</i>					
					<i>Mancozeb</i>	<i>667 g/kg</i>					
	Basagran SG	05192	BASF plc	Herbicide			0.4 kg/ha	0.8 kg/ha	-	Before 3 true leaf stage	
					<i>Bentazone</i>	<i>870 g/kg</i>					
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Butisan S	04631	BASF plc	Herbicide			1.5 l/ha	1.5 l/ha	-	5-21 days after planting	
					<i>Metazachlor</i>	<i>500 g/l</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.2 l/ha	0.6 l/ha	3 per year	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			250 mls/ha	750 mls/ha	3 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	1.26 l/ha	-	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before 5th leaf visible	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide			0.3 l/ha	0.3 l/ha	1 per year	After transplanting	
					<i>Isoxaben</i>	<i>500 g/l</i>					
	Folio Gold	02363	Syngenta UK Ltd	Fungicide			2.0 l/ha	6.0 l/ha	3	14 days before harvest	
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
					<i>Metalaxyl M</i>	<i>37.5 g/l</i>					
	Infinito	02532	Bayer CropScience Limited	Fungicide			1.6 l/ha	1.6 l/ha	1 per crop	14 days before harvest	14 days
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Leeks</i>											
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			100 mls/ha	200 mls/ha	2 per crop	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Lentagran WP	03082	Belchim Crop Protection NV	Herbicide			2.0 kg/ha	2.0 kg/ha	1	28 days before harvest	
					<i>Pyridate</i>	<i>45 % w/w</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.36 kg/ha	0.36 kg/ha	* 1 per crop	21 days before harvest	21 days
					<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
	NEMGuard DE	05368	ECOspray Ltd	Nematicide			20 kg/ha	20 kg/ha	1	At drilling	
					<i>Garlic Extract</i>	<i>450 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			100 mls/ha	200 mls/ha	2	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 15	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Starane Hi-Load H	04549	Dow AgroSciences Ltd	Herbicide			0.3 l/ha	0.3 l/ha		28 days before harvest	28 days
					<i>Fluroxypyr</i>	<i>333 g/l</i>					
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	Pre-emergence / Pre-transplanting	
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					
					<i>Pendimethalin</i>	<i>250g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Lettuce - field</i>											
	Acrobat MZ	04059	BASF plc	Fungicide			2.0 kg/ha	6.0 kg/ha	-	21 days before harvest	21 days
					<i>Dimethomorph</i>	<i>75 g/kg</i>					
					<i>Mancozeb</i>	<i>667 g/kg</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	<i>220 g/kg</i>					
					<i>GHA / ATCC 74040</i>						
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20 % w/w</i>					
	Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	1.6 l/ha	1 per crop	Before GS 49	14 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Kerb Flo	02830	Dow AgroSciences Ltd	Herbicide			1.875 l/ha	1.875 l/ha	1	24 days before harvest	24 days
					<i>Propyzamide</i>	<i>400 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Lettuce - field</i>											
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide	<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>	0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
	PDM 330 EC	03582	BASF plc	Herbicide	<i>Pendimethalin</i>	<i>330 g/l</i>	4.5 l/ha	4.5 l/ha	1	6 weeks pre harvest	
	Pendifin 400 SC	05731	Finchimica S.p.A	Herbicide	<i>Pendimethalin</i>	<i>400 g/l</i>	3.75 l/ha	3.75 l/ha	1 per crop	6 weeks before harvest	6 weeks
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide	<i>Pymetrozine</i>	<i>50 %w/w</i>	0.4 kg/ha	1.2 kg/ha	3	7 days before harvest	7 days
	Scala	03916	BASF plc	Fungicide	<i>Pyrimethanil</i>	<i>400 g/l</i>	1.33 l/ha	4.0 l/ha	-	14 days before harvest	14 days
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide	<i>Difenoconazole</i>	<i>250 g/l</i>	0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
	Stomp 400 SC	02879	BASF plc	Herbicide	<i>Pendimethalin</i>	<i>400 g/l</i>	3.75 l/ha	3.75 l/ha	1 / crop	6 weeks pre harvest	6 weeks
	Stomp Aqua	03750	BASF plc	Herbicide	<i>Pendimethalin</i>	<i>455 g/l</i>	3.3 l/ha	3.3 l/ha	1 / crop	6 weeks pre harvest	
	Switch	03761	Syngenta UK Ltd	Fungicide	<i>Fludioxonil</i> <i>Cyprodinil</i>	<i>25.0 % w/w</i> <i>37.5 % w/w</i>	0.7 kg/ha	1.4 kg/ha	2	7 days before harvest	
	Systhane 20 EW	04944	Dow AgroSciences Ltd	Fungicide	<i>Myclobutanil</i>	<i>200 g/l</i>	0.3 l/ha	0.6 l/ha	2	14 days before harvest	14 days
	Teldor	02003	Bayer CropScience Limited	Fungicide	<i>Fenhexamid</i>	<i>50 %w/w</i>	1.5 kgs/ha	3.0 kgs/ha	2 / crop	3 days before harvest	3 days
	Tracer	02649	Dow AgroSciences Ltd	Insecticide	<i>Spinosad</i>	<i>480 g/l</i>	200 ml/ha	600 ml/ha	3 per crop	3 days before harvest	3 days
	Wing P	03590	BASF plc	Herbicide	<i>Dimethenamid-P</i> <i>Pendimethalin</i>	<i>212.5 g/l</i> <i>250g/l</i>	1.25 l/ha	1.25 l/ha	-	Before transplanting	35 days

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Lettuce - protected</i>											
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	<i>220 g/kg</i>					
					<i>GHA / ATCC 74040</i>						
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.25 l/ha	1 per crop	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	1.66 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Dynamec	02640	Syngenta UK Ltd	Acaricide/Insecticid			50 mls/product/100 litres water	-	4	14 days before harvest	14 days
					<i>Abamectin</i>	<i>18 g/l</i>					
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20 % w/w</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Kerb Flo	02830	Dow AgroSciences Ltd	Herbicide			1.875 l/ha	1.875 l/ha	1	24 days pre harvest	24 days
					<i>Propyzamide</i>	<i>400 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Paraat	03614	BASF plc	Fungicide			0.36 kg/ha	0.72 kg/ha	2 per crop	14 days before harvest	14 days
					<i>Dimethomorph</i>	<i>500 g/kg</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kg/ha	1.2 kgs/ha	3	14 days before harvest	14 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Lettuce - protected</i>											
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide			1.9 kg/ha	5.7 kg/ha*		35 days before harvest	35 days
					<i>Mancozeb</i>	<i>64 %w/w</i>					
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
	Scala	03916	BASF plc	Fungicide			1.33 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			0.7 kg/ha	1.4 kg/ha	2	7 days before harvest	7 days
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	Sythane 20 EW	04944	Dow AgroSciences Ltd	Fungicide			0.3 l/ha	0.6 l/ha	2	14 days before harvest	14 days
					<i>Myclobutanil</i>	<i>200 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha	2 / crop	3 days before harvest	3 days
					<i>Fenhexamid</i>	<i>50 %w/w</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha	600 ml/ha	3 per crop	3 days before harvest	3 days
					<i>Spinosad</i>	<i>480 g/l</i>					
<i>Linseed</i>											
	Butisan S	04631	BASF plc	Herbicide			1.5 l/ha	1.5 l/ha	-	Before GS 19	
					<i>Metazachlor</i>	<i>500 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	500 ml/ha	1 per crop	GS 81	
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Rapsan 500SC	05886	Globachem nv	Herbicide			1.5 l/ha	1.5 l/ha	-	GS 19	
					<i>Metazachlor</i>	<i>500 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Loganberries - field</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis</i>	375 g/kg					
					<i>subspecies kurstaki EG</i>						
					<i>2348</i>						
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 g/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	200 g/kg					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium</i>	2% w/w					
					<i>anisoplice - ST</i>						
					<i>BIPESCO 5/F52</i>						
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus</i>	1015.1 g/l					
					<i>amyloliquefaciens str.</i>						
					<i>QST 713</i>						
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha	-	21 days before harvest	21 days
					<i>Carfentrazone-ethyl</i>	60 g/l					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Loganberries - protected</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 l/ha	3.0 l/ha	-	N/A	
					<i>Bacillus thuringiensis</i>	375 g/kg					
					<i>subspecies kurstaki EG</i>						
					<i>2348</i>						
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 gms/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	200 g/kg					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>			pre-planting	
					<i>Metarhizium</i>	2% w/w					
					<i>anisoplice - ST</i>						
					<i>BIPESCO 5/F52</i>						
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6	-	0 days
					<i>Bacillus</i>	1015.1 g/l					
					<i>amyloliquefaciens str.</i>						
					<i>QST 713</i>						
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha	-	21 days before harvest	21 days
					<i>Carfentrazone-ethyl</i>	60 g/l					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m <sup>3</sup>	10 g/m <sup>3</sup>	1	Pre planting / sowing	
					<i>Trichoderma</i>	12% w/w					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	12% w/w					
					<i>Asperellum (strain T34)</i>						

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Lupins</i>											
	Anthem	04761	Adama Agricultural Solutions UK Ltd	Herbicide			3.75 l/ha	3.75 l/ha		Pre crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	PDM 330 EC	03582	BASF plc	Herbicide			4.5 l/ha	4.5 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>330 g/l</i>					
	Pendifin 400 SC	05731	Finchimica S.p.A	Herbicide			3.75 l/ha	3.75 l/ha	1 per crop	Pre-crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
<i>Maize - Forage - open ground</i>											
	Comet	02354	BASF plc	Fungicide			1.0 l/ha	2.0 l/ha	2	Tasel/silk visible stage	
					<i>Pyraclostrobin</i>	<i>250 g/l</i>					
	Comet 200	03696	BASF plc	Fungicide			1.25 l/ha	2.5 l/ha	2	Tasel/silk visible stage (BBCH 51-59)	
					<i>Pyraclostrobin</i>	<i>200 g/l</i>					
	Modem	02355	BASF plc	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	Tasel/silk visible stage	
					<i>Pyraclostrobin</i>	<i>250 g/l</i>					
	Modem 200	03695	BASF plc	Fungicide			1.25 l/ha	2.5 l/ha	2	Tasel/silk visible stage (BBCH 51-59)	
					<i>Pyraclostrobin</i>	<i>200 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Maize - Forage - under plastic</i>											
	Comet 200	03696	BASF plc	Fungicide			1.25 l/ha	2.5 l/ha	2	Tassel/silk visible stage (BBCH 51-59)	
					<i>Pyraclostrobin</i>	<i>200 g/l</i>					
	Modem	02355	BASF plc	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	Tassel/silk visible stage	
					<i>Pyraclostrobin</i>	<i>250 g/l</i>					
	Modem 200	03695	BASF plc	Fungicide			1.25 l/ha	2.5 l/ha	2 per crop	Tassel/silk visible stage (BBCH 51-59)	
					<i>Pyraclostrobin</i>	<i>200 g/l</i>					
<i>Maize - Grain - open ground</i>											
	Comet	02354	BASF plc	Fungicide			1.0 l/ha	2.0 l/ha	2	Tassel/silk visible stage	
					<i>Pyraclostrobin</i>	<i>250 g/l</i>					
<i>Mange tout peas</i>											
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
<i>Mangolds</i>											
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Marrows</i>										
Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
				<i>Azoxystrobin</i>	<i>200 g/l</i>					
				<i>Difenoconazole</i>	<i>125 g/l</i>					
CaLypso	02441	Bayer CropScience Limited	Insecticide			0.45 l/ha	0.9 l/ha	2 per year	3 days before harvest	3 days
				<i>Thiacloprid</i>	<i>480 g/l</i>					
Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3 mls per sq m	6 mls per sq m	-	Before transplanting	
				<i>Propamocarb</i>	<i>530 g/l</i>					
				<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
				<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
				<i>Propamocarb</i>	<i>530 g/l</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
				<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Melon - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
	Amistar Top	04582	Syngenta UK Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	2.0 l/ha	4 per crop	14 days before harvest	14 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Revus	05154	Syngenta UK Ltd	Fungicide			0.6 l/ha	2.4 l/ha	4 per crop	3 days before harvest	3 days
					<i>Mandipropamid</i>	<i>250 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
<i>Mint</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	<i>50 %w/w</i>					
<i>Mushrooms</i>											
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			5ml/100 cubic metre	-	2 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Pyrethrum 5 EC	06255	Pelgar International	Insecticide			0.65 l/ha	1.95 l/ha	3 per crop	1 day before harvest	1 day
					<i>Pyrethrins</i>	<i>50 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Mustard</i>										
Fathom	04688	FMC Agro Ltd	Fungicide	<i>Tebuconazole</i>	<i>200 g/l</i>	1.25 l/ha	1.25 l/ha	1	GS 65	21 days
<i>Oilseed Rape - spring</i>										
Galera	05188	Dow AgroSciences Ltd	Herbicide	<i>Clopyralid</i>	<i>267 g/l</i>	0.35 l/ha	0.35 l/ha	-	Before flower buds visible above crop canopy	
				<i>Picloram</i>	<i>67 g/l</i>					
Proline	03786	Bayer CropScience Limited	Fungicide	<i>Prothioconazole</i>	<i>250 g/l</i>	0.7 l/ha	1.4 l/ha	-	56 days before harvest	56 days

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Onions</i>											
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	Basagran SG	05192	BASF plc	Herbicide			1.65 kg/ha	1.65 kg/ha	-	21 days before harvest	21 days
					<i>Bentazone</i>	<i>870 g/kg</i>					
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	56 days before harvest	56 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before 5th leaf visible	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Digital	03147	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	-	42 days before harvest	42 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Emerger	06190	Bayer CropScience Limited	Herbicide			2.5 l/ha	2.5 l/ha	2 per crop	GS 12	
					<i>Aclonifen</i>	<i>600 g/l</i>					
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide			0.2 l/ha	0.2 l/ha	1 per year	After transplanting or beginning of root formation	
					<i>Isoxaben</i>	<i>500 g/l</i>					
	Folio Gold	02363	Syngenta UK Ltd	Fungicide			2.0 l/ha	6.0 l/ha	3	14 days before harvest	
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
					<i>Metalaxyl M</i>	<i>37.5 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Onions</i>											
	Guillotine	03146	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	42 days before harvest	42 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	1.6 l/ha	1 per crop	7 days before harvest	7 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			75 mls/ha	225 mls/ha	3 per crop	21 days before harvest	21 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	NEMGuard DE	05368	ECOSpray Ltd	Nematicide			20 kg/ha	20 kg/ha	1	At drilling	
					<i>Garlic Extract</i>	<i>450 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			75 mls/ha	225 mls/ha	3	21 days before harvest	21 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 15	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Starane Hi-Load H	04549	Dow AgroSciences Ltd	Herbicide			0.3 l/ha	0.3 l/ha		Before GS 15	
					<i>Fluroxypyr</i>	<i>333 g/l</i>					
	Sumimax	02764	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	42 days before harvest	42 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Valbon	03411	Certis Europe BV	Fungicide			1.6 kg/ha	9.6 kg/ha	6 per crop	28 days pre-harvest	28 days
					<i>Benthiavdicarb</i>	<i>17.5 g/kg</i>					
					<i>Mancozeb</i>	<i>700 g/kg</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Onions</i>											
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	Pre-emergence / Pre-transplanting	
					<i>Pendimethalin</i>	<i>250g/l</i>					
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Onions - salad</i>											
	Acrobat MZ	04059	BASF plc	Fungicide			2.0 kg/ha	10.0 kg/ha	-	14 days before harvest	14 days
					<i>Mancozeb</i>	<i>667 g/kg</i>					
					<i>Dimethomorph</i>	<i>75 g/kg</i>					
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	21 days before harvest	21 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	1.0 l/ha	-	7 days before harvest	7 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	21 days before harvest	21 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	21 days before harvest	21 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Basagran SG	05192	BASF plc	Herbicide			0.55 kg/ha	1.1 kg/ha	-	21 days before harvest	21 days
					<i>Bentazone</i>	<i>870 g/kg</i>					
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.2 l/ha	0.6 l/ha	3 per year	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	21 days before harvest	21 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	1.26 l/ha	3	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha		Before 3rd true leaf	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Folio Gold	02363	Syngenta UK Ltd	Fungicide			2.0 l/ha	6.0 l/ha		14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
					<i>Metalaxyl M</i>	<i>37.5 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Onions - salad</i>											
	Globazar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	21 days before harvest	21 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	1.6 l/ha	1 per crop	14 days before harvest	14 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			75 mls/ha	150 mls/ha	2 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha		N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Manfil 75 WG	03598	Indofil Industries Limited	Fungicide			2.1 kg/ha	8.4 kg/ha	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75 % w/w</i>					
	Manfil 80 WP	03563	Indofil Industries Limited	Fungicide			2.0 kg/ha	8.0 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.36 kg/ha	0.36 kg/ha	* 1 per crop	21 days before harvest	21 days
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
					<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			75 mls/ha	150 mls/ha	2	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Penncozeb 80 WP	03609	UPL Europe Ltd	Fungicide			2.0 kg/ha	8.0 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
	Penncozeb WDG	03608	UPL Europe Ltd	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>750 g/kg</i>					
	Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide			1.5 kgs/ha	4.5 kgs/ha		14 days pre harvest	14 days
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
					<i>Mancozeb</i>	<i>64 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Onions - salad</i>											
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 13	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.5 kg/ha	3.0 kg/ha	-	14 days before harvest	14 days
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 per crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	21 days before harvest	
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	Pre-emergence	
					<i>Pendimethalin</i>	<i>250g/l</i>					
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	21 days before harvest	21 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
<i>Onions - sets</i>											
	Fathom	04688	FMC Agro Ltd	Fungicide			0.066 l/ha	0.066 l/ha	1	Prior to planting	
					<i>Tebuconazole</i>	<i>200 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Orchards - top fruit</i>											
Digital	03147	Sumitomo Chemical Agro Europe SAS	Herbicide				100 mls/ha	100 mls/ha		365 days before first harvest	
					<i>Flumioxazin</i>	<i>300 g/l</i>					
Dow Shield 100	03861	Dow AgroSciences Ltd	Herbicide				2.0 l/ha	2.0 l/ha	-	-	
					<i>Clopyralid</i>	<i>100 g/l</i>					
Fastnet	06253	Sipcam U.K. Ltd	Herbicide				3.75 l/ha	3.75 l/ha	1 per year	N/A	
					<i>Pendimethalin</i>	<i>365 g/l</i>					
Guillotine	03146	Sumitomo Chemical Agro Europe SAS	Herbicide				100 mls/ha	100 mls/ha	1	365 days before first harvest	
					<i>Flumioxazin</i>	<i>300 g/l</i>					
Most Micro	04837	Sipcam U.K. Ltd	Herbicide				3.75 l/ha	3.75 l/ha	1 per year	N/A	
					<i>Pendimethalin</i>	<i>365 g/l</i>					
Serenade ASO	03847	Bayer CropScience Limited	Fungicide				8 l/ha	48 l/ha	6		0 days
					<i>Bacillus anyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
Sumimax	02764	Sumitomo Chemical Agro Europe SAS	Herbicide				100 mls/ha	100 mls/ha	1	365 days before first harvest	
					<i>Flumioxazin</i>	<i>300 g/l</i>					
Vivendi 200	05122	UPL Europe Ltd	Herbicide				1.0 l/ha	1.0 l/ha	-	-	
					<i>Clopyralid</i>	<i>200 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsley - field</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	<i>220 g/kg</i>					
					<i>GHA / ATCC 74040</i>						
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Emerger	06190	Bayer CropScience Limited	Herbicide			2.5 l/ha	2.5 l/ha	1 per crop	Before crop germination	
					<i>Aclonifen</i>	<i>600 g/l</i>					
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20 % w/w</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsley - field</i>											
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
<i>Parsley - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20 % w/w</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsley Root</i>										
Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
				<i>Difenoconazole</i>	<i>125 g/l</i>					
				<i>Azoxystrobin</i>	<i>200 g/l</i>					
AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
				<i>Mancozeb</i>	<i>75% w/w</i>					
Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	14 days before harvest	14 days
				<i>Thiacloprid</i>	<i>240 g/l</i>					
Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
				<i>Clethodim</i>	<i>120 g/l</i>					
Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
				<i>Azoxystrobin</i>	<i>250 g/l</i>					
Corbel	90438	BASF plc	Fungicide			1.0 l/ha	3.0 l/ha	3	28 days before harvest	28 days
				<i>Fenpropimorph</i>	<i>750 g/l</i>					
Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
				<i>Deltamethrin</i>	<i>25 g/l</i>					
Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
				<i>Deltamethrin</i>	<i>15 g/l</i>					
Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
				<i>Prosulfocarb</i>	<i>800 g/l</i>					
Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
				<i>Mancozeb</i>	<i>75% w/w</i>					
Emerger	06190	Bayer CropScience Limited	Herbicide			2.5 l/ha	2.5 l/ha	2 per crop	GS 12	
				<i>Aclonifen</i>	<i>600 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsley Root</i>											
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	8 weeks before harvest	56 days
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.3 kg/ha	0.9 kg/ha	* 3 per crop	21 days before harvest	21 days
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
					<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsley Root</i>											
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Stomp 400 SC	02879	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsnips</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Corbel	90438	BASF plc	Fungicide			1.0 l/ha	3.0 l/ha	3	28 days before harvest	28 days
					<i>Fenpropimorph</i>	<i>750 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Digital	03147	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	-	28 days before harvest	28 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
					<i>Mancozeb</i>	<i>75% w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsnips</i>											
	Emerger	06190	Bayer CropScience Limited	Herbicide			1.5 l/ha	2.5 l/ha	2 per crop	GS 12	
	Fazor Star	03729	Arysta LifeScience Great Britain Ltd	Growth Regulator	<i>Aclonifen</i>	<i>600 g/l</i>	5.0 kg/ha	5.0 kg/ha	1 per crop	21 days before harvest	
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide	<i>Maleic hydrazide</i>	<i>600 g/kg</i>	0.15 l/ha	0.15 l/ha	1 per year	Pre-emergence	
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide	<i>Isoxaben</i>	<i>500 g/l</i>	3 l/ha	3 l/ha	1	8 weeks before harvest	8 weeks
	Globaztar SC	05529	Globachem nv	Fungicide	<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>	1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
	Goltix 70 SC	04668	Adama Agricultural Solutions UK Ltd	Herbicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	2.0 l/ha	5.0 l/ha	-	Pre crop emergence	
	Guillotine	03146	Sumitomo Chemical Agro Europe SAS	Herbicide	<i>Metamitron</i>	<i>700 g/l</i>	100 mls/ha	100 mls/ha	1	28 days before harvest	28 days
	Hurricane SC	04357	Adama Agricultural Solutions UK Ltd	Herbicide	<i>Flumioxazin</i>	<i>300 g/l</i>	0.24 l/ha	0.24 l/ha	-	Pre-emergence	
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide	<i>Diflufenican</i>	<i>500 g/l</i>	0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
	Microthiol Special	90548	UPL Europe Ltd	Fungicide	<i>Difenoconazole</i>	<i>250 g/l</i>	0.01kgs/5 litres water	-	2	-	0 days
	Movento	05349	Bayer CropScience Limited	Insecticide	<i>Sulphur</i>	<i>80 %w/w</i>	0.5 l/ha	1.0 l/ha	2 per crop	21 days before harvest	21 days
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide	<i>Spirotetramat</i>	<i>150 g/l</i>	0.3 kg/ha	0.9 kg/ha	* 3 per crop	21 days before harvest	21 days
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide	<i>Trifloxystrobin</i>	<i>250 g/kg</i>	2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Parsnips</i>											
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.0 kg/ha	2.0 kgs/ha	-	14 days before harvest	
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
	SL567A	03106	Syngenta UK Ltd	Fungicide			1.3 l/ha	1.3 l/ha	1	no later than 6 weeks after drilling	30 days
					<i>Metalaxyl M</i>	<i>465.2 g/l</i>					
	Sumimax	02764	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	28 days before harvest	28 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Pears</i>											
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.375 l/ha	0.75 l/ha	2 per year	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide			2.0 kg/ha	8.0 kg/ha	-	Before GS 07	
					<i>Copper oxychloride</i>	<i>87.8% w/w</i>					
	Exilis	04702	Fine Agrochemicals Ltd	Growth Regulator			75 ml per 10 litres water *	5 ml per 10 litres water	1 per year	Up to fruitlet diameter of 15 mm	75 days
					<i>6-benzyladenine</i>	<i>20.0 g/l</i>					
					<i>6-benzyladenine</i>	<i>20.0 g/l</i>					
	Fathom	04688	FMC Agro Ltd	Fungicide			0.325 l/ha	0.325 l/ha	1	After final harvest	
					<i>Tebuconazole</i>	<i>200 g/l</i>					
					<i>Tebuconazole</i>	<i>200 g/l</i>					
	Masai	03585	BASF plc	Insecticide			125 gms/1 m crown height		1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.25 l/ha	1.0 l/ha	-	28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Scala	03916	BASF plc	Fungicide			1.125 l/ha	4.5 l/ha	4 per year	56 days before harvest	56 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.25 l/ha	1.0 l/ha		28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			0.8 kg/ha	2.4 kg/ha	3	3 days before harvest	
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
	Talius	04604	DuPont (UK) Ltd	Fungicide			0.25 l/ha	0.5 l/ha	2 per year	49 days before harvest	49 days
					<i>Proquinazid</i>	<i>200 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Peas - combining</i>											
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			330 mls/ha	330 mls/ha	2	14 days pre harvest	
					<i>Esfenvalerate</i>	<i>25 g/l</i>					
	Teppeki	02683	ISK Biosciences Europe N.V.	Insecticide			0.14 kg/ha	0.14 kg/ha	1 per crop	14 days before harvest	14 days
					<i>Flonicamid</i>	<i>500 g/kg</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Peas - vining</i>											
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			1.0 l/ha	1.0 l/ha	-	30 days before harvest	30 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Digital	03147	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	-	42 days before harvest	42 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Guillotine	03146	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	42 days before harvest	42 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Stallion Sync Tech	05483	FMC Chemical s.p.r.l.	Herbicide			3.0 l/ha	3.0 l/ha	1 per crop	Pre-crop emergence	
					<i>Clomazone</i>	<i>30 g/l</i>					
					<i>Pendimethalin</i>	<i>333 g/l</i>					
	Sumi-Alpha	02334	Sumitomo Chemical Agro Europe SAS	Insecticide			330 mls/ha	330 mls/ha	2	14 days pre harvest	
					<i>Esfenvalerate</i>	<i>25 g/l</i>					
	Sumimax	02764	Sumitomo Chemical Agro Europe SAS	Herbicide			100 mls/ha	100 mls/ha	1	42 days before harvest	42 days
					<i>Flumioxazin</i>	<i>300 g/l</i>					
	Teppeki	02683	ISK Biosciences Europe N.V.	Insecticide			0.14 kg/ha	0.14 kg/ha	1 per crop	14 days before harvest	14 days
					<i>Flonicamid</i>	<i>500 g/kg</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product</i>					<i>Max Ind.</i>	<i>Max. Total</i>	<i>Max No of</i>	<i>Latest Time</i>	
<i>Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Dose:</i>	<i>Dose:</i>	<i>Applic's:</i>	<i>of Applic:</i>	<i>PHI:</i>
<i>Peppers - Chilli - protected</i>										
Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide	<i>Bacillus thuringiensis</i> <i>subspecies kurstaki EG</i> 2348	375 g/kg	1.0 kg/ha	3.0 kg/ha	-	N/A	

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Peppers - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	3 days before harvest	3 days
	Amistar Top	04582	Syngenta UK Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	2.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	3 days before harvest	3 days
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.45 l/ha	0.9 l/ha	2 per crop	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Dynamec	02640	Syngenta UK Ltd	Acaricide/Insecticid			50 mls/product/100 litres water	-	5	3 days before harvest	3 days
					<i>Abamectin</i>	<i>18 g/l</i>					
	FLiPPER	05027	Alpha Biopesticides Ltd	Insecticide			16.0 l/ha	80.0 l/ha	5	N/A	
					<i>Fatty Acid - C7-C20</i>	<i>479.8 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Microthiol Special	90548	UPL Europe Ltd	Fungicide			0.01kgs/5 litres water	-	-	-	0 days
					<i>Sulphur</i>	<i>80 %w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	2.0 l/ha	4 per crop	14 days before harvest	14 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.04 kgs/100 l water	-	3	3 days before harvest	3 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Peppers - protected</i>											
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3.0 l/ha	12.0 l/ha	-	3 days before harvest	3 days
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	-	7 days before harvest	
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	4.5 kgs/ha	3 / crop	1 day before harvest	1 day
					<i>Fenhexamid</i>	<i>50 %w/w</i>					
	Vivando	05685	BASF plc	Fungicide			0.3 l/ha	0.6 l/ha	-	3 days before harvest	3 days
					<i>Metrafenone</i>	<i>500 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
<i>Plums</i>											
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.3 l/ha	0.6 l/ha	2 per year	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide			2.0 kg/ha	8.0 kg/ha	-	Before GS 07	
					<i>Copper oxychloride</i>	<i>87.8% w/w</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			0.75 kg/ha	2.25 kg/ha	-	7 days before harvest	7 days
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
					<i>Boscalid</i>	<i>26.7 % w/w</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Potatoes - seed crops</i>											
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide			1.0 kg/ha	6.0 kg/ha	-	8 days before harvest	8 days
					<i>Copper oxychloride</i>	<i>87.8% w/w</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			670 ml/ha	1.34 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			5 l/ha	5 l/ha	1	GS 5	
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
<i>Potatoes - ware crops</i>											
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide			1.0 kg/ha	6.0 kg/ha	-	8 days before harvest	8 days
					<i>Copper oxychloride</i>	<i>87.8% w/w</i>					
<i>Pumpkins - field</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Pumpkins - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3 mls per sq m	6 mls per sq m	-	Before transplanting	
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6	-	0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
<i>Radicchio</i>											
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2	3 days pre harvest	3 days
					<i>Acetamiprid</i>	<i>20 % w/w</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Radish</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Butisan S	04631	BASF plc	Herbicide			1.0 l/ha	1.0 l/ha	-	Pre-emergence	
					<i>Metazachlor</i>	<i>500 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	8 weeks before harvest	56 days
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Rapsan 500SC	05886	Globachem nv	Herbicide			1.0 l/ha	1.0 l/ha	-		
					<i>Metazachlor</i>	<i>500 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Radish</i>											
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Isopyrazam</i>	<i>125 g/l</i>	8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	SL567A	03106	Syngenta UK Ltd	Fungicide	<i>Metalaxyl M</i>	<i>465.2 g/l</i>	1.3 l/ha	1.3 l/ha	1	30 days before harvest	30 days
	Sparviero	04469	Sipcam Oxon SpA	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Stratos Ultra	04936	BASF plc	Herbicide	<i>Cycloxydim</i>	<i>100 g/l</i>	4.0 l/ha	4.0 l/ha	1 per crop	8 weeks before harvest	8 weeks
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide	<i>Oxamyl</i>	<i>100 g/kg</i>	0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
<i>Rape - fodder</i>											
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>	8 l/ha	48 l/ha	6		0 days

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Raspberries - field</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
	Amistar Top	04582	Syngenta UK Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Apollo 50SC	04610	Adama Agricultural Solutions UK Ltd	Acaricide			0.4 l/ha	0.4 l/ha	1 per year	9 days before harvest	9 days
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide	<i>Clofentezine</i>	<i>500 g/l</i>	0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 g/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kgs/ha	1.2 kgs/ha	3	28 days before harvest	28 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
	Pyrimala	05429	Servem Ltd	Fungicide			2.0 l/ha	4.0 l/ha	2	7 days before harvest	7 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Scala	03916	BASF plc	Fungicide			2.0l/ha	4.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha	-	21 days before harvest	
					<i>Carfentrazone-ethyl</i>	<i>60 g/l</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200ml/ha in 1000L water	400 ml/ha	2 per year	1 day before harvest	1 day
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Raspberries - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	7 days before harvest	7 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Apollo 50SC	04610	Adama Agricultural Solutions UK Ltd	Acaricide			0.4 l/ha	0.4 l/ha	1 per year	9 days before harvest	9 days
					<i>Clofentezine</i>	<i>500 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.75 l/ha	3 per crop	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 g/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
	Pyrimala	05429	Servem Ltd	Fungicide			2.0 l/ha	4.0 l/ha	2	7 days before harvest	7 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Scala	03916	BASF plc	Fungicide			2.0 l/ha	4.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha		21 days before harvest	21 days
					<i>Carfentrazone-ethyl</i>	<i>60 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	14 days before harvest	
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Raspberries - protected</i>											
T34	Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
Tracer		02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha in 1000L water	600 ml/ha	3 per year	1 day before harvest	1 day
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Redcurrants - field</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.75 l/ha	3 per year	3 days before harvest	3 days
					<i>Thiacloprid</i>	480 g/l					
	Chikara Weed Cont	04203	ISK Biosciences Europe N.V.	Herbicide			120 g/ha	120 g/ha	1 per year	Before bud burst	
					<i>Flazasulfuron</i>	25 % w/w					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			400 ml/ha	800 ml/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	15 g/l					
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	Before flowering or after harvest	
					<i>Fluazifop-P (-butyl)</i>	125 g/l					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			100 mls/ha	200 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	100 g/l					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	375 g/kg					
	Manfil 75 WG	03598	Indofil Industries Limited	Fungicide			1.5 kg per 1000 litres water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	75 % w/w					
	Manfil 80 WP	03563	Indofil Industries Limited	Fungicide			1.4 kg per 1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	800 g/kg					
	Masai	03585	BASF plc	Insecticide			1.5 kgs/ha	1.5 kgs/ha	-	7 days before harvest	
					<i>Tebufenpyrad</i>	200 g/kg					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			100 mls/ha	200 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	100 g/l					
	Penncozeb 80 WP	03609	UPL Europe Ltd	Fungicide			1.4 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	800 g/kg					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Redcurrants - field</i>											
	Penncozeb WDG	03608	UPL Europe Ltd	Fungicide			1.5 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>750 g/kg</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kgs/ha	1.2 kgs/ha	3	28 days pre harvest	28 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
	Scala	03916	BASF plc	Fungicide			2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.5 kgs/ha	4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest)	-	3 days before harvest, N/A for post harvest	
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha in 250L water	400 ml/ha	2 per year	7 days before harvest	7 days
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Redcurrants - protected</i>											
AQ 10	05801	CBC (Europe) S.r.l.	Fungicide	<i>Ampelomyces Quisqualis</i>	58 %w/w	70 g/ha	-	-	12	-	
Botanigard WP	06264	Mycotech Europe Ltd	Insecticide	<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg	0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour		
Scala	03916	BASF plc	Fungicide	<i>Pyrimethanil</i>	400 g/l	2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days	
Serenade ASO	03847	Bayer CropScience Limited	Fungicide	<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l	8 l/ha	48 l/ha	6		0 days	
T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide	<i>Trichoderma Asperellum (strain T34)</i> <i>Trichoderma Asperellum (strain T34)</i>	12% w/w 12% w/w	0.5 g/m2*	0.5 g/m2	1	Immediately after sowing seeds or rooting cuttings		
Tracer	02649	Dow AgroSciences Ltd	Insecticide	<i>Spinosad</i>	480 g/l	200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days	

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Rhubarb</i>											
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.5 l/ha	2 per year	14 days before harvest	14 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide			0.4 l/ha	0.4 l/ha	1 per year	Dormant stage	
					<i>Isoxaben</i>	<i>500 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	PDM 330 EC	03582	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>330 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Rocket</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	3 days pre harvest	3 days
					<i>Acetamiprid</i>	20% w/w					
	Devrinol	05234	UPL Europe Ltd	Herbicide			0.85 l/ha	0.85 l/ha	1 per crop	Pre-planting	
					<i>Napropamide</i>	450 g/l					
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide			250 g/ha	500 g/ha	2	3 days pre harvest	3 days
					<i>Acetamiprid</i>	20 % w/w					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis</i>	375 g/kg					
					<i>subspecies kurstaki EG</i>						
					<i>2348</i>						
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str.</i>	1015.1 g/l					
					<i>QST 713</i>						
<i>Rosemary</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str.</i>	1015.1 g/l					
					<i>QST 713</i>						
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
					<i>Cycloxydim</i>	100 g/l					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	50 %w/w					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Rubus Hybrid - field</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	220 g/kg					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	375 g/kg					
	Masai	03585	BASF plc	Insecticide			375 g/ha	375 g/ha	1 per year	21 days before harvest	21 days
					<i>Tebufenpyrad</i>	200 g/kg					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	2% w/w					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	1015.1 g/l					
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha	-	21 days before harvest	
					<i>Carfentrazone-ethyl</i>	60 g/l					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Rubus Hybrid - protected</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	When leaves begin to discolour	
					<i>Beauveria bassiana</i>	220 g/kg					
					<i>GHA / ATCC 74040</i>						
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis</i>	375 g/kg					
					<i>subspecies kurstaki EG</i>						
					<i>2348</i>						
	Masai	03585	BASF plc	Insecticide			375 g/ha	375 g/ha	1 per year	21 days before harvest	21 days
					<i>Tebufenpyrad</i>	200 g/kg					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium</i>	2% w/w					
					<i>anisoplice - ST</i>						
					<i>BIPESCO 5/F52</i>						
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus</i>	1015.1 g/l					
					<i>amyloliquefaciens str.</i>						
					<i>QST 713</i>						
	Spotlight Plus	03294	FMC Chemical s.p.r.l.	Herbicide			0.8 l/ha	1.6 l/ha	-	21 days before harvest	21 days
					<i>Carfentrazone-ethyl</i>	60 g/l					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m <sup>3</sup>	10 g/m <sup>3</sup>	1	Pre planting / sowing	
					<i>Trichoderma</i>	12% w/w					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	12% w/w					
					<i>Asperellum (strain T34)</i>						

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Rye - spring</i>											
	Compitox Plus	02843	Nufarm UK Limited	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			250 mls/ha	750 mls/ha		Before GS 85	
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	840 mls/ha	2	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1 per crop	Before GS 22	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Duplosan KV	02842	Nufarm UK Limited	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Optica	02661	Nufarm UK Limited	Herbicide			2.0 l/ha	2.0 l/ha	-	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Twinprop	04427	DHM Agrochemicals Ltd.,	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Rye - winter</i>											
	Compitox Plus	02843	Nufarm UK Limited	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Deca 2.5% EC	04952	Sharda Europe b.v.b.a	Insecticide			250 mls/ha	750 mls/ha	-	Before GS 85	
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			250 mls/ha	750 mls/ha	-	Before GS 85	
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	840 mls/ha	2	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1 per crop	Before GS 22	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Duplosan KV	02842	Nufarm UK Limited	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	GAT Decline 2.5 E	04422	FMC Chemical s.p.r.l.	Insecticide			250 mls/ha	750 mls/ha	-	Before GS 85	
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Optica	02661	Nufarm UK Limited	Herbicide			2.0 l/ha	2.0 l/ha	-	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Twinprop	04427	DHM Agrochemicals Ltd.,	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
<i>Sage</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana</i>	<i>220 g/kg</i>					
					<i>GHA / ATCC 74040</i>						
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Salsify</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
	Amistar Top	04582	Syngenta UK Ltd	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide	<i>Azoxystrobin</i> <i>Difenoconazole</i>	<i>200 g/l</i> <i>125 g/l</i>	2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide	<i>Mancozeb</i>	<i>75% w/w</i>	1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
	Azoxystar	05534	Life Scientific Limited	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
	Biscaya	03602	Bayer CropScience Limited	Insecticide	<i>Azoxystrobin</i>	<i>250 g/l</i>	0.4 l/ha	0.8 l/ha	2	14 days before harvest	14 days
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide	<i>Thiacloprid</i>	<i>240 g/l</i>	2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
	Chamane	05002	UPL Europe Ltd	Fungicide	<i>Clethodim</i>	<i>120 g/l</i>	1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
	Corbel	90438	BASF plc	Fungicide	<i>Azoxystrobin</i>	<i>250 g/l</i>	1.0 l/ha	3.0 l/ha	3	28 days before harvest	28 days
	Decis	02693	Bayer CropScience Limited	Insecticide	<i>Fenpropimorph</i>	<i>750 g/l</i>	300 mls/ha	900 mls/ha	3 per crop	21 days before harvest	21 days
	Decis Protech	05269	Bayer CropScience Limited	Insecticide	<i>Deltamethrin</i>	<i>25 g/l</i>	500 mls/ha	1.5 l/ha	3 per crop	21 days before harvest	21 days
	Defy	04277	Syngenta UK Ltd	Herbicide	<i>Deltamethrin</i>	<i>15 g/l</i>	5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide	<i>Prosulfocarb</i>	<i>800 g/l</i>	2.1 kg/ha	8.4 kg/ha	4 per crop	30 days before harvest	30 days
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide	<i>Mancozeb</i> <i>lambda-Cyhalothrin</i>	<i>75% w/w</i> <i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Salsify</i>											
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	8 weeks before harvest	56 days
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
	Globazar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.3 kg/ha	0.9 kg/ha	* 3 per crop	21 days before harvest	21 days
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
					<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	PDM 330 EC	03582	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1		
					<i>Pendimethalin</i>	<i>330 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Salsify</i>											
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 11	80 days
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha		14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Stomp 400 SC	02879	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	8 weeks before harvest	8 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Switch	03761	Syngenta UK Ltd	Fungicide			1.0 kg/ha	3.0 kg/ha	3	7 days before harvest	
					<i>Cyprodinil</i>	<i>37.5 % w/w</i>					
					<i>Fludioxonil</i>	<i>25.0 % w/w</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	3.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Shallot</i>											
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	1.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Basagran SG	05192	BASF plc	Herbicide			1.65 kg/ha	1.65 kg/ha	-	21 days before harvest	21 days
					<i>Bentazone</i>	<i>870 g/kg</i>					
	Bravo 500	03452	Syngenta UK Ltd	Fungicide			2.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	56 days before harvest	56 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Daconil	05748	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			5.0 l/ha	5.0 l/ha	-	Before 5th leaf visible	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Emerger	06190	Bayer CropScience Limited	Herbicide			2.5 l/ha	2.5 l/ha	2 per crop	GS 12	
					<i>Aclonifen</i>	<i>600 g/l</i>					
	Flexidor	04963	Dow AgroSciences Ltd	Herbicide			0.2 l/ha	0.2 l/ha	1 per year	After transplanting or beginning of root formation	
					<i>Isoxaben</i>	<i>500 g/l</i>					
	Folio Gold	02363	Syngenta UK Ltd	Fungicide			2.0 l/ha	6.0 l/ha	-	14 days before harvest	14 days
					<i>Metalaxyl M</i>	<i>37.5 g/l</i>					
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Fusilade Max	01472	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	1	4 weeks before harvest	28 days
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Shallot</i>											
	Infinito	04309	Bayer CropScience Limited	Fungicide			1.6 l/ha	1.6 l/ha	1 per crop	7 days before harvest	7 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Jupital	04503	Syngenta UK Ltd	Fungicide			2.0 l/ha	4.0 l/ha	-	14 days before harvest	14 days
					<i>Chlorothalonil</i>	<i>500 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			75 mls/ha	225 mls/ha	3 per crop	21 days before harvest	21 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Lentagran WP	03082	Belchim Crop Protection NV	Herbicide			2.0 kg/ha	2.0 kg/ha	1	Before 7 leaf stage	
					<i>Pyridate</i>	<i>45 % w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	2.0 l/ha	4 per crop	7 days before harvest	7 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	NEMGuard DE	05368	ECOspray Ltd	Nematicide			20 kg/ha	20 kg/ha	1	At drilling	
					<i>Garlic Extract</i>	<i>450 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			75 mls/ha	225 mls/ha	4	21 days before harvest	21 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Roxy 800 EC	05739	Globachem nv	Herbicide			5.0 l/ha	5.0 l/ha	-	Before GS 15	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Starane Hi-Load H	04549	Dow AgroSciences Ltd	Herbicide			0.3 l/ha	0.3 l/ha		Before GS 15	
					<i>Fluroxypyr</i>	<i>333 g/l</i>					
	Stomp 400 SC	02879	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1	Before crop emergence	
					<i>Pendimethalin</i>	<i>400 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			2.9 l/ha	2.9 l/ha	1 / crop	Before crop emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Valbon	03411	Certis Europe BV	Fungicide			1.6 kg/ha	9.6 kg/ha	6 per crop	28 days pre-harvest	28 days
					<i>Mancozeb</i>	<i>700 g/kg</i>					
					<i>Benthiavalicarb</i>	<i>17.5 g/kg</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Shallot</i>	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	Pre-emergence / Pre-transplanting	
					<i>Pendimethalin</i>	<i>250g/l</i>					
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Spinach</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Bonalan	05763	Gowan Crop Protection Ltd	Herbicide			8.0 l/ha	8.0 l/ha	-	Incorporation pre sowing/planting	
					<i>Benfluralin</i>	<i>180 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Devrinol	05234	UPL Europe Ltd	Herbicide			0.85 l/ha	0.85 l/ha	-	Pre crop emergence	
					<i>Napropamide</i>	<i>450 g/l</i>					
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Infinito	02532	Bayer CropScience Limited	Fungicide			1.6 l/ha	1.6 l/ha	1 per crop	Before GS 49	14 days
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
					<i>Propamocarb</i>	<i>625 g/l</i>					
					<i>Fluopicolide</i>	<i>62.5 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			75 mls/ha	150 mls/ha	4	7 days pre harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Spinach</i>											
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide			75 mls/ha	150 mls/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Markate 50	05001	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	1.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			75 mls/ha	150 mls/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3 mls per sq m	6 mls per sq m	-	Immediately post drilling	
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	7 days before harvest	7 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			0.5 g/m <sup>2</sup> *	0.5 g/m <sup>2</sup>	1	Immediately after sowing seeds or rooting cuttings	
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
					<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 mls/ha	600 mls/ha	3		3 days
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Spinach</i>											
	Venzar 500 SC	04805	DuPont (UK) Ltd	Herbicide			0.8 l/ha	0.8 l/ha	-	Pre-emergence	
					<i>Lenacil</i>	<i>500 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Spinach beet</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Bonalan	05763	Gowan Crop Protection Ltd	Herbicide			8.0 l/ha	8.0 l/ha	-	Incorporation pre sowing/planting	
					<i>Benfluralin</i>	<i>180 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Globaztar SC	05529	Globachem nv	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Met 52 Granular	04811	Novozymes France S.A.S.	Insecticide			0.5 kgs/m <sup>3</sup>	-	-	Pre-planting	
					<i>Metarhizium anisoplice - ST BIPESCO 5/F52</i>	<i>2% w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	1.0 l/ha	2 per crop	7 days before harvest	7 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 mls/ha	600 mls/ha	3		3 days
					<i>Spinosad</i>	<i>480 g/l</i>					
	Venzar 500 SC	04805	DuPont (UK) Ltd	Herbicide			0.8 l/ha	0.8 l/ha	-	Pre-emergence	
					<i>Lenacil</i>	<i>500 g/l</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Squash - protected</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.45 l/ha	0.9 l/ha	2 per year	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Previcur Energy	04041	Bayer CropScience Limited	Fungicide			3 mls per sq m	6 mls per sq m	-	Before transplanting	
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Propamocarb</i>	<i>530 g/l</i>					
					<i>Fosetyl (-aluminium)</i>	<i>310 g/l</i>					
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide			10 g/m3	10 g/m3	1	Pre planting / sowing	
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
					<i>Trichoderma</i>	<i>12% w/w</i>					
					<i>Asperellum (strain T34)</i>						
	Vivando	05685	BASF plc	Fungicide			0.2 l/ha	0.4 l/ha	-	3 days before harvest	3 days
					<i>Metrafenone</i>	<i>500 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Strawberries - field</i>											
	Apollo 50SC	04610	Adama Agricultural Solutions UK Ltd	Acaricide			0.4 l/ha	0.4 l/ha	1 per year	9 days before harvest	9 days
					<i>Clofentezine</i>	<i>500 g/l</i>					
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.50 l/ha	2 per year	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Chess WG	03365	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kg/ha	1.2 kg/ha	3 per crop	3 days before harvest	3 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			500 ml/ha	1.0 l/ha	2 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			0.7 l/ha	1.3 l/ha	-	30 days before harvest	30 days
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			0.7 l/ha	1.3 l/ha	-	30 days before harvest	30 days
					<i>S-Metolachlor</i>	<i>960 g/l</i>					
	Fortress	02364	Dow AgroSciences Ltd	Fungicide			0.25 l/ha	0.5 l/ha	2	14 days before harvest	14 days
					<i>Quinoxifen</i>	<i>500 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			125 ml/ha	250 ml/ha	4 per crop	3 days before harvest	3 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 g/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
	Nimrod	90789	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	3	3 days before harvest	3 days
					<i>Bupirimate</i>	<i>250 g/l</i>					
	Takumi SC	06060	Nisso Chemical Europe GmbH	Fungicide			0.15 l/ha	0.3 l/ha	2 per crop	3 days before harvest	3 days
					<i>Cyflufenamid</i>	<i>100 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product</i>					<i>Max Ind.</i>	<i>Max. Total</i>	<i>Max No of</i>	<i>Latest Time</i>	
<i>Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Dose:</i>	<i>Dose:</i>	<i>Applic's:</i>	<i>of Applic:</i>	<i>PHI:</i>
<i>Strawberries - field</i>										
Tracer	02649	Dow AgroSciences Ltd	Insecticide			150 ml/ha in 200L water	450 ml/ha	3 per year	1 day before harvest	1 day
				<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Strawberries - protected</i>											
	Apollo 50SC	04610	Adama Agricultural Solutions UK Ltd	Acaricide			0.4 l/ha	0.4 l/ha	1 per year	9 days before harvest	9 days
					<i>Clofentezine</i>	<i>500 g/l</i>					
	Benevia 10 OD	05354	Headland Agrochemicals Ltd	Insecticide			0.75 l/ha	1.5 l/ha	2	1 day before harvest	1 day
					<i>Cyantraniliprole</i>	<i>100 g/l</i>					
	Borneo	04235	Sumitomo Chemical Agro Europe SAS	Acaricide			500mls/ha - max conc. 35 mls/100 litres water	500mls/ha - max conc. 35 mls/100 litres water	1/crop	3 days before harvest	3 days
					<i>Etoxazole</i>	<i>110 g/l</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.25 l/ha	0.50 l/ha	2 per crop	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Chess WG	03365	Adama Agricultural Solutions UK Ltd	Insecticide			0.4 kg/ha	1.2 kg/ha	3 per crop	3 days before harvest	3 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Dynamec	02640	Syngenta UK Ltd	Acaricide/Insecticid			50 mls/product/100 litres water	-	3	3 days before harvest	3 days
					<i>Abamectin</i>	<i>18 g/l</i>					
	Fortress	02364	Dow AgroSciences Ltd	Fungicide			0.25 l/ha	0.5 l/ha	2	14 days before harvest	14 days
					<i>Quinoxifen</i>	<i>500 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			75 ml/ha	150 ml/ha	4 per crop	3 days before harvest	3 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Masai	03585	BASF plc	Insecticide			375 gms/ha	375 g/ha	1 / year	21 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
	Microthiol Special	90548	UPL Europe Ltd	Fungicide							
					<i>Sulphur</i>	<i>80 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Strawberries - protected</i>											
	Nimrod	90789	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	3	3 days before harvest	3 days
	Sabik-1.8%	04907	Servem Ltd	Acaricide/Insecticid	<i>Bupirimate</i>	<i>250 g/l</i>	50 mls / 100 l water	4.0 l/ha	3	3 days before harvest	3 days
	Sequoia	05915	Dow AgroSciences Ltd	Insecticide	<i>Abamectin</i>	<i>18 g/l</i>	200 mls/ha	400 mls/ha	2	1 day before harvest	1 day
	T34 Biocontrol	05620	Biocontrol Technologies S.L.	Fungicide	<i>Sulfoxaflor</i>	<i>120 g/l</i>	10 g/m3	10 g/m3	1	Pre planting / sowing	
	Takumi SC	06060	Nisso Chemical Europe GmbH	Fungicide	<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>	0.15 l/ha	0.3 l/ha	2 per crop	3 days before harvest	3 days
	Talius	04604	DuPont (UK) Ltd	Fungicide	<i>Trichoderma Asperellum (strain T34)</i>	<i>12% w/w</i>	0.19 l/ha	0.19 l/ha	1 per crop	3 days before harvest	3 days
					<i>Cyflufenamid</i>	<i>100 g/l</i>					
					<i>Proquinazid</i>	<i>200 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Swedes</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Benevia 10 OD	05354	Headland Agrochemicals Ltd	Insecticide			0.75 l/ha	1.5 l/ha	2 per crop	14 days before harvest	14 days
					<i>Cyantraniliprole</i>	<i>100 g/l</i>					
	Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	35 days before harvest	35 days
					<i>Thiacloprid</i>	<i>240 g/l</i>					
	Bonalan	05763	Gowan Crop Protection Ltd	Herbicide			8.0 l/ha	8.0 l/ha	-	Incorporation pre sowing/planting	
					<i>Benfluralin</i>	<i>180 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	14 days pre harvest	14 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide			1 kg/ha	6 kg/ha	-	8 days before harvest	8 days
					<i>Copper oxychloride</i>	<i>87.8% w/w</i>					
	Devrinol	05234	UPL Europe Ltd	Herbicide			1.25 l/ha	1.25 l/ha	-	Pre crop emergence	
					<i>Napropamide</i>	<i>450 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					

**Warning EAMU approvals:**

***The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.***

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Swedes</i>											
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide	<i>S-Metolachlor</i>	<i>960 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Fortress	02364	Dow AgroSciences Ltd	Fungicide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	0.3 l/ha	0.4 l/ha	2	28 days pre harvest	28 days
	Gamit 36 CS	05357	FMC Chemical s.p.r.l.	Herbicide	<i>Quinoxifen</i>	<i>500 g/l</i>	0.25 l/ha	0.25 l/ha	-	pre-emergence	
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide	<i>Clomazone</i>	<i>360 g/l</i>	250 g/ha	500 g/ha	2 per crop	14 days before harvest	14 days
	Globaztar SC	05529	Globachem nv	Fungicide	<i>Acetamiprid</i>	<i>20 % w/w</i>	1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide	<i>Azoxystrobin</i>	<i>250 g/l</i>	150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	1.0 kg/ha	3.0 kg/ha		N/A	
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide	<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>	75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
	Markate 50	05001	Globachem nv	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide	<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>	0.5 l/ha	0.5 l/ha	-	28 days before harvest	28 days
	Movento	05349	Bayer CropScience Limited	Insecticide	<i>Difenoconazole</i>	<i>250 g/l</i>	0.5 l/ha	1.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Swedes</i>											
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide			0.36 kg/ha	0.72 kg/ha	* 2 per crop	21 days before harvest	21 days
					<i>Trifloxystrobin</i>	<i>250 g/kg</i>					
					<i>Tebuconazole</i>	<i>500 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha		28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Venzar 500 SC	04805	DuPont (UK) Ltd	Herbicide			0.4 l/ha	0.4 l/ha	-	Pre-emergence	
					<i>Lenacil</i>	<i>500 g/l</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Sweetcorn</i>											
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide			100 ml/ha	400 ml/ha	4	7 days before harvest	
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Lepinox Plus	05254	CBC (Europe) S.r.l.	Insecticide			1.0 kg/ha	3.0 kg/ha	-	N/A	
					<i>Bacillus thuringiensis subspecies kurstaki EG 2348</i>	<i>375 g/kg</i>					
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide			100 ml/ha	400 ml/ha	4	7 days before harvest	
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Stomp Aqua	03750	BASF plc	Herbicide			3.3 l/ha	3.3 l/ha	1 per crop	Pre-emergence	
					<i>Pendimethalin</i>	<i>455 g/l</i>					
	Wing P	03590	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	-	Pre-emergence	
					<i>Dimethenamid-P</i>	<i>212.5 g/l</i>					
					<i>Pendimethalin</i>	<i>250g/l</i>					
<i>Thyme</i>											
	Botanigard WP	06264	Mycotech Europe Ltd	Insecticide			0.6 kg/ha	3.0 kg/ha	5 per year	-	
					<i>Beauveria bassiana GHA / ATCC 74040</i>	<i>220 g/kg</i>					
	Stratos Ultra	04936	BASF plc	Herbicide			4.0 l/ha	4.0 l/ha	1 per crop	4 weeks before harvest	4 weeks
					<i>Cycloxydim</i>	<i>100 g/l</i>					
	Teldor	02003	Bayer CropScience Limited	Fungicide			1.5 kgs/ha	3.0 kgs/ha		3 days before harvest	3 days
					<i>Fenhexamid</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Tomatoes - protected</i>											
	Acrobat MZ	04059	BASF plc	Fungicide			2.4 kg/ha	7.2 kg/ha	-	3 days before harvest	3 days
					<i>Dimethomorph</i>	<i>75 g/kg</i>					
					<i>Mancozeb</i>	<i>667 g/kg</i>					
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Difenoconazole</i>	<i>125 g/l</i>					
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
	AVTAR 75 NT	06297	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	6.3 kg/ha	-	3 days before harvest	3 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	CaLypso	02441	Bayer CropScience Limited	Insecticide			0.45 l/ha	0.9 l/ha	2 per crop	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
	Cerone	03933	Bayer CropScience Limited	Growth Regulator			100 ml in 100 litre water. To run off	100 ml in 100 litre water. To run off	1 per crop	5 days before harvest	5 days
					<i>Ethephon</i>	<i>480 g/l</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Coragen	05295	DuPont (UK) Ltd	Insecticide			250 ml/ha	500 ml/ha	-	1 day before harvest	1 day
					<i>Chlorantraniliprole</i>	<i>200 g/l</i>					
	Curzate M WG	04723	DuPont (UK) Ltd	Fungicide			2.3 kg/ha	6.9 kg/ha	-	3 days before harvest	3 days
					<i>Mancozeb</i>	<i>68 % w/w</i>					
					<i>Cymoxanil</i>	<i>4.5 % w/w</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			830 ml/ha	2.49 l/ha	3 per crop	3 days before harvest	3 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Tomatoes - protected</i>											
	Dithane 945	03705	Indofil Industries (Netherlands) B.V	Fungicide			2.0 kg/ha	6.0 kg/ha	-	3 days before harvest	3 days
					<i>Mancozeb</i>	<i>80 % w/w</i>					
	Dithane Dry Flowa	03707	Indofil Industries (Netherlands) B.V	Fungicide			2.1 kg/ha	6.3 kg/ha	-	3 days before harvest	3 days
					<i>Mancozeb</i>	<i>75% w/w</i>					
	Electis 75 WG	01821	Gowan Crop Protection Ltd	Fungicide			1.8 kg/ha	5.4 kg/ha	3 per crop	3 days before harvest	3 days
					<i>Zoxamide</i>	<i>8.3 %w/w</i>					
					<i>Mancozeb</i>	<i>66.7 %w/w</i>					
	Globazar SC	05529	Globachem nv	Fungicide			1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Karma	05116	Certis Europe BV	Fungicide			3.0 kg/ha	24.0 kg/ha	8 per crop	1 day before harvest	1 day
					<i>Potassium Hydrogen Carbonate</i>	<i>850 g/kg</i>					
	Luna Privilege	05916	Bayer CropScience Limited	Fungicide			1.0 l/ha	4.0 l/ha	4	1 day before harvest	1 day
					<i>Fluopyram</i>	<i>500 g/l</i>					
	Manfil 75 WG	03598	Indofil Industries Limited	Fungicide			2.1 kg/ha	6.3 kg/ha	-	3 days before harvest	3 days
					<i>Mancozeb</i>	<i>75 % w/w</i>					
	Manfil 80 WP	03563	Indofil Industries Limited	Fungicide			2.0 kg/ha	4.0 kg/ha	-	3 days before harvest	3 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
	Microthiol Special	90548	UPL Europe Ltd	Fungicide			0.01kgs/5 litres water	-	-	-	0 days
					<i>Sulphur</i>	<i>80 %w/w</i>					
	Movento	05349	Bayer CropScience Limited	Insecticide			0.5 l/ha	2.0 l/ha	4 per crop	14 days before harvest	14 days
					<i>Spirotetramat</i>	<i>150 g/l</i>					
	Nimrod	90789	Adama Agricultural Solutions UK Ltd	Fungicide			0.2L per 100L water	-	6 per crop	2 days before harvest	2 days
					<i>Bupirimate</i>	<i>250 g/l</i>					
	Option	04722	DuPont (UK) Ltd	Fungicide			0.4 kg/ha	1.6 kg/ha	4 per crop	3 days before harvest	3 days
					<i>Cymoxanil</i>	<i>600 g/kg</i>					
	Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide			0.06 kgs/100 litres water	-	3	3 days before harvest	3 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Tomatoes - protected</i>											
Revus	05154	Syngenta UK Ltd	Fungicide				0.6 l/ha	2.4 l/ha	4 per crop	3 days before harvest	3 days
					<i>Mandipropamid</i>	<i>250 g/l</i>					
Ridomil Gold MZ 6	01516	Syngenta UK Ltd	Fungicide				2.5 kg/ha	7.5 kg/ha	-	3 days before harvest	3 days
					<i>Metalaxyl M</i>	<i>4.0 %w/w</i>					
					<i>Mancozeb</i>	<i>64 %w/w</i>					
Scala	03916	BASF plc	Fungicide				0.5 l/ha	1.0 l/ha	2 per crop	3 days before harvest	3 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
SL567A	03106	Syngenta UK Ltd	Fungicide				0.3 l/ha	0.9 l/ha	3 per crop		3 days
					<i>Metalaxyl M</i>	<i>465.2 g/l</i>					
Teldor	02003	Bayer CropScience Limited	Fungicide				1.5 kgs/ha	4.5 kgs/ha	3 / crop	1 day before harvest	1 day
					<i>Fenhexamid</i>	<i>50 %w/w</i>					
Vivando	05685	BASF plc	Fungicide				0.45 l/ha	0.9 l/ha	-	3 days before harvest	3 days
					<i>Metrafenone</i>	<i>500 g/l</i>					
Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide				1.0 l/ha	3.0 l/ha	-	3 days before harvest	3 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Triticale</i>											
	Barclay Bolt XL	04102	Barclay Chemicals Ltd	Fungicide			0.5 l/ha	1.0 l/ha	2 per crop	Before GS 71	
					<i>Propiconazole</i>	<i>250 g/l</i>					
	Barclay Propizole	04099	Barclay Chemicals Ltd	Fungicide			0.5 l/ha	1.0 l/ha	2 per crop	Before GS 71	
					<i>Propiconazole</i>	<i>250 g/l</i>					
	Bumper	03292	Adama Agricultural Solutions UK Ltd	Fungicide			0.5 l/ha	1.0 l/ha	2 per crop	Before GS 71	35 days
					<i>Propiconazole</i>	<i>250 g/l</i>					
	Cliophar 600 SL	04864	Arysta LifeScience Benelux SPRL	Herbicide			0.12 l/ha	0.12 l/ha	-	Before GS 33	
					<i>Clopyralid</i>	<i>600 g/l</i>					
	Compitox Plus	02843	Nufarm UK Limited	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Deca 2.5% EC	04952	Sharda Europe b.v.b.a	Insecticide			250 mls/ha	750 mls/ha	-	Before GS 85	
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis	02693	Bayer CropScience Limited	Insecticide			250 mls/ha	750 mls/ha		before GS 85	
					<i>Deltamethrin</i>	<i>25 g/l</i>					
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	840 mls/ha	2	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
	Defy	04277	Syngenta UK Ltd	Herbicide			3.0 l/ha	3.0 l/ha	-	Before GS22	
					<i>Prosulfocarb</i>	<i>800 g/l</i>					
	Duplosan KV	02842	Nufarm UK Limited	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Empire Supreme	05061	Croplink (Ireland) Ltd	Herbicide			100 g/ha	100 g/ha	1 per crop	Before GS 39	
					<i>Thifensulfuron methyl</i>	<i>400 g/kg</i>					
					<i>Metsulfuron (-methyl)</i>	<i>40 g/kg</i>					
	Finish SX	02329	DuPont (UK) Ltd	Herbicide			75 g/ha	75 g/ha	1	Before GS 39	
					<i>Metsulfuron (-methyl)</i>	<i>67 g/kg</i>					
					<i>Thifensulfuron methyl</i>	<i>333 g/kg</i>					
	GAT Decline 2.5 E	04422	FMC Chemical s.p.r.l.	Insecticide			250 mls/ha	750 mls/ha	-	Before GS 85	
					<i>Deltamethrin</i>	<i>25 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Triticale</i>											
	Harmony M SX	02330	DuPont (UK) Ltd	Herbicide			100 g/ha	100 g/ha	1	Before GS 39	
					<i>Metsulfuron (-methyl)</i>	<i>40 g/kg</i>					
					<i>Thifensulfuron methyl</i>	<i>400 g/kg</i>					
	Optica	02661	Nufarm UK Limited	Herbicide			2.0 l/ha	2.0 l/ha	-	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					
	Presite SX	02888	DuPont (UK) Ltd	Herbicide			75 g/ha	75 g/ha	1	Before GS 39	
					<i>Thifensulfuron methyl</i>	<i>333 g/kg</i>					
					<i>Metsulfuron (-methyl)</i>	<i>67 g/kg</i>					
	Propi 25 EC	04205	Sharda Europe b.v.b.a	Fungicide			0.5 l/ha	1.0 l/ha	2 per crop	Before GS 71	
					<i>Propiconazole</i>	<i>250 g/l</i>					
	Twinprop	04427	DHM Agrochemicals Ltd.,	Herbicide			2.3 l/ha	2.3 l/ha	1 per crop	Before GS 33	
					<i>Mecoprop-P</i>	<i>600 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Turnip</i>											
	Amistar	05072	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Amistar Top	04582	Syngenta UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>200 g/l</i>					
					<i>Difenoconazole</i>	<i>125 g/l</i>					
	Azoshy	05618	Sharda Cropchem Espana S.L.	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Azoxystar	05534	Life Scientific Limited	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Benevia 10 OD	05354	Headland Agrochemicals Ltd	Insecticide			0.75 l/ha	1.5 l/ha	2 per crop	14 days before harvest	14 days
					<i>Cyantraniliprole</i>	<i>100 g/l</i>					
	Biscaya	03602	Bayer CropScience Limited	Insecticide			0.4 l/ha	0.8 l/ha	2	35 days before harvest	35 days
					<i>Thiacloprid</i>	<i>240 g/l</i>					
	Bonalan	05763	Gowan Crop Protection Ltd	Herbicide			8.0 l/ha	8.0 l/ha	-	Incorporation pre sowing/planting	
					<i>Benfluralin</i>	<i>180 g/l</i>					
	Centurion Max	04948	Arysta Life Science S.A.S.	Herbicide			2.0 l/ha	2.0 l/ha	-	48 days before harvest	48 days
					<i>Clethodim</i>	<i>120 g/l</i>					
	Ceta SG	06252	Unichem Limited	Insecticide			250 g/ha	500 g/ha	2 per crop	14 days pre harvest	14 days
					<i>Acetamiprid</i>	<i>20% w/w</i>					
	Chamane	05002	UPL Europe Ltd	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
	Curenox 50 WP	05829	Industrias Quimicas Del Valles S.A (IQV, S.A.)	Fungicide			1 kg/ha	6 kg/ha	-	8 days before harvest	8 days
					<i>Copper oxychloride</i>	<i>87.8% w/w</i>					
	Devrinol	05234	UPL Europe Ltd	Herbicide			1.25 l/ha	1.25 l/ha	-	Pre crop emergence	
					<i>Napropamide</i>	<i>450 g/l</i>					
	Dual Gold	04192	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 leaf stage	
					<i>S-Metolachlor</i>	<i>960 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**



<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Turnip</i>											
	Efica 960 EC	05942	Syngenta UK Ltd	Herbicide			1.4 l/ha	1.4 l/ha	-	Up to 4 true leaf stage	
	Eminentos 10 CS	04933	FMC Chemical s.p.r.l.	Insecticide	<i>S-Metolachlor</i>	<i>960 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Fortress	02364	Dow AgroSciences Ltd	Fungicide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	0.3 l/ha	0.4 l/ha	2	28 days pre harvest	28 days
	Gazelle SG	03248	Nisso Chemical Europe GmbH	Insecticide	<i>Quinoxifen</i>	<i>500 g/l</i>	250 g/ha	500 g/ha	2 per crop	14 days before harvest	14 days
	Globazar SC	05529	Globachem nv	Fungicide	<i>Acetamiprid</i>	<i>20 % w/w</i>	1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
	Karate Zeon	04084	Syngenta UK Ltd	Insecticide	<i>Azoxystrobin</i>	<i>250 g/l</i>	150 mls/ha	450 mls/ha	4	14 days pre harvest	14 days
	Karis 10 CS	04454	FMC Chemical s.p.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Laidir 10 CS	04932	FMC Chemical s.p.r.l.	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
	Life Scientific Lam	04868	Life Scientific Limited	Insecticide	<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>	75 mls/ha	150 mls/ha	4	14 days before harvest	14 days
	Markate 50	05001	Globachem nv	Insecticide	<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>	75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
	Mavita 250 EC	05941	Syngenta UK Ltd	Fungicide	<i>Difenoconazole</i>	<i>250 g/l</i>	0.5 l/ha	0.5 l/ha	-	28 days before harvest	28 days
	Movento	05349	Bayer CropScience Limited	Insecticide	<i>Spirotetramat</i>	<i>150 g/l</i>	0.5 l/ha	1.0 l/ha	2 per crop	21 days before harvest	21 days
	Nativo 75 WG	04642	Bayer CropScience Limited	Fungicide	<i>Trifloxystrobin</i>	<i>250 g/kg</i>	0.36 kg/ha	0.72 kg/ha	* 2 per crop	21 days before harvest	21 days
	Ninja 10CS	05178	Syngenta UK Ltd	Insecticide	<i>Tebuconazole</i>	<i>500 g/kg</i>	150 mls/ha	450 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Turnip</i>											
	Pilot Ultra	04955	Nissan Chemical Europe	Herbicide			2.5 l/ha	2.5 l/ha	-	60 days before harvest	60 days
					<i>Quizalofop-P-ethyl</i>	<i>50 g/l</i>					
	Ravane	05536	Globachem nv	Insecticide			75 ml/ha	150 ml/ha	4 per crop	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>50 g/l</i>					
	Reflect	05462	Adama Agricultural Solutions UK Ltd	Fungicide			1.0 l/ha	2.0 l/ha	2 per crop	14 days before harvest	14 days
					<i>Isopyrazam</i>	<i>125 g/l</i>					
	Score 250 EC	04566	Syngenta UK Ltd	Fungicide			0.5 l/ha	0.5 l/ha		28 days before harvest	28 days
					<i>Difenoconazole</i>	<i>250 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Sparviero	04469	Sipcam Oxon SpA	Insecticide			75 ml/ha	150 ml/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
	Venzar 500 SC	04805	DuPont (UK) Ltd	Herbicide			0.4 l/ha	0.4 l/ha	-	Pre-emergence	
					<i>Lenacil</i>	<i>500 g/l</i>					
	Vydate 10G	03698	DuPont (UK) Ltd	Nematicide			0.09 kg/100 m row	0.09 kg/100 m row	1 per crop	At drilling	
					<i>Oxamyl</i>	<i>100 g/kg</i>					
	Zoxis	05348	Arysta LifeScience Benelux SPRL	Fungicide			1.0 l/ha	2.0 l/ha	-	14 days before harvest	14 days
					<i>Azoxystrobin</i>	<i>250 g/l</i>					
<i>Wheat - durum</i>											
	Decis Protech	05269	Bayer CropScience Limited	Insecticide			420 mls/ha	840 mls/ha	2	30 days before harvest	30 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Whitecurrants - field</i>											
CaLypso	02441	Bayer CropScience Limited	Insecticide				0.25 l/ha	0.75 l/ha	3 per year	3 days before harvest	3 days
					<i>Thiacloprid</i>	<i>480 g/l</i>					
Chikara Weed Cont	04203	ISK Biosciences Europe N.V.	Herbicide				120 g/ha	120 g/ha	1 per year	Before bud burst	
					<i>Flazasulfuron</i>	<i>25 % w/w</i>					
Decis Protech	05269	Bayer CropScience Limited	Insecticide				400 ml/ha	800 ml/ha	2 per crop	7 days before harvest	7 days
					<i>Deltamethrin</i>	<i>15 g/l</i>					
Fusilade Max	01472	Syngenta UK Ltd	Herbicide				3.0 l/ha	3.0 l/ha	1	Before flowering or after harvest	
					<i>Fluazifop-P (-butyl)</i>	<i>125 g/l</i>					
Karate Zeon	04084	Syngenta UK Ltd	Insecticide				100 mls/ha	200 mls/ha	4	14 days pre harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
Manfil 75 WG	03598	Indofil Industries Limited	Fungicide				1.5 kg per 1000 litres water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>75 % w/w</i>					
Manfil 80 WP	03563	Indofil Industries Limited	Fungicide				1.4 kg per 1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
Masai	03585	BASF plc	Insecticide				1.5 kgs/ha	1.5 kgs/ha	-	7 days before harvest	
					<i>Tebufenpyrad</i>	<i>200 g/kg</i>					
Ninja 10CS	05178	Syngenta UK Ltd	Insecticide				100 mls/ha	200 mls/ha	4	14 days before harvest	14 days
					<i>lambda-Cyhalothrin</i>	<i>100 g/l</i>					
Penncozeb 80 WP	03609	UPL Europe Ltd	Fungicide				1.4 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>800 g/kg</i>					
Penncozeb WDG	03608	UPL Europe Ltd	Fungicide				1.5 kg/1000L water	-	-	28 days before harvest	28 days
					<i>Mancozeb</i>	<i>750 g/kg</i>					
Plenum	02238	Adama Agricultural Solutions UK Ltd	Insecticide				0.4 kgs/ha	1.2 kgs/ha	3	28 days pre harvest	28 days
					<i>Pymetrozine</i>	<i>50 %w/w</i>					
Scala	03916	BASF plc	Fungicide				2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					

**Warning EAMU approvals:**

**The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.**

<i>Crop</i>	<i>Product Name</i>	<i>PCS No</i>	<i>Company Name</i>	<i>Function</i>	<i>Substance</i>	<i>Content</i>	<i>Max Ind. Dose:</i>	<i>Max. Total Dose:</i>	<i>Max No of Applic's:</i>	<i>Latest Time of Applic:</i>	<i>PHI:</i>
<i>Whitecurrants - field</i>											
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Signum	03799	BASF plc	Fungicide			1.5 kgs/ha	4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest)	-	3 days before harvest, N/A for post harvest	
					<i>Boscalid</i>	<i>26.7 % w/w</i>					
					<i>Pyraclostrobin</i>	<i>6.7 % w/w</i>					
	Stroby WG	05092	BASF plc	Fungicide			0.2 kg/ha	0.6 kg/ha	-	14 days before harvest	14 days
					<i>Kresoxim Methyl</i>	<i>50% w/w</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha in 250L water	400 ml/ha	2 per year	7 days before harvest	7 days
					<i>Spinosad</i>	<i>480 g/l</i>					
<i>Whitecurrants - protected</i>											
	AQ 10	05801	CBC (Europe) S.r.l.	Fungicide			70 g/ha	-	12	-	
					<i>Ampelomyces Quisqualis</i>	<i>58 %w/w</i>					
	Scala	03916	BASF plc	Fungicide			2.0 l/ha	4.0 l/ha	2 per crop	21 days before harvest	21 days
					<i>Pyrimethanil</i>	<i>400 g/l</i>					
	Serenade ASO	03847	Bayer CropScience Limited	Fungicide			8 l/ha	48 l/ha	6		0 days
					<i>Bacillus amyloliquefaciens str. QST 713</i>	<i>1015.1 g/l</i>					
	Tracer	02649	Dow AgroSciences Ltd	Insecticide			200 ml/ha in 250L water	400 ml/ha	2 per year	3 days before harvest	3 days
					<i>Spinosad</i>	<i>480 g/l</i>					

**Warning EAMU approvals:**

*The use of any product in accordance with an EAMU approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.*