

Apply CREDO under good growing conditions with adequate soil moisture. Avoid poor growing conditions, which may give less reliable results.

TIMING

CREDO controls diseases most effectively when diseases are at low levels and before they spread to new growth. Application can be repeated after 3-4 weeks if necessary.

Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made as a protectant treatment or in the earliest stages of disease development following a disease risk assessment or the use of appropriate decision support systems.

DOSE

Apply CREDO at 2.0 litres per hectare. Use CREDO as part of a disease management programme. Control of many diseases may be improved by using with an appropriate triazole mixture partner. This is also good practice for resistance management.

VOLUME AND APPLICATION

Apply in 200-300 litres of water per hectare using a medium quality spray (BCPC) at a pressure of at least 2 Bar. Apply through conventional crop spraying equipment. The higher volumes are recommended when the crop is dense.

MIXING

CREDO containers should be shaken well before use. CREDO mixes easily with water and should be added directly to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation during filling and when spraying.

SPRAYER CLEANING

Thoroughly wash out sprayer according to manufacturer's guidelines and dispose of washings and clean containers according to DEFRA Code of Practice (UK only) and local water authority guidelines.

NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

All trademarks duly acknowledged.

Section 6 of the Health and Safety at Work Act
Additional Product Safety Information
(This section does not form part of the product label under the Control of Pesticide Regulations 1986)

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has "off-label" approval or is otherwise permitted under the Control of Pesticides Regulations.

The information on this label is based on the best available information including data from test results.

IMPORTANT INFORMATION			
FOR USE ONLY AS A PROFESSIONAL FUNGICIDE			
For use on:	Maximum individual dose (L/ha)	Maximum number of treatments (per crop)	Latest time of application
wheat	2.0	2	GS71
barley	2.0	2	GS51
Method of application: Tractor mounted sprayer Other specific restrictions: <ul style="list-style-type: none">• The container must not be re-used for any purpose.• To reduce the risk of resistance developing in target diseases the total number of applications of products containing QoI fungicides made to any cereal crop must not exceed two.			
READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR PLANT PROTECTION PRODUCTS			

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work (UK only).

Du Pont (U.K.) Limited, Crop Protection Products,
4th Floor, Kings Court, London Road, Stevenage,
Herts. SG1 2NG Technical enquiries Tel: 01438 734450
or enquiry.agproducts@dupont.com
Emergency 24-hour contact Tel: 0870 820 0418(UK) or 01 901 4670 (Ireland)
National Poisons Information Service; 111 (England), 08454 24 24 24 (Scotland)
and 01 837 9964 (Ireland)

SHAKE WELL BEFORE USE. PROTECT FROM FROST FOR PROFESSIONAL USE ONLY

Manufactured in the EU


K-41224/31602 - UK/IE

Batch No.: see on the pack

To access the Safety Data Sheet for this product scan the code, visit www.cropprotection.dupont.co.uk or alternatively contact your supplier

3 576960 035934

The Voluntary Initiative
This label is compliant with the CPA Voluntary Initiative Guidance (UK only)



MAPP 14577
PCS 02503

A suspension concentrate containing 100 g/l picoxystrobin and 500 g/l chlorothalonil for use as a broad spectrum fungicide for disease control in wheat and barley.

® Registered trademark of E. I. du Pont de Nemours and Company

Contents: 5 litres e

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment (UK only):
WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or contaminated surfaces.
However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection (UK only).
AVOID CONTACT WITH SKIN AND EYES.
WASH ALL PROTECTIVE CLOTHING thoroughly after use especially the insides of gloves.
WASH SPLASHES from skin or eyes immediately.
WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work

Environmental protection

(UK only) DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from hand held sprayers to fall within 1 m of the top of the bank of a static or flowing waterbody. Aim spray away from water. This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme. Before each spraying operation from a horizontal boom sprayer or broadcast air assisted sprayer either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.
(Ireland only) To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies
DO NOT CONTAMINATE surface waters and ditches with chemical or used container.

Storage and disposal

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.
RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add the washings to the sprayer tank at the time of filling and dispose of container safely.




RISK AND SAFETY INFORMATION

CREDO®
Fungicide containing 100g/l picoxystrobin and 500g/l chlorothalonil

Harmful if inhaled
May cause an allergic skin reaction
May cause respiratory irritation
Suspected of causing cancer
Very toxic to aquatic life with long lasting effects

Avoid breathing spray.
Wear protective gloves / protective clothing.
IF ON SKIN: Wash with plenty of soap and water.
IF exposed or concerned: Get medical advice/attention.
Wash contaminated clothing before reuse.
Collect spillage.
Store in a well-ventilated place. Keep container tightly closed.
Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste
Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).
To avoid risks to human health and the environment, comply with the instructions for use

MAPP 14577
PCS 02503



OPEN HERE

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

Avoid application to any crop suffering from stress as a result of drought, waterlogging, low temperatures, nutrient or lime deficiency or other factors reducing crop growth.
Avoid application in either frosty or hot, sunny weather conditions.

DISEASE CONTROL

CREDO contains chlorothalonil (a phthalonitrile fungicide) and picoxystrobin (a QoI fungicide) and controls a range of diseases of wheat and barley. At the recommended rate of application, ‘CREDO’ is active against the following:

Crop	Disease	Level of control	
		UK	Ireland
Wheat	Septoria leaf blotch	C	MC
	Glume blotch	C	-
	Yellow rust	MC	MC
	Brown Rust	C	C
	Tan spot	C	C
Barley	Net Blotch	C	C
	Brown Rust	C	
	<i>Rhynchosporium</i>	C	C
	Powdery Mildew	-	MC

Key
C = control, MC = moderate control

Barley spotting control is also achieved when used as part of a programme with other fungicides.
Disease control is generally more effective if CREDO is applied at an early stage of disease development.

DISEASE RESISTANCE

CREDO contains fungicides from the QoI and chloronitrile groups. CREDO should be used preventatively and should not be relied on for its curative potential. Use CREDO as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action
When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. A pathogen is considered resistant to a fungicide if it survives a correctly applied treatment at the recommended dose and timing, under normal weather conditions. Development of resistance within a pathogen can be avoided or delayed by alternating or mixing products having a different mode of action. The Fungicide Resistance Action Group (FRAG) has produced guidelines on the management and avoidance of fungicide resistance. Details of these can be obtained from your supplier or DuPont.
There is significant risk of widespread resistance occurring in Septoria populations in the UK and Ireland. Failure to follow resistance management action may result in reduced levels of disease control. For further advice on resistance management, contact your agronomist, specialist advisor or TEAGASC and visit FRAG – UK website. You must not apply more than two foliar applications of QoI containing products to any cereal crop.

CROPS

CREDO can be used on all winter and spring varieties of wheat and barley.

SPECIMEN -
2016 to date