# Comet<sup>®</sup>200

A protectant fungicide with curative properties for use in winter wheat, winter barley, spring wheat, spring barley, winter oats and spring oats, rye and triticale.

An emulsifiable concentrate containing 200 g/litre (19.2% w/w) pyraclostrobin. Pyraclostrobin is a strobilurin fungicide.

Risk and Safety Information

#### Danger

Harmful if swallowed. May be fatal if swallowed and enters airways.

Causes skin irritation.

May cause an allergic skin reaction. Causes serious eve irritation.

Harmful if inhaled. May cause respiratory irritation.

Very toxic to aquatic life with long lasting effects.

Use only outdoors or in a well-ventilated area.

Wear protective gloves/protective clothing/eye protection/face protection

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenges, if present and easy to do. Continue rinsing. Call a POISON CENTRE or doctor/physician if you feel unwell.

Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container to a licensed hazardous waste disposal contrar of or collection site except for triple

rinsed empty containers which can be disposed of as non-hazardous waste. To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. : 03696

UFT - BCU1-80U9-0008-0VN2

Supplied by: BASF Ireland Limited, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, UK, Telephone: 00 44 (0)161 475 3000 Emergency Information (24 hours freephone): 0049 180 227 3112 Technical Enquiries: 0044 (0)845 602 2553 (office hours)

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, UK, Telephone: 00 44 (0)161 475 3000

BASF

φ

trademark

Registered

#### FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

l	Crops	Maximum individual dose	Maximum number of applications	Latest time of application	
	Winter and spring wheat, rye and triticale	1.25 litre product per hectare	2 per season	Before watery ripe stage (GS 71)	
	Winter and spring barley, winter and spring oats	1.25 litre product per hectare	2 per season	Before end of heading (GS 59)	

#### **READ ALL PRECAUTIONS BEFORE USE**

PCS No.: 03696

#### **PRECAUTIONS**

For professional use only

WEAR SUITABLE PROTECTIVE GLOVES when handling the corce trate

WASH HANDS AND EXPOSED SKIN before meals and after york

WHEN USING DO NOT FAT, DRINK OR SMOKE.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING S UFFS.

VERY TOXIC TO AQUATIC ORGANISMS, may cause long-term adverse effects in the aquatic environment.

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

IF YOU FEEL UNWELL, seek medical advice (show label where possible).

#### STORAGE

Store in a suitable pesticide store. Keep dry and protect from frost.

#### **DIRECTIONS FOR USE**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Apply as a MEDIUM spray, as defined by BCPC.

Comet 200 is a fungicide with protectant and curative properties for disease control in winter and spring wheat, winter and spring barley, winter and spring oats, rye and triticale, as summarised below:

	Winter Wheat	Spring Wheat	Winter Barley Spring Barley	Winter and Spring Oats	Rye	Triticale
Yellow rust	С	С	C		С	С
Brown rust	С	С	C		С	С
Crown rust			-(/ xO	С		
Net blotch			CCC			
Rhynchosporium		()	MC MC		MC	

C = Control

MC = Moderate Control

Yield response may be obtained in the absence of visual disease.

#### Time of Application

Apply Comet 200 at the start of foliar disease attack. A maximum of two applications can be made up to and including flowering just complete in winter and spring wheat, rye and triticale and up to and including emergence of ear just complete in winter and spring barley and winter and spring oats.

### Rate of Application

Apply 1.25 litres Comet 200 in a minimum of 200 litres of water per hectare.

### **Resistance Management**

- A maximum of 2 applications of any strobilurin type (Ocl) product can be made to any cereal crop.
- In cereals this product must always be applied in a mixture vitin an approved fungicide that has a different mode of action. Both products must be applied at ecommercied rates.

### **Important Notes**

- 1. Avoid spray drift on to neighbouring crops
- 2. Wash equipment thoroughly after use.

# **Mixing and Application**

### Mixing

Half fill the spray tank with clean water and start the agitation. SHAKE THE CONTAINER WELL before use and pour in the required amount of product. Rinse any empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed.

When tank mixes are to be used, each product should be added separately to the spray tank.

# Compatibility

For details of compatibilities contact your distributor, local BASF representative or the BASF Technical Services Hotline: 0044 0845 602 2553. Alternatively, visit our viebsite www.agricentre.basf.ie.

#### Trade Mark Acknowledgments

Comet® is a registered trademark of BASF.

### **Additional Product Safety Information**

This section does not form part of the product label under S.I. No. 159 of 2012.

The information on this label is based on the best available information including data from test results.

## Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.basf.ie/Comet-200-IE/MSDS Alternatively, contact your supplier.