



Hygeia

HYMEC TRIPLE

FOR THE CONTROL OF BROAD-LEAVED WEEDS IN CEREALS AND GRASSLAND

A soluble concentrate formulation 310 g/L (26.2% w/w) Dichlorprop-P acid, 160 g/L (13.5% w/w) MCPA acid and 130 g/L (11.0% w/w) Mecoprop-P acid formulated as the dimethylamine salt for the control of broad-leaved weeds in cereals.

Always read labels carefully before use. Use pesticides safely.
For use only as an Agricultural Herbicide

PROTECT FROM FROST - FOR PROFESSIONAL USE ONLY

MAX DOSE
2.5L/HA
READ LABEL FOR DETAILS



DANGER

RISK AND SAFETY INFORMATION

Harmful if swallowed. Causes serious eye damage. May cause an allergic skin reaction. Toxic to aquatic life with long lasting effects. Do not eat, drink or smoke when using this product. Wear protective gloves/protective clothing/eye protection/face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON

CENTER or doctor/physician. If medical advice is needed, have product container or label at hand. Keep out of reach of children. Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste. **To avoid risks to human health and the environment, comply with the instructions for use**

PCS No. 03890

ADDITIONAL SAFETY PHRASES: Do not contaminate water with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads)

- Winter and Spring Wheat
- Barley, Oats, Rye

10 Litre e



5 390182 040563

MARKETED BY:
Hygeia Chemicals Ltd.
Oranmore, Co. Galway, Ireland.
Tel: 091-794722 Fax: 091-794738
Email: enquiries@hygeia.ie Web: www.hygeia.ie

Irish 24 hour emergency number:
National Poisons Centre in Ireland: 01-8092566
Approval holder Nufarm UK Limited, Wyke Lane,
Wyke, Bradford, West Yorkshire, BD12 9EJ, UK

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE Crop Winter and spring Wheat, Barley, Oats, Rye, Triticale, Spelt and Durum Wheat **Maximum Individual Dose** 2.5 L product/ha **Maximum Total Dose** 2.5 L product/ha **Latest Time of Application** Before second node Detectable (GS 32). **Method of Application:** Tractor mounted/trailed sprayer **Other specific restrictions:** The earliest time of application for all crops is the 1st March in the year of harvest. The total amount of Mecoprop-P applied to an individual crop, or in a single year in the case of a perennial crop, must not exceed the maximum total dose of Mecoprop-P approved for application to that crop by any single Mecoprop-P containing product.

DIRECTIONS FOR USE IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

Do not spray cereals that have been, or are to be, undersown with clover.
Do not roll or harrow crops within a week of spraying.
Avoid overlapping spray swathes.
Do not apply during cold weather or drought.
Do not apply during rain or if rain is expected.
Do not spray in windy weather and avoid drift onto broad-leaved plants outside the target area.

ADDITIONAL SAFETY PHRASES

Keep livestock out of treated areas for at least 2 weeks and until poisonous weeds such as ragwort have died and become unpalatable.

WEED CONTROL Weeds Controlled in Cereals

Susceptible at 1.5 L product/ha

Charlock	<i>Sinapis arvensis</i>	5cm across/high
Common Chickweed	<i>Stellaria media</i>	5cm across/high
Crane's-bill species	<i>Geranium spp.</i>	6 true leaves
Fat Hen	<i>Chenopodium album</i>	5cm across/high
Shepherd's Purse	<i>Capsella bursa-pastoris</i>	4 true leaves

Susceptible at 2.0 L product/ha

Common Poppy	<i>Papaver rhoeas</i>	5cm across/high
--------------	-----------------------	-----------------

Susceptible at 2.5 L product/ha

Buttercups	<i>Ranunculus species</i>	6 true leaves
Cleavers	<i>Galium aparine</i>	5cm across/high
Common Mouse Ear	<i>Cerastium holosteoides</i>	5cm across/high
Corn Spurry	<i>Spergula arvensis</i>	5cm across/high
Hempnettles	<i>Galeopsis spp.</i>	4 true leaves

Moderately Susceptible at 1.5 L product/ha

White clover	<i>Trifolium repens</i>	5cm across/high
--------------	-------------------------	-----------------

Moderately Susceptible at 2.0 L product/ha

Creeping Thistle #	<i>Cirsium arvense</i>	6 true leaves
Deadnettle species	<i>Lamium spp.</i>	Before flowers open
Fumitory	<i>Fumaria officinalis</i>	8 true leaves
Runch	<i>Raphanus raphanistrum</i>	Before side shoots form

Moderately Susceptible at 2.5 L product/ha

Field Pansy	<i>Viola arvensis</i>	4 true leaves
Mayweeds	<i>Matricaria species</i>	2 true leaves
Oil Seed Rape	<i>Brassica napus</i>	3 true leaves
Redshank	<i>Polygonum persicaria</i>	2 true leaves
Speedwells	<i>Veronica species</i>	6 true leaves

Creeping thistle will be effectively controlled, but some regrowth may occur from rhizomes.

MANUFACTURED BY: Hygeia Chemicals Ltd.
 Oranmore, Co. Galway, Ireland.
 Tel: 091-794722 Fax: 091-794738
 Email: enquiries@hygeia.ie
 Web: www.hygeia.ie



SPECIFIC CROP INFORMATION CEREALS HYMEC TRIPLE can be used on all commercial varieties of winter and spring sown wheat, winter and spring sown barley, winter and Spring Sown Oats, Rye, Triticale, Spelt, Durum wheat.

RATE OF USE Apply HYMEC TRIPLE at 1.5 - 2.5 L product/ha. Use the appropriate rate as indicated in the weed control table.

WATER VOLUME 200 to 400 L water/ha. Use the higher rate in dense stands of cereals, or if weed growth is dense.

Timing: **Winter Sown Wheat, Barley, Durum Wheat, Rye and Oats:** Apply HYMEC TRIPLE in the spring from leaf sheath erect stage until before the second node detectable stage (GS 32). **Spring Sown Wheat, Barley, Rye and Oats:** Apply HYMEC TRIPLE from first leaf fully expanded stage until before the second node detectable stage (GS 32).

Spray weeds when the crop is actively growing and is at the correct growth stage. The best results will be obtained if spraying is done while the majority are at the seedling stage. However, spraying at the seedling stage is not advised if temperatures are low and should be delayed until temperatures increase and growth becomes more active. Optimum control will be achieved by spraying when the temperature is above 10°C.

APPLICATION Spray Quality Apply as a medium quality spray (as defined by BCPC). A spray pressure of 2-3 bar is recommended. **MIXING** Apply using conventional ground vehicle mounted/drawn equipment. Ensure that all application equipment is clean. Make sure the sprayer is set to give an even application at the correct volume. Add half the required volume of water and start agitation, add the required quantity of Hymec Triple. Fill the tank to the required volume of water whilst maintaining agitation. Triple rinse containers with water and add washing to the spray tank. Continuous agitation must be maintained until spraying is complete. After use, the spraying machine must be thoroughly cleaned. Wash equipment thoroughly with water and wetting agent or liquid detergent immediately after use. Wash out again with water, drain and allow drying. Spray out, fill with clean water and leave overnight. Spray out again before storing or using for another product. Traces of Hymec Triple can cause harm to susceptible crops sprayed later.

DRIFT Do not spray in windy conditions as the spray drift may cause damage to neighbouring crops. The following crops are particularly susceptible sugar beet, oilseed rape, peas, turnips and most market garden crops including lettuce and tomatoes under glass.