CARPOVIRUSINE EV02

INSECTICIDE

PCS 04825

A suspension concentrate (SC) formulation containing *Cydia pomonella* Granulosis Virus M (CpGV-M) 1 x 10¹³ GV/l

Risk and Safety Information

Warning

May cause an allergic skin reaction

If medical advice is needed, have product container or label at hand. Keep out of reach of children. Avoid breathing spray

Wear protective gloves/protective clothing/eye protection/face protection.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except . triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

Micro-organisms may have the potential to provoke sensitizing reactions. 10 percent of the mixture consists of ingredients of unknown inhalation toxicity

To avoid risks to human health and the environment, comply with the instructions for use.

Approval holder and Marketing company:

Arvsta LifeScience S.A.S. Route d'Artix BP 80 64150 NOGUÈRES – France Phone number : + 33 (0) 559 60 92 92

24 hours emergency telephone no. +44 (01) 1235 239670

Manufacturing date and batch no.: see packaging

NET CONTENTS



CARPOVIRUSINE® EVO2

IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL INSECTICIDE

| Crops | Maximum single dose | Max. n°. of applications | Maximum total dose | Latest time of application |
|---|---------------------|--------------------------|--------------------|----------------------------|
| Apple, pear | 1 L/ha | 3 | 3L/Ha | 1 day before harvest |
| READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCI | | | | |

SAFETY PRECAUTIONS

Operator protection

WEAR SUITABL PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES Consult your specialist advisor for further information. AND SUITABLE RE. TRATORY EQUIPMENT* when handling the concentrate or applying the product. Disposed litering facepiece respirator to at least EN149 FFP3 or equivalent. MIXING AND SPRAYING

Envirgimente protection

equanter chear surface wale work in the second of the second se

Sturage and disposal

DO NOT RE-USE ONT INF fo any purpose. KEEP AWAY FROM DOD, RIM, AND ANIMAL FEEDINGSTUFFS. KEEP IN ORIGE AL CO. TAINER, tightly closed, in a safe place.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions Out of the freezer, CARPOVIRUSINE EVO 2 can be kept in the refrigerator or cold room wit in this ection must be read carefully in order to obtain safe and successful use of (+4°C) for maximum 8 months, or at room temperature for 1 month (not exceeding 25°C). his produce

RESTRICTIONS OR WARNING

Consult processor before using on crops for processing. CARPOVIRUSINE EVO 2 has been tested on a range of common apple varieties but russeting cannot be ruled out on varieties prone to this problem, e.g. Cox.

RESISTANCE

Total reliance on one pesticide will hasten the development of resistance.

Alternating insecticides with different modes of action and alternative control measures Conditions of Supply

in the planned programme is a recognised anti-resistance strategy. Applying CpGV Seller warrants that this product conforms to the chemical description on the label products with different isolates in alternation within a treatment programme will reduce thereof and is reasonably fit for purposes stated on such labels, only when used in the possibility of resistance development. CROP SPECIFIC INFORMATION Useful control of Codling Moth larvae and reduction in damage to apple and pear.

Application rate: 1 L/ha in 1000 L/ha water. Maximum number of treatments per year: 3 Pre-Harvest Interval: 1 day

Timing of Applications: GS 71-87

CARPOVIRUSINE EVO 2 may be applied from fruit setting until fruit maturation (GS 71-87), on young larvae for optimal efficacy. Timing of applications should be based on information from pheromone traps and/or specialist advice. Do not apply when rain is expected.

Apply the recommended dose of 1.0 L/ha to full height trees typical of orchards at full canopy density. Where tree height and/or canopy density is reduced, the dose (and water volume) should be adjusted in accordance with an appropriate dose adjustment scheme.

Apply with a foliar spray system, ensuring a uniform spraying. CARPOVIRUSINE EVO 2 only acts by ingestion and does not have any contact activity. Therefore, it is crucial to ot contininate water with the product or its container. Do not clean application have very good spray coverage of foliage and fruits. Timing of applications should be rain is expected.

Storage Conditions

CARPOVIRUSINE EVO 2 must be kept frozen (-18°C) for long period storage (at least 2 vears following manufacturing date). Thaw at least 1 day before the treatment at ambient temperature. CARPOVIRUSINE EVO 2 can be frozen several times.

COMPATIBILITY

Do not apply CARPOVIRUSINE EVO 2 in tank mixture with other pesticides or foliar fertilisers

This product is to be used only in accordance with the recommendations and instructions given on the labels provided with this pack. Use in any other circumstances is entirely at user's risk.

accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or manner of use or application, all of which are beyond the control of the seller. In no case shall seller be liable for consequential, special or indirect damages resulting from the use of handling of this product. All such risks shall be assumed by the buyer. Seller makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

()

