lima gold 5%

Granular bait containing 5 % w/w metaldehyde Molluscicide for control of slugs in agricultural and horticultural crops.

25 Kg

Safety Information

Suspected of damaging fertility. If medical advice is needed, have product container or label at hand

Keep out of the reach of children.

Do not get in eyes, on skin, or on clothing. Wear protective gloves.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can

be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use

PCS No. 0 '961

PRECAUTIONS

If you feel unwell seek medical advice immediately (s' ow the , bel w, ere possible).

In case of accident, call the National Poisons Information Centre Tel: +353 (0)1 809 2566 or +353 (0)1 837 9964

Do not contaminate water with the productor its container. Do not cl an application equipment near surface water/Ave contamination via dl ins from farmyards and roads.

To protect aquatic organisms do not one on permeable sur aces uch as asphalt, concrete, cobblestones, railway tracks and other cituations with a high risk of run-off.

To protect wild birds and wild mammals remove spillages.

Keep poultry out of treated areas for at least 7 day follo ving nor cation.

Protect from frost, store in a dry, cool, frost-free ventilate 1 - ea.

24 hour emergency number - National Poisons Centre: +353 (01) 809 2566 or +353 (01) 837 9964

UFI: 7J1E-V7TJ-1C0A-MXD5

Batch number: see container **Registration holder:** Sharda Europe b.v.b.a, Jozef Mertensstraat 142, 1702 Dilbeek, Belgium Phone no: +91 22 6678 2800 **Emergency number: +91 22 6678 2800 (8.00 – 16.00h)**

Registered logos of Sharda Cropchem Ltd.



READ LABEL CAREFULLY BEFORE USE

IMPORTANT INFORMATION

| FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL/HORTICULTURALMOLLUSCIDE | | | | |
|---|----------------------|--------------------------|----------------|---|
| Сгор | Max. individual dose | Max. no. of applications | Max.total dose | Latest timing of application |
| Wheat, barley, oats, rye & triticale | 4 kg/ha | | 12 kg/ha | GS 29 (end of tillering) |
| Oilseed rape | 4 kg/ha | | 12 kg/ha | GS 29 (9 or more side buds visible) |
| Maize | 4 kg/ha | | 12 kg/ha | GS 15 (5th leaf unfolded) |
| Lettuce | 4 kg/ha | | 12 kg/ha | GS 39 (rosettes complete) |
| Field Berries (Blackberry, Blackcurrant, Blueberry, Cranberry, Gooseberry, Loganberry, Raspberry, Redcurrant, Whitecurrant) | 4 kg/ha | | 12 kg/ha | GS 69 (End of flowering) |
| Pome fruit & Stone fruit (apple, pear, plum, cherry) | 4 kg/ha | | 12 kg/ha | GS 69 (End of flowering) |
| Grapes (Wine and the) | 4 kg/ha | | 12 kg/ha | GS 69 (End of flowering) |

A GOV is a molluscicide or the control of slugs in crops.

D. PECTIONS FOR USF

IMPORTANT: This inform turn is approved as part of the Product Label. All instructions with in this section must be read carefully in order to obtain safe and successfor use of this product.

RESTRICTIONS

Prof ssic ral use only

DO NOT at bly before heavy rainfall, which may reduce the persistence of ranking cate application of pellets to broad-leaved plants may result in lowing of the pellets in foliage. Care should be taken to avoid this during application in edible crops.

Keep dogs out of the field during application. When the applicators stops/starts, ensure no small heaps of pellets are left on the soil surface as they may be consumed by dogs and other animals. Spillages must be buried or removed.

APPLICATION RECOMMENDATIONS

Best results are achieved when application is made during mild, damp weather when slugs are the most active. When slug attack is heavy or prolonged, repeat treatments may be required. **All crops**

4 kg/ha broadcast evenly over soil or standing crop using any suitable slug pellet applicator. For crops to be drilled best results are from application a few days before drilling. Alternatively, apply just before or just after drilling. For emerged or established crops, apply immediately damage appears. Do not apply more than three times per season (maximum of 12 kg/ha) Minimum interval between applications 7 days

COMPANY ADVISORY INFORMATION

Slug pellets are pesticides, keep pesticides out of water

• Pellets should not fall within 5m of the top of the bank of a watercourse or ditch. To achieve this, users should determine the spread width of the applicator with the product to be applied prior to any application (to the rear as well as the side of the applicator).

- Calibrate your applicator before use
- Fill and clean application equipment in the field
- Do not fill or clean down in the yard
- Avoid applying slug pellets when heavy rain is forecast
- . Check that product packaging is intact and clear up spillages immediately

