
# EU 513 CCL FRANCE - CCL UK - CCL ITALY

Page 1



15mm

## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

TO OBTAIN THE PROPERTY OF THE

Use of an antifoam is compulsory with this product

Do not plant potatoes, sugar beet, vegetables, beans or other leguminous crops in the calendar year following an application of HALCYON. Following good agricultural practice ensure that plant remains have completely decayed before planting susceptible crops.

If application of HALCYON is followed by dry weather conditions, particularly on sandy soil, consult the manufacturer for specific following

### ! Neighbouring Crops/Plants

Avoid damage by drift onto susceptible crops, non-target plants or waterways. Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings, gardens, ponds, lakes or watercourses.

Animal Waste/Farmyard/Slurry/Compost/Mulch Management
HALCYON residues in plant tissues which have not completely decayed may affect succeeding susceptible crops eg peas, beans and other legumes, sugar beet, carrots and umbelliferae, potatoes and tomatoes, lettuce and other compositae.

Do not use any plant material treated with HALCYON for composting or mulching Do not use animal waste (eg manure, slurry) from animals fed on grass treated with HALCYON, or fodder resulting from grass treated with

HALCYON, on susceptible crops eg. peas, beans and other legumes, sugar beet, carrots and umbelliferae, potatoes and tomatoes, lettuce and other compositae, or land intended for growing such crops, until all plant tissues have completely decayed.

On not use animal waste (eg manure, slurry) from animals fed on grass treated with HALCYON for composting or mulching susceptible

## CONSULT THE MANUFACTURER FOR SPECIFIC ADVICE.NOTES

Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass. Therefore do not spray in drought, very hot or very cold weather conditions.

Do not roll or harrow grass for 10 days before or 7 days after application.

Late applications of HALCYON may lead to a slight leaning of the grass but this is transient and does not affect the yield of the grass.

HALCYON is safe to grass. Very occasionally some yellowing of the sward may occur; this is transient and quickly outgrown. To allow maximum translocation of HALCYON to the roots do not cut grass for 7 days after application.

If the grass has been cut for hay or silage or grazed leave 2-3 weeks to allow sufficient regrowth to occur before spraying.

Where there is a high reservoir of weed seed in the soil and/or a historically high weed population, a programmed weed control approach may be needed involving a second application the following year.

Control may be reduced if rain falls within 1 hour of application.

## Wash equipment thoroughly with water and detergent immediately after use. CLOVER MANAGEMENT

Clover will be killed by application of HALCYON.

Do not drill clover within 4 months of applying HALCYON.

Once control of target weeds has been achieved, and you have ensured that plant remains have completely decayed, clover can be re-introduced 4 months after application

## WEED CONTROL

	Weeds	Rates of use litres/ha	Optimum timing of application	
	Broad-leaved dock	2.0	Treat when the docks are actively growing, in the rosette stage up to 25 cm high or wide	
	Curled dock	2.0	Treat when the docks are actively growing, in the rosette stage up to 25 cm high or wide	
	Creeping thistle	2.0	Treat when the thistles are actively growing, in the rosette stage up to 25 cm high	
	Spear thistle	2.0	Treat when the thistles are actively growing, in the rosette stage up to 25 cm high	
	Common nettle	2.0	Treat when the nettles are actively growing, up to 30 cm high	
g	Creeping buttercup	2.0	Treat when the buttercups are actively growing, before flowering	
	Dandelion	2.0	Treat when the dandelions are actively growing, before flower bud	
	Common chickweed	1.0	Treat when the chickweed is actively growing, up to 30 cm wide	

The timing of application of HALCYON is crucial. For good results the product must be applied to actively gr growth stage as detailed in the WEED CONTROL section of this label.

## HALCYON should be applied through a tractor-mounted hydraulic sprayer provided it is in good y

Fill the spray tank half full with water and add required amount of HALCYON mixing well. To

## spray tank is full. Maintain agitation while spraying. Use the spray immediately.

For overall application HALCYON should be used in a spray volume of 200 litre ec re to g good coverage of the weeds.

A higher water volume of 250-300 litres of water per hectare may be require tion is high and wh dense at the time of application

Apply as a MEDIUM quality spray as defined by the BCPC systematics

## Corteva Conditions of Supply

All goods supplied by us are of high grade and we believe hem to be suita, but, as we cannot exercise control over the handling, mixing or use, or the weather conditions before, or ng or after application which may affect the performance of the good conditions and warranties, statutory or otherwise of the quantum or fitness from the purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any fair or in performance of injury whatsoever arising from a storage, handling, application or use. These conditions canno

To access the Safety Data Sher or this placed so the code or use the weblink below:

## SAFETY PRECAUTIONS Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or contaminated

WEAR SUITABLE PROTECTIVE COVERALLS, SUITABLE PROTECTIVE GLOVES AND RUBBER BOOTS when applying by hand-held equipment. WHEN USING DO NOT FAT DRINK OR SMOKE

WASH CONCENTRATE from skin or eyes immediately.

To protect groundwater do not apply to levs less than 1 year old Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target

DO NOT CONTAMINATE WATER with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads.) KEEP LIVESTOCK out of treated areas for at least 7 days or until foliage of any poisonous

h as ragwort has died and become unpalatable. prote aquatic organisms, respect an unsprayed buffer zone of 5m to surface water

ect or undwater do not apply to grassland within one year of sowing from seed

EEP OUT OF REACH OF CHILDREN.

AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

manually rinsing three times. Add washings to sprayer at time of filling and dispose of fluroxypyr (present as 144 g/litre fluroxypyr-methyl heptyl ester).

## Distributed by: Whelehan Crop Protection

T P Whelehan Son & Co. Ltd Suite 11. Bunkilla Plaza.

Bracetown Business Park,

Co Meath, Ireland +353 1 4688900

## **CORTEVA**agriscience



# **Halcyon**<sup>®</sup>

™ Trademarks of Corteva Agriscience and its affiliated companies.

## GROUP O HERBICIDES Product Registration number : PCS No: 05393

For use as an agricultural herbicide for the long term control of ANNUAL and PERENNIAL BROAD-LEAVED WEEDS in GRASSLAND.

A water in oil emulsion containing 30 g /litre (3.5% w/w) aminopyralid RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or (present as 36 g/litre aminopyralid potassium) + 100 g /litre (14.14% w/w)

## READ DIRECTIONS FOR USE ON ATTACHED LEAFLET. PROFESSIONAL USE ONLY

Triple rinse containers, Puncture and Invert to Dry at Time of Use PROTECT FROM FROST.

## Authorization Holder:

Corteva Agriscience UK Limited CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE, United Kingdom Telephone: (01462) 457272 Fax: (01462) 426605

24 Hour Emergency Telephone Number: 076 680 5288 e-mail:ukhotline@corteva.com

Technical Hotline: 0800 689 8899 National Poison Centre (Dublin): 01 8092166

## **NET CONTENTS: 5 LITRES (2)**

## IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Maximum No. of applications: 1 per year

Latest time of application 7 days before grazing

Other specific restrictions: Do not use between 31st July and 1st March

or applications made between 1st March and 31st May, 1 application may be made per calendar year or applications made between 1st June and 31st July, only 1 application may be made in a 2 year period his product must not be used on grassland that will be grazed by animals other than cattle or sheep

Ihis product must not be used on grassland that will be grazed by animals other than cattle or sheep The product must not be used on land where vegetation will be cut for animal feed, fodder or bedding r for composting or mulching within one calendar year of treatment Manure from animals grazed on 'Halcyon' treated grassland should not leave the farm. Manure from animals fed silage, hay or haylage produced from grassland treated in the previous calendar year shot not leave the farm. This manure should only be spread onto agricultural grassland. This product must be used with an antifoaming agent

The product must not be used on airfields

Spot applications can be applied between Mar 1st and Aug 31st Do not apply spot applications to more tha 20% of the area in August: if infestation levels exceed this figure then a broadcast application should be made Users must have received adequate instruction, training and guidance in the safe use of the product an must take all reasonable precautions to protect the health of human beings, creatures and plants and

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for Using Plant Protection Products.

May be fatal if swallowed and e Cause skin irritation Causes serious eye damage

Very toxic to aquatic life with long lasting effects

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No 05393