

WHEN USING DO NOT EAT, DRINK OR SMOKE. WASH CONCENTRATE from skin or eyes immediately.

Environmental protection:

hand-held equipment.

Extreme care must be taken to avoid spray drift on non-crop plants outside of the target area. DO NOT CONTAMINATE WATER with the product or container. (Do not clean application equipment not a surface water / Avoid contamination via drains from fainyards and roads.)

AND SUITABLE PROTECTIVE GLOVES when applying by

Storage and disposal:

KEEP OUT OF REACH OF CHILDREN. KEEP AWAY FROM FOOD, DRINK AN AMMAL TEEDING

KEEP IN ORIGINAL CONTAINF 1, tight closes, 1 a safe place RINSE CONTAINER THOROG HLY by using an integrate I pressure rinsing devices manually rinsing three times. Activashings to sprayer at time of filling and dispose of more

Distributed by:



TP Whelehan Son & Co Ltd Suite 12 Bunkilla Plaza, Bracetown Business Park, Clonee, Co Meath Tel: +353 14688900





roduct Registration N mber: 'S No. 05396

A soluble concentrate co. taining 12 g ae/litre aminopyralid and 120 g ae/litre trialogy For the control of cortain PERENNIAL, HERBACEOUS and WOODY WEEL in all enity glissland.

READ VIL SCTICILY FOR USE ON ATTACHED LEAFLET. PROTECT F. 971 FROST.

) IR PROFESSIONAL USE ONLY

T ple Rinse Containers, Puncture And Invert To Dry At Time Of Use th isation Holder:

C reva Agriscience UK Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE Telephone: + 44 1462 457272

Technical enquiries: +44 800 6898899, e-mail: ukhotline@corteva.com 24 Hour Emergency Telephone Number: 076 680 5288

National Poison Centre (Dublin): 01 8092166

NET CONTENTS: 1 LITRE Θ



IMPORTANT INFORMATION. FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Amenity grassland, non-crop areas 4.0 litres product per hectare or Maximum Individual Dose: 200 mL per 10 litres water

Maximum number of applications: 1 per year Maximum Total Dose: 4.0 litres product per hectare per year Latest time of application 31st August

Do not use between 31st August and 1st March

For applications made between 1st March and 31st May, 1 application may be made per calendar

For applications made between 1st June and 31st July, only 1 application may be made in a 2

or applications made between 1st August and 31st August, only 1 application may be made in a 3 year period

DO NOT USE on land that will be grazed by livestock
DO NOT USE on land where vegetation will be cut for animal feed, fodder or bedding nor for sting or mulching within one calendar year of treatment

DO NOT USE on airfields

DO NOT apply spot applications to more than 20% of the area Users must have received adequate instruction, training and guidance in the safe use of the product and must take all reasonable precautions to protect the health of human beings,

creatures and plants and safeguard the environment.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for Using Plant Protection Products.



Causes serious eye irritation. May cause damage to kidneys through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects

Do not breathe mist/vapours/spray. Wear protective gloves / protective clothing / eye protection / face protection

IF IN FYPS: Rinse cautiously with water for several

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do - continue rinsing

Contains Triclopyr. May produce an allergic reaction. Dispose of contents / container to a licensed

hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste
To avoid risks to human health and the

environment, comply with the instructions for use.







DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

WARNINGS

Take extreme care to avoid drift onto susceptible crops, non-target plants or waterways. Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings, gardens, ponds, lakes or watercourses.

SENSITIVE PLANTS may be harmed by residues of GARLON ULTRA in soil and treated vegetation. Do not apply GARLON ULTRA on or adjacent to soil which may be used as garden top-soil, potting soil, etc. or to grass which may be cut and used as mulch or for compost for horticultural or garden crops. Do not use cuttings from treated grass for mulching or composting.

Cut treated grass must not be removed from site, or used for animal feed, animal WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION bedding, composting or mulching

Treated grass must not be used for grazing.

On level ground there is negligible lateral movement but do not apply GARLON UL around desirable trees or shrubs, do not spray under the canopy and 1 metr least from the trunk of desirable trees or shrubs. Care should be taken on slopes prevent leaching into areas where desirable shrubs, etc are present.

Grass and weeds must be actively growing to ensure good weed control minimal check to the grass. Therefore do not spray in drought, hot or very cold weather conditions.

Do not use on grass less than one year old.

To allow maximum translocation to the roots do not cut grass for 7 days after application. Clover will be killed by application of GARLON ULTŘA.

Wash equipment thoroughly with water and detergent immediately after use.

HERBACEOUS WEED CONTROL BROADCAST APPLICATION

WEEDS CONTROLLED. RATES OF USE AND TIMING OF APPLICATION

	Weeds	Rates of use litres/ha	Optimum timing of application
	Bramble ¹	4.0	Treat when the bramble is actively growing but is less than 50 cm hig
	Common mugwort	4.0	Treat when the mugwort is actively growing and less than 70cm high.
	Common nettle ¹	2.0	Treat when the nettles are actively growing and less than 70cm high.

Weeds	Rates of u litres/ha	
Creeping thi	stle 4.0	Treat when the thistle is actively growing and less than 70cm high.
Hogweed ²	4.0	Treat when the hogweed is actively growing and less than 70cm high.
Rosebay willowherb	4.0	Treat when the willowherb is actively growing and less than 15cm high.

A second application may be required the following year.

WOODY AND HERBACEOUS WEED CONTROL LOCALISED SPOT APPLICATION

TRA	Weeds	Rate of use mL/10 litres of water	Optimum timing of application
e at	Bramble	150	Up to 1 metre high.
s to	Broom	200	Up to 1 metre high.
	Buddleia	200	Up to 1 metre high.
	Gorse	150	Up to 1 metre high.
and	Japanese knotweed	200	1 metre high, with good foliage cover.

TIMING OF APPLICATION

GARLON ULTRA should be applied between March and the end of Augustiming of application of GARLON ULTRA is crucial and for good results. ARLON ULTRA must be applied to actively growing weeds.

APPLICATION

Broadcast Treatment

provided it is in good working order and has been calibrated according to the weblink weblink manufacturers' recommendations.

Fill the spray tank half full with water and add the required amount of GARLON ULTRA mixing well. Top up with water and continue agitation until the spray tank is full. Maintain agitation while spraying. Use the spray immediately.

For overall application GARLON ULTRA should be used in a spray volume between 300L and 600L per hectare to give good coverage of the weeds. Higher water

volumes are recommended where weed densities are high and where the grass is dense at the time of application.

Spray Quality Apply as a MEDIUM quality spray as defined by the BCPC system

Spot Treatment

For localised treatment using a suitable lance from a hapsack thractor mounted sprayer, use a solution of 150-200 mL of GARLON LTRA depends, on the target weed, per 10 litres of water. Woody weeds struld be coroughly wetted with the spray solution, but spraying until "run-off" w. d. ease act into the the spray solution, but spraying until "run-off" w. d. ease act into the flood jets is recommended to prevent dri are six id be ken to avoid local overdosing. Woody weeds should not excee a metre boyht.

COMPATIBILITY

ct Corteva Agrisciance or our For the latest advice on tank mixes distributor.

TRADEMARK ACY JOWLEDGEM, 'TS

Garlon Ultra is a tra emark of Co eva Agriscience. All other brand i mes used this document are trad manufactur s in which propriets y rights may exist.

Cortex AgriSci ace Conditions of Supply

All good, surplied by are of high grade an tive brieve than to be suitable: we anot expuse control over their stora a handley, mixing or use, or the we ther colditions before, during or after application which may affect the performance of the goods, all cormula, and warranties, statutory or otherwise, s of the quality or fitness fo any p ose of our goods are excluded. No consibility will be accepted by us or sellers for any failure in performance,

dain be or injury whatso we rising from the storage, handling, application or se. These conditions of not by varied to storage, staff or agents whether or not they inpervise or assist in the use of such go. as.

IRELAND GARLON ULTRA should be applied through a tractor-mounted hydraulic sprayer To access the afety Dai. She of or this product scan the QR code or use the

.tps://www.corteva.ie/content/dam/dpagco/corteva/eu/ie/en/ files/sds/GARLON-ULTRA-IRL-SDS.pdf

Alternatively contact your supplier

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. ■ WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin or eyes immediately.

Environmental protection:

Extreme care must be taken to avoid spray drift onto Trademarks of Corteva Agriscience and its affiliated companies. non-crop plants outside of the target area.

DO NOT CONTAMINATE WATER with the product or its GROUP 0 container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads.)

Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Distributed by:



Suite 12 Bunkilla Plaza, Bracetown Business Park, Clonee, Co Meath Tel: +353 14688900

CORTEVA[™]



HERBICIDE



Product Registration Number: PCS No. 05396

A soluble concentrate containing 12 g ae/litre aminopyralid and 120 g ae/litre triclopyr

For the control of certain PERENNIAL, HERBACEOUS and WOODY WEEDS in amenity grassland.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET. PROTECT FROM FROST.

FOR PROFESSIONAL USE ONLY

Triple Rinse Containers, Puncture And Invert To Dry At Time Of Use

Authorisation Holder:

Corteva Agriscience UK Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE Telephone: + 44 1462 457272

Technical enquiries: +44 800 6898899, e-mail: ukhotline@corteva.com

NET CONTENTS: 1 LITRE \bigcirc

24 Hour Emergency Telephone Number: 076 680 5288 National Poison Centre (Dublin): 01 8092166

IMPORTANT INFORMATION. FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Crons / Situations: Amenity grassland, non-crop areas

4.0 litres product per hectare or Maximum Individual Dose:

200 mL per 10 litres water

Maximum number of applications: 1 per year
Maximum Total Dose:
4.0 litres product per hectare per year

Latest time of application 31st August

Do not use between 31st August and 1st March For applications made between 1st March and 31st May, 1 application may be made per calendar

For applications made between 1st June and 31st July, only 1 application may be made in a 2

For applications made between 1st August and 31st August, only 1 application may be made in

a 3 year period.

DO NOT USE on land that will be grazed by livestock

DO NOT USE on land where vegetation will be cut for animal feed, fodder or bedding nor for composting or mulching within one calendar year of treatment DO NOT USE on airfields

DO NOT apply spot applications to more than 20% of the area

Users must have received adequate instruction, training and guidance in the safe use of the product and must take all reasonable precautions to protect the health of human beings, creatures and plants and safeguard the environment.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for Using Plant Protection Products.



Causes serious eye irritation. May cause damage to kidneys through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects Wear protective gloves / protective clothing / eye protection / face protection

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do - continue rinsing

If eye irritation persists: get medical advice / attention. IF ON SKIN: Wash with soap and water

Contains Triclopyr. May produce an allergic reaction. Dispose of contents / container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 05396