

4 mm

**Last updated by :** PUBLI H

▼

®

HERBICIDE

GROUP	4	HERBICIDES
-------	---	------------

**Whelehan Crop Protection**  
Suite 12 Bunkilla Plaza, Bracetown Business Park  
Clonee, Ireland D15 WR13  
Tel: + 353 (1) 574 2410

**Triple Rinse Containers, Puncture and Invert to Dry at time of Use**

**Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the code of practice for using plant protection products.**

2311



# EU513 Leaflet Label

## Inside pages

245 mm  
233 mm

4 mm

4 mm

Last updated by : PUBLI H

### DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

#### NOTES

Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass.  
DO NOT spray in drought, very hot or very cold weather.  
Control may be reduced if rain falls within 2 hours of application.  
DO NOT roll or harrow grass for 10 days before or 7 days after application.  
Clover will be killed or severely checked by application of DOXSTAR® PRO.  
DO NOT sow kale, swedes, turnips or grass mixtures containing clover by direct drilling or minimum cultivation techniques within 6 weeks of applying DOXSTAR PRO.  
DOXSTAR PRO is safe to established grassland. Very occasionally some yellowing of the sward may occur; this is transient and quickly outgrown.  
Occasionally, e.g. in hot conditions, vapour drift may occur, making it particularly important to assess the risk to neighbouring vegetation.  
Take care to avoid drift onto susceptible crops, non-target plants or waterways. Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings, gardens, ponds, lakes or watercourses.

Wash out spray equipment thoroughly with water and detergent immediately after use. Traces of DOXSTAR PRO could cause harm to susceptible crops sprayed later.

#### Grazing Interval

Exclude livestock during treatment and do not allow livestock to graze treated grassland for at least 7 days following treatment, and until foliage of any poisonous weeds which may have been affected by application has died and become unpalatable.

#### GENERAL INFORMATION

DOXSTAR PRO is a foliar acting herbicide for the control of broad-leaved dock and curled dock in established grassland.



### ESTABLISHED GRASSLAND

#### WEEDS CONTROLLED

DOXSTAR PRO is recommended for the control of broad-leaved dock and curled dock, in the rosette stage up to 200 mm high or across.

Weed	Rate of Use litres product/ha	Optimum timing of application
Broadleaved dock, Curled dock	2.0	At the rosette stage up to 200 mm tall or across

#### RATE OF USE AND TIMING

Apply DOXSTAR PRO at 2.0 litres/ha in 300 to 400 litres of water as an overall treatment in the autumn or spring for the control of docks.

Use the higher volume of water where the weeds are large at the time of application and where the grass is dense.

Application timing of DOXSTAR PRO is crucial for good control and must be made when the weeds and grass are actively growing to ensure good weed control and minimal check to the grass. Docks should be treated in the rosette stage and within the size given above.

It is important that there is sufficient leaf area for uptake of the herbicide. This is particularly true on well-established docks which will have an extensive tap root, but which early in the spring or soon after cutting may only have a few small leaves.

If the grass (and hence the weeds) has been cut for conservation hay or silage, or has been grazed, leave for 2 to 3 weeks to allow sufficient regrowth to occur before spraying.

To allow maximum translocation of DOXSTAR PRO to the roots do not cut grass for 28 days after application.

On large well-established docks and where there is a large reservoir of seed in the soil a further control programme in the following year may be required.

#### APPLICATION

DOXSTAR PRO may be applied only through tractor-mounted hydraulic sprayers providing it is in good working order and have been calibrated according to the manufacturers' recommendations.

Do not apply in knapsack sprayers.

#### MIXING

Fill the spray tank half full with water and add the required amount of DOXSTAR PRO. Continue filling with water and maintain agitation during filling, while spraying and during any stoppages. Mix up only sufficient spray volume for the area to be treated and use the spray immediately.

#### SPRAY QUALITY

Apply DOXSTAR PRO as a MEDIUM spray as defined by the BCPC system.

# EU513 Leaflet Label

## Inside pages

245 mm

233 mm

4 mm

**Trademark Acknowledgement**  
DOXSTAR PRO is a trademark of Corteva Agriscience and its affiliated companies.  
All other brand names used in this document are trademarks of other manufacturers in which proprietary rights may exist.

**Corteva Agriscience Conditions of Supply**  
All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<https://www.corteva.ie/content/dam/dpagco/corteva/eu/ie/en/files/sds/DOXSTAR-PRO-IRL-SDS.pdf>

Alternatively contact your supplier

4 mm

Last updated by : PUBLI H