

Safety Information

Suspected of damaging fertility.

If medical advice is needed, have product container or label at hand.

Keep out of the reach of children.

Do not get in eyes, on skin, or on clothing. Wear protective gloves.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.



To avoid risks to human health and the environment, comply with instructions for use

PC' NU. 0555

PRECAUTIONS

If you feel unwell seek medical advice immediately show the label where possible).

In case of accident, call the National Poisons Information of the Tel: +353 (0)1 809 2566 or +353 (0)1 837 9964

Do not contaminate water with the project on s container. Do not coan application equipment near surface water/^-oid ontamination has ains from farmyards and roads.

To protect aquatic organisms do not apply or impermeable orfaces such as asphalt, concrete, cobblestones, railway tracks and other situations with a high risk of run-off.

To protect wild birds and wild mammals remove stillage

Keep poultry out of treated areas for at least 7 days follo ving application.

Protect from frost, store in a dry, cool, frost-free ventilated area.

 $24\ hour\ emergency\ number -$ National Poisons Centre: +353 (01) 809 2566 or +353 (01) 837 9964

UFI: FF1E-D744-QC0U-YPHP

Net content: 25 Kg

Batch number: see container Registration no.: PCS No. 05533

Registration holder: Sharda Europe b.v.b.a, Jozef Mertensstraat 142,

1702 Dilbeek, Belgium

Distributed by:

Phone no: +91 22 6678 2800

Emergency number: +91 22 6678 2800 (8.00 - 16.00h)

Registered logos of Sharda Cropchem Ltd.



SHARDA CROPCHEM LIMITED Domnic Holm 29'" Road, Bandra (West) Mumbai - 400050, India, Tel. +39 02 66101029

READ LABEL CAREFULLY BEFORE USE

IMPORTANT INFORMATION

FOR PROFESSIONA USE ONLY AS AN AGRICULTURAL / HORTICULTURAL MOLLUSCICIDE				
Crop	Max single dose	Max. no. of applications	Max.total dose	Latest timing of application
Grapes (Wine and Table)	7 kg/ha		21 kg/ha	GS 69 (End of flowering)
Field Berries (Blackberry, Blackcurrant, Blueberry, Cranberry, Gooseberry, Loganberry, Raspberry, Redcurrant, Whitecurrant)	7 kg/ha		21 kg/ha	GS 69 (End of flowering)
Stone & pome fruit (apple, pear, plum, cherry)	7 kg/ha		21 kg/ha	GS 69 (End of flowering)
Lettuce	7 kg/ha		21 kg/ha	GS 39 (rosettes complete)
Oilseed rape	7 kg/ha		21 kg/ha	GS 29 (9 or more side buds visible)
Wheat barley, oat, rye & triucale	7 kg/ha		21 kg/ha	GS 29 (end of tillering)
Maize	7 kg/ha		21 kg/ha	GS15 (5th Leaf unfolded)
Spe_ifir_pests controlled: igs _ snailr				

DESTROYER is a my lust led the for the control of slugs in crops.

DIRECTIONS OR CSE.

IMPORTANT: his information is approved as part of the Product Label. All instructions with models section must be read carefully in order to obtain safe and successful use of this product.

יES, יירי ONS

Prefessional use only

DO NOT apply before heavy rainfall, which may reduce the persistence of granules. Late application of pellets to broad-leaved plants may result in lodging of the pellets in foliage. Care should be taken to avoid this during application in edible crops.

Keep dogs out of the field during application. When the applicators stops/starts, ensure no small heaps of pellets are left on the soil surface as they may be consumed by dogs and other animals. Spillages must be buried or removed.

APPLICATION RECOMMENDATIONS

Best results are achieved when application is made during mild, damp weather when slugs are the most active. When slug attack is heavy or prolonged, repeat treatments may be required.

All crops

7 kg/ha broadcast evenly over soil or standing crop using any suitable slug pellet applicator. For crops to be drilled best results are from application a few days before drilling. Alternatively, apply just before or just after drilling. For emerged or established crops, apply immediately damage appears. Do not apply more than three times per season (maximum of 21 kg/ha) Minimum interval between applications 7days

COMPANY ADVISORY INFORMATION

Slug pellets are pesticides, keep pesticides out of water

- Pellets should not fall within 5m of the top of the bank of a watercourse or ditch. To achieve this, users should determine the spread width of the applicator with the product to be applied prior to any application (to the rear as well as the side of the applicator).
- Calibrate your applicator before use
- Fill and clean application equipment in the field
- Do not fill or clean down in the yard
- Avoid applying slug pellets when heavy rain is forecast
- Check that product packaging is intact and clear up spillages immediately