THEFT FENCE[®] H SHAKE WELL BEFORE USE PROTECT FROM FROST

A suspension concentrate containing 480 g/litre flufenacet for the pre- and early post-emergence control of specified grass and broad-leaved weeds in winter wheat and winter barley.

SAFETY INFORMATION HARMFUL IF SWALLOWED MAY CAUSE DAMAGE TO ORGANS THROUGH PROLONGED OR REPEATED EXPOSURE VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction

Contains flufenacet. May produce an allergic reaction Do not eat, drink or smoke when using this product

Wear protective gloves/protective clothing

IF SWALLOWED: immediately call a POISON CENTER or doctor/ physician

Collect spillage

Dispose of contents/container to a licensed hazardous-waste disposal contractor

or collection site except for triple rinsed empty containers which can be disposed of

as non-hazardous waste

Do not contaminate water with the product or its container (Do not clean application

equipment near surface water/Avoid contamination via drains from farmyards and roads)

Horizontal boom sprayer: To protect aquatic organisms respect at unsprayed buffer zone of 5m to surface water bodies. Direct spray away from water

Wash all protective clothing after use

To avoid risks to human health and the environment, comply with the instructions for use Professional use only

PCS No. 05873

KI-EL_Fence_LBL_05-10-17

Albaugh

WARNING

Operator protection Since Bit Note Description <thdescription< th=""> Description</thdescription<>			
operator's normal working position is within a closed cab on a tractor or on a self-propeiled sprayer. Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) suITABLE PROTECTIVE GLOVES AND FACE PROTECTIVE GLOVES AND FACE PROTECTIVE WEAR SUITABLE PROTECTIVE GLOVES WHAT SUITABLE PROTECTIVE GLOVES WEAR SUITABLE PROTECTIVE CLOTHING (FACESHIELD) when handling the MEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) during application. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) during application. Strate data Storage and disposal Storage and and disposal Storage and d	Operator protection	SMOKE.	its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads
sprayer. Engineering control of operator exposure must be used where reasonably practicable must be used where reasonable protections must be used where reasonable protections must be used where reasonable protections must be used where reasonable preasonable protections must			Storage and disposal
Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the manually rinsing three times. Add washings to concentrate. WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) during application. KEEP OUT OF REACH OF CHILDREN. WASH HANDS before meals and after work. Environmental protection to protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water RINSE CONTAINER THOROUGHLY by wasy form water. RINSE CONTAINER THOROUGHLY by RINSE CONTAINER THOROUGHLY by RINSE CONTAINER THOROUGHLY by manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. NOTO NON-TARGET PLANTS OUTSIDE OF THE TARGET AREA Do not contaminate water with the product or			
mušt be ušed where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) during application. WEAR SUITABLE PROTECTIVE GLOVES Wear SUITABLE PROTECTIVE CLOTHING WEAR SUITABLE PROTECTIVE GLOVES Wear SUITABLE PROTECTIVE CLOTHING WEAR SUITABLE PROTECTIVE GLOVES Wear SUITABLE PROTECTIVE CLOTHING WEAR SUITABLE PROTECTIVE CLOTHING WEAR SUITABLE PROTECTIVE GLOVES When handling contaminated surfaces. Wear SUITABLE PROTECTIVE CLOTHING WEAR SUITABLE PROTECTIVE SUITABLE Manufacturing date and batch number: see pack Do not contaminate water with the product or COVERALLS) during application. STORE IN ORGINAL CONTAINER tightly STORE STORE STANCES STORE STORE STORE STORE STORE STORE STORE STORE STORE STORE STORES STORE STORE STORES STORE STORES STORES STORE STORES STOR			
equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTIVE (FACESHIELD) when handling the concentrate. WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces. WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces. WEAR SUITABLE PROTECTIVE CLOTHING WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces. WEAR SUITABLE PROTECTIVE CLOTHING WEAR SUITABLE PROTECTIVE SUITABLE PROTECTIVE SUITABLE NTO NON-TARGET PLANTS OUTSIDE OF THE TARGET AREA Do not contaminate water with the product or (COVERALLS) during application.	must be used where reasonably practicable		STORE IN ORIGINAL CONTAINER tightly
WEAR SUITABLE PROTECTIVE CLOTHING Do not contaminate water with the product or (COVERALLS) during application.	in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. WEAR SUITABLE PROTECTIVE GLOVES	Environmental protection To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies. Aim spray away from water. RISK TO NON-TARGET INSECTS OR OTHER ARTHROPODS. See Directions for Use. TAKE EXTREME CARE TO AVOID DRIFT ONTO NON-TARGET PLANTS OUTSIDE OF	closed, in a safe place. DO NOT RE-USE CONTAINER for any purpose. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. Keep dry and protect from frost in a suitable pesticide store.
	WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) during application.	Do not contaminate water with the product or	

For advice on medical emergencies, fires, spillages or chemical hazards, telephone: +44 (0)1235 239 670 (24 hour) Approval Holder and Marketing Company: Albaugh TKI d.o.o., Grajski trg 21, 2327 Rače, Slovenia, Tel: +386 (0)2 60 90 211; Fax: +386 (0)2 60 90 410 Fence® is a trademark of Albaugh. Albaugh is a trading style of Albaugh Europe Sàrl and Albaugh TKI d.o.o.

Fence®

PCS No. 05873

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE.

Crops/Situations	Maximum Individual Dose	MaximumNumber of Treatments	Maximum Total Dose	Latest Time of Application
Winter wheat, winter barley	0.5 litres product /ha	-	0.5 litres product /ha	3 leaves unfolded stage (GS13) Before 31st December of the year of planting

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

GENERAL INFORMATION

Fence is a contact and residual herbicide for use alone or in specified tank-mixtures and sequences for the control of certain annual grass and broad-leaved weeds in winter wheat and winter barley. Fence may be applied pre-emergence of the crop in the autumn up to the 31st December and before GS13 in winter wheat and winter barley. Best results will be obtained from pre-emergence or early post-emergence applications to weeds that have not yet emerged on have emerged and are rapidly growing but still at early growth stages.

WARNINGS AND RESTRICTIONS

EXTREME CARE MUST BE TAKEN TO AVOID SPRAY DRIFT to avoid damage to plants outside the target area.

DO NOT APPLY under windy conditions.

DO NOT ALLOW SPRAY SWATHS TO OVERLAP as crop damage may occur.

Do not apply to broadcast crops.

Do not treat undersown cereals or those due to be undersown

Do not cultivate or disturb the soil after application

Do not use on sands, very light soils or stony or gravelly soils, or on soils with more than 10% organic matter content.

Do not use on any soil with poor or compacted soil structure or that is subject to waterlogging.

A reduced level of weed control may result if a prolonged dry period follows application.

Weed control is likely to be unsatisfactory if application is made to dry cloddy seedbeds.

Under certain climatic conditions, some transient discolouration of the crop foliage may occur, which can be particularly severe if heavy rain falls soon after application. These transient symptoms are quickly outgrown and have no adverse effect on crop yield. Do not apoly to croos that are under stress from any cause including pest or disease attack. frost damage or nutrient deficiency.

Avoid making applications during prolonged or severe frosty conditions as there is a risk of transitory crop damage from which the crop will normally recover.

Trace amounts of Fence remaining in the sprayer after use can cause damage to other crops subsequently treated with the same equipment. Immediately after use thoroughly clean the sprayer and all equipment as detailed in the section "SPRAYER DECONTAMINATION".

Contract agents should be consulted before using on crops grown for seed.

Effects on processed crops have not been established. Consult processor before using Fence on crops intended for processing.

RESISTANCE MANAGEMENT

There is a moderate risk for the development of weed resistance to Fence. Strains of some annual weeds (e.g. Black-grass, Wild oats and Italian Ryegrass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures and alternating the use of Fence, with other herbicides with a different mode of action and a similar spectrum of activity. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor or crop adviser. Implement a weed resistance strategy based on Good Agricultural Practice and including the following:

DO NOT use Fence as a stand-alone treatment against black-grass, Use only in tank-mix or sequence with other herbicides approved for use against black-grass and with alternative modes of action.

DO NOT use Fence as the sole means of weed control in successive crops

Follow the label recommendations.

Adopt complementary weed control practices.

Minimise the risk of spreading weed infestations.

Implement good spraying practice to maintain effective weed control.

Use the correct spray nozzles to maximise coverage.

Apply only under appropriate weather conditions.

Monitor herbicide performance and report any unexpected results to Albaugh (www.albaugh.eu).

APPLICATION

Before use, make sure the spraying equipment is cleaned from any previous use and that it's calibrated to apply the recommended spray volume and pressure. After use, make sure the sprayer and all equipment are thoroughly deaned as detailed in the section "SPRAYER DECONTAMINATION".

Apply in 200-400 litres of water per hectare as a MEDIUM spray (BCPC). Do not allow the spray boom to bounce or yaw, and ensure an even coverage of the weed foliage and soil surface.

MIXING

Half fill the sprayer tank with clean water and begin gentle agitation. Add the required quantity of product into the sprayer tank and allow to disperse fully. Rinse the empty containers thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add the washings to the sprayer and continue agitation whilst bepring up the tank with water to the required level. Continue agitation until the mix is sprayed out. Spray, integrated after mixing, do not allow the mixture to stand in the sprayer. When using tank-mixes follow any specified order of addition on the partner product label(s).

SOIL CONDITIONS

Apply to moist level seedbeds that are of a fine fith and free from clods. Loose or puffy seedbeds should be consolidated by rolling or other cultivations prior to application. Good weed control depends on the removal of trash from a previous crop and adequate soil moisture.

The seed should be covered by at least 2-4 cm of soil to ensure adequate crop safety. Where a crop has been direct-drilled, the drill sits must be closed by harrowing across the direction of drilling prior to application. Where direct drilled crops are treated then ploughing and cultivating to a depth of 15 cm must take place before the planting of any subsequent crop. Do not apply to broadcast crops

WEEDS CONTROLLED

The following weeds are controlled by a pre-or early post-emergence application of Fence at 0.5 litres/ha.

Annual meadow-grass	Highly susceptible
Black-grass	Susceptible
Cleavers	Moderately resistant
Loose silky bent-grass	Susceptible

Best results will be obtained if rain falls within 7 days after application. Residual control may be reduced if there is a prolonged period of dry conditions.

TANK MIXTURES

Fence may be used in tank-mixture with the following herbicides for improved control of Blackgrass and other weeds.

Approved formulations of:

Pendimethalin

Diflufencan

Follow the label recommendations of the partner product when using any tank-mix.

FOLLOWING CROPS

After normal harvest of any crop treated with Fence, any crop may be planted.

Potato, peas, wheat, barley		Ploughing is not required	
	All other crops not listed above	Plough to a depth of at least 15cm	

REPLACEMENT CROPS

In the event that a crop treated with Fence should fail for any reason, the soil must be ploughed to a depth of at least 15cm and only the crops listed below may be planted as a replacement for the failed crop:

Potato, peas, wheat, barley

SPRAYER DECONTAMINATION

Trace amounts of Fence remaining in the sprayer after use can cause damage to other crops subsequently treated with the same equipment. Immediately after use, thoroughly clean the sprayer and all equipment with a proprietary detergent cleaner. It is essential that all nozzles, filters, tubing, strainers, pumps and the spray tank are thoroughly cleaned after use to avoid the risk of damage to crops subsequently treated with the same equipment. Spray equipment should not be drained or flushed onto land planted with or intended for planting with trees or crops other than oilsear rage, beans and pees.

TERMS AND CONDITIONS OF SUPPLY, SALE AND USE

Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors, and the occurrence and development of strains resistant to the active ingredient. Under certain circumstances, changes in activity or crop damage can occur. The manufacturer or supplier is unable to accept any fability in these circumstances. All goods supplied by us are of a high grade and we believe them to be suitable for the purpose for which we expressly supply them: but as we cannot exercise any control over their mixing, use or application which may affect the performance of the goods all conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff, our agents or the re-sellers of the product whether or not they supervise or assist in the use of such goods.

Fence® is a registered trademark of Albaugh.