DW X-OJYMA

AMYLO-X° WG

DIRECTIONS FOR USE

GENERAL INFORMATION

AMYLO-X® is a broad-spectrum preventive bio fungicide for control or suppression of fungal plant diseases. The active ingredient of AMYLO-X, Bacillus amyloliquefaciens subsp. plantarum D747, is a naturally occurring strain of the beneficial soil bacterium B. amyloliquefaciens, which colonises leaves, roots and other plant surfaces, killing pathogenic organisms by means of antibiosis and release of lytic enzymes. Moreover, B. amyloliquefaciens quickly colonises plant surfaces preventing establishment of diseases such as fungi.

AMYLO-X is applied as a foliar spray using air assisted or hydraulic pressure sprayer.

Apply up to 2.5 kg/ha, diluted in 200 to 1500 litres of water/ha, depending on the crop and its development stage.

CROPS, APPLICATION RATES AND **APPLICATION TIMING**

Leafy Vegetables

Lettuce, Endive, Cress, Land cress, Lamb's Lettuce, Rocket, Red Mustard, Spinach, Spinach Beet, Watercress and Chicory

1.5 - 2.5 kg/ha in a water volume of 200 - 1000 L/ha Number of applications: max 6

Diseases controlled: Sclerotinia spp. and Botrytis cinerea Timing: apply at the beginning or prior to infection or from 4 true leaf stage to lettuce hearts have reached typical size (BBCH 79).

Strawberries, Cane fruit and Bush/small fruit Strawberry, Blackberry, Loganberry and Rubus hybrid,

Raspberry, Blueberry, Gooseberry, Blackcurrant and Redcurrant 1.5 – 2.5 kg/ha in a water volume of 500 - 1000 L/ha

Number of applications: max 6 Diseases controlled: Botrytis cinerea and powdery mildew

(Podosphaera aphanus)

Timing: apply at the beginning or prior to infection or from the beginning of leaf development up to (BBCH 89) harvest.

Fruiting Vegetables

Tomato, Pepper, Chilli, Aubergines, Courgette and Summer Squash, Cucumber, Melon, Winter Squash, Pumpkin and

1.5 – 2.5 kg/ha in a water volume of 500 – 1000 L/ha Number of applications: max 6

Diseases controlled: Botrytis cinerea and powdery mildew (Levelillula taurica) for Tomato, Pepper, Chilli and Aubergine and Botrytis cinerea only for all other crops listed Timing: apply at the beginning or prior to infection or from the beginning of leaf development up to (BBCH 89) harvest.

Including Mushrooms 15 g/100 kg compost

Diseases controlled: Trichoderma aggressivum Timing: once at set up of cultivation, spray onto growth substrate.

PREPARATION OF THE SPRAY SOLUTION

AMYLO-X may be applied with conventional sprayer, boom sprayers, power sprayers or knapsack sprayers. To ensure homogenous dispersion of the product in the tank-mix, half-fill the tank with water, add the required amount of the product and fill the tank under continuous agitation. Do not use highly alkaline or highly acidic water to mix sprays. Use a buffering agent if necessary to maintain neutrality (pH 6 to 8) of water in the tank. Maintain agitation during application. Apply immediately after mixing; do not allow spray mix to stand overnight.

COMPATIBILITY

AMYLO-X can be applied alone or in combination and/or rotation with chemical fungicides. Do not mix with products which could alter the spores' viability. It is always advisable to verify the compatibility or the potential phytotoxicity caused by a mixture with an untested product, since the possible incompatibility may depend on factors outside the control of the manufacturer, such as crop stage, climatic conditions, the used water volumes, etc.

STORAGE AND DISPOSAL

Store in a cool and dry place; high humidity and temperatures over 30°C decrease the activity of the product. The product is stable at room temperature (21-24°C) for 2 years. The disposal of product has to be performed in accordance with all applicable national and local environmental regulations. Completely empty the container into application equipment. Clean container promptly after emptying by triple rinsing. Dispose of empty container according to local regulations, alternatively contact a licensed disposal contractor. Avoid contamination of the environment from empty containers. not re-use or refill empty container

CLEANING OF THE SPRAY EQUIPMENT

Rinse the application equipment thoroughly with Distribute washings over the treated area or least unsprayed area in the crop for applying washings. After u clean the tank with clean water. Special cleaning agents

The product is intended for professional horticultural units ONLY. The responsibility of using the product remains with end user. All directions of use should be adhered to in order to ensure safe use of the product, including the safety of treated crops, persons and protection of the environme Follow all directions for use in order to avoid risks for human health and the environment.

The development of resistance to the active Bacillus amylolique faciens subsp. plantarum strain D747 can be considered a low risk, but cannot be excluded. To minimise the risk of resistance developing in the future it is good practice to use products with a range of active ingredients as part of advisable to use AMYLO X continually in sequential crops grown in the same physical structure.



MAPP 17976 PCS 06311

ranule (WG) containing 250 g/kg (5 x 1013 CFU/kg) ns subsp. plantarum strain D747 for the control of mildew and other fungal plant diseases on a range of crops grown panent protection with full enclosure.

(COSHH) CONTROL OF SUBSTANCES AZARDOVS TO HEALTH REGULATIONS
MAY APPLY TO THE USE OF THIS PRODUCT
AT WORLD - UK only

ETY PRECAUTIONS

Certis Beichim

ntal protection INATE SURFACE WATER OR DITCHES with

Storage and disposal SE CONTAINER THOROUGHLY by using an integrated essure device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. DO NOT RE-USE EMPTY CONTAINER for any purpose. KEEP AWAY FROM FOOD, DRINK AND ANIMAL

KEEP OUT OF THE REACH OF CHILDREN

PORTANT INFORMATION

USE ONLY AS A PROFESSIONAL

ximum individual dose: 2.5 kg product /ha Maximum number of applications: 6

Minimum interval between treatments: 7 days Latest time of application: BBCH 89 for strawberrie

For fungi (mushrooms): 1 application at set up of cultivation with a maximum application rate of 15 g per 100 kg compost.

Other specific restrictions:

- 1) Treatment must only be made under 'permanent protection' situations which provide full enclosure (including continuous top and side barriers down to below ground level) and which are present and maintained over a number of years.
- Reasonable precautions must be taken to prevent access of birds, wild mammals and honey bees to) To minimise airborne environmental exposure,
- READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE

FOR USING PLANT PROTECTION PRODUCTS.

Water dispersible granules (WG) containing 250 g/kg (5 x 10¹³ CFU/kg) Bacillus amyloliquefaciens subsp. plantarum

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, triple rinsed clean containers which can be disposed of as non-hazardous waste.

Bacillus amyloliquefaciens subsp. plantarum strain D747 may have the potential to provoke sensitising

To avoid risks to human health and the environment, comply with the instructions for use. MAPP 17978 / PCS 06311



DIRECTIONS FOR USE:

READ CAREFULLY BEFORE OPENING THE PACK

Material Safety Data Sheet

To access the Safety Data Sheet for this product, scan the QR code or use the web link.

https://cutt.lv/rviN7v

Batch No.: Date of Manufacture: - See container

For advice on medical emergencies, fires, spillages phone +44 (0)870 190 6777

For technical and non-emergency calls phone +44 (0)845 373 0305

CONDITIONS OF SUPPLY: The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumer'





Marketing Company: Certis Belchim BV Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridgeshire CB21 6AD Tel: +44 (0)1223 652500 | email: info.uk@certisbelchim.com | www.certisbelchim.co.uk Authorisation Holder: Mitsui AgriScience International S.A./B.V. Certis Belchim

Authorisation Profeet. William Agrissicate International States

Belchim

Belchi

Net content 1kg

