

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Basagran SG (PCS No 05192)

CROPS:

[Outdoor grown: Bulb onion, garlic, shallot & salad onion]

For this EAMU the following crop definitions apply:

Bulb onion includes onions grown from seed, sets or transplants.

This approval provides for the use of the product **Basagran SG (PCS No 05192)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 01.07.21. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Basagran SG
<i>Active Substance:</i>	Bentazone
<i>Formulation:</i>	870g/kg SG
<i>Authorisation Holder:</i>	BASF plc
<i>Marketing Company:</i>	BASF Ireland Ltd
<i>PCS No:</i>	05192

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

User type/Function: Professional Use Only / herbicide.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Outdoor grown: onion, garlic & shallot	0.55 kg/ha <u>Or</u> 1.1 kg/ha	2 per crop <u>Or</u> 1 per crop	1.1 kg/ha		21 days before harvest	General weed control
Outdoor salad onion	0.55 kg/ha	2	1.1 kg/ha	7 days	21 days before harvest	General weed control

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed/air assisted sprayer only. Application by hand held sprayer is NOT allowed.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 1m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Basagran SG as a horticultural herbicide for the control of; Charlock, common chickweed, cleavers, cranes bill, fools parsley, field forget me not, goosefoot, corn marigold, mayweeds, black & white mustard, small

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

nettle, black nightshade, volunteer oilseed rape, field penny cress, pale persicaria, scarlet pimpernel, wild radish, redshank, shepherds purse & corn spurrey on the crops and situations listed above. This product is also useful in controlling a number of other difficult weeds if applied when the weed is not more than the 2 true leaf stage. Please refer to the product label for more information.

For bulb onion apply in 220 to 450 litres of water per ha.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other herbicides from different cross-resistant groups [different mode of action, HRAC group] with satisfactory efficacy against the target.

Basagran SG contains bentazone.

Bentazone is classed HRAC group: C3.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.