

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Trianum P (PCS No. 05486)

CROPS: strawberry (protected), bilberry (protected), blackcurrant (protected), and redcurrant (protected), blueberry (protected), cranberry (protected), elderberry, (protected), gooseberry (protected), mulberry (protected), and rosehips (protected), cane fruit (protected), and table and wine grapes (protected).

For this EAMU the following crop definitions apply:

Cane fruit include: blackberry, raspberry, loganberry and rubus hybrid.

This approval provides for the use of the product **Trianum P (PCS No. 05486)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 04.06.20. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Trianum P
<i>Active Substance:</i>	<i>Trichoderma harzianum</i> strain T22
<i>Formulation:</i>	10 g / kg /WG
<i>Authorisation Holder:</i>	Koppert BV.
<i>Marketing Company:</i>	Koppert (UK) Limited
<i>PCS No:</i>	05486

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

User type/Function:

Professional Use Only / horticultural fungicide

Application:

Crop	Max. individual dose	Max. Number of applications	Max. Total dose	Minimum interval between applications	Latest time of application	Target
Redcurrant (protected), blackcurrant (protected), bilberry (protected), cranberry (protected), blueberry (protected), elderberry (protected), mulberry (protected), rosehips (protected) & gooseberry (protected).	15 kg/ha	2 per crop	30kg/ha	14 days	- At propagation	fusarium, pythium & <i>Rhizoctonia</i>
	And					
	15g per 1000 plants at cultivation	12 per crop	-	4 weeks	-	
Blackberry (protected), loganberry (protected), rubus hybrid (protected) & raspberry (protected).	15 kg/ha	2 per crop	30kg/ha	14 days	At propagation	fusarium, pythium & <i>Rhizoctonia</i>
	And					
	15g per 1000 plants at cultivation	12 per crop	-	4 weeks	-	
Strawberry (protected)	15 kg/ha	2 per crop	30kg/ha	14 days	At propagation	fusarium , pythium & <i>Rhizoctonia</i>
	And					
	15g per 1000 plants at cultivation	12 per crop	-	4 weeks	-	
Table and wine grapes (protected)	15 kg/ha	2 per crop	30kg/ha	14 days	At propagation	fusarium, pythium & <i>Rhizoctonia</i>
	And					
	15g per 1000 plants at cultivation	12 per crop	-	4 weeks	-	

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application may only be made using a drench, drip irrigation or glass house sprayer only. Note application by handheld sprayer is NOT allowed.
- (4) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.
- (5) This approval extends to professional use only.
- (6) This use is restricted to all off labels authorisations being grown under permanent protection with full enclosure at a water volume of 2.5 - 5 litres/ m² via drench application to growing medium and through drip irrigation systems.
- (7) A minimum interval of 14 days must be observed between applications at propagation
- (8) A minimum interval of 4 weeks must be observed between applications at cultivation
- (9) Reasonable precautions must be taken to prevent access of birds, wild mammals and honey bees to treated crop
- (10) To minimise airborne environmental exposure, vents, doors and other openings must be closed during and after application until the applied product has fully settle.
- (11) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (12) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Trianum P as a horticultural fungicide for the control of: fusarium, pythium & *Rhizoctonia* on the crops and situation above.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, HRAC group] with satisfactory efficacy against the target.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.