

Chapter 7

List of off label approvals which have been granted up to 1/1/18 for use on food crops

Current list of off-label approvals for use on Food Crops as of *22 February 2018*

Warning: Off-Label Approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

Instruction sheets for off-label approvals can be obtained from your Teagasc office, or Bord Bia, or local advisor.

Disclaimer

The Department of Agriculture Food and the Marine maintains its websites to enhance public access to information about the Department and its activities and to provide services to its customers. Every effort has been made to ensure that the information provided on the Department's websites is accurate and up-to-date. If you notice any errors or omissions please let the Department know as soon as possible. The Department gives no guarantees, undertakings or warranties concerning the accuracy, completeness or up-to-date nature of information provided on its websites and does not accept any liability whatsoever arising from any errors or omissions.

The Department reserves the right to change all or any of the information provided at any time, but shall not be responsible for or liable in respect of any such change in information. The Department has no liability for any loss occasioned from the use of information appearing on these websites. Any links from these websites to external websites are provided as a matter of convenience only, and should not be taken as an endorsement of the contents or practices of those external websites.

Unless otherwise stated reproduction is authorised except for commercial purposes, provided the source is acknowledged.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|---------------------|---------------|--|------------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Apples</i> | | | | | | | | | | | |
| | Cerone | 03933 | Bayer CropScience Limited | Growth Regulator | | | 0.75 l/ha | 0.75 l/ha | - | 10 days before harvest | 10 days |
| | Curenox 50 WP | 05829 | Industrias Quimicas Del Valles S.A (IQV, S.A.) | Fungicide | <i>Ethephon</i> | <i>480 g/l</i> | 2.0 kg/ha | 8.0 kg/ha | - | Before GS 07 | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | <i>Copper oxychloride</i> | <i>87.8% w/w</i> | 0.25 l/ha | 1.0 l/ha | | 28 days before harvest | 28 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Difenoconazole</i> | <i>250 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.8 kg/ha | 2.4 kg/ha | 3 | 3 days before harvest | |
| | Talius | 04604 | DuPont (UK) Ltd | Fungicide | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | 0.25 l/ha | 0.5 l/ha | 2 per year | 49 days before harvest | 49 days |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Proquinazid</i> | <i>200 g/l</i> | | | | | |
| <i>Apples - post harvest</i> | | | | | | | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 0.6 l/ha | 1.8 l/ha | 3 | After final harvest but before final leaf fall | |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------|---------------------|---------------|----------------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|--|---|-------------|
| <i>Asparagus</i> | | | | | | | | | | | |
| | Amistar Opti | 03408 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | Before senescence | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | Before senescence | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Curator | 04504 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | Before senescence | |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | Before senescence | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | Before senescence | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | | | 2.0 kg/ha | 4.0 kg/ha | 2 - 1 to growing crop. 1 - post harvest underneath | 28 days before harvest | |
| | | | | | <i>Pyridate</i> | <i>45 % w/w</i> | | | | | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | After harvest and before end of September | |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Vertik | 04030 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | Before senescence | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-------------------------------|---------------------|---------------|---------------------------------|-----------------|-----------------------------|-------------------|---------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Aubergines - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 100 mls/100 litres water | - | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.45 l/ha | 0.9 l/ha | 2 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | 0.01kgs/5 litres water | - | - | - | 0 days |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.06 kgs/100 litres water | 0.18 kgs/ha | 3 / crop | 3 days before harvest | 3 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 3.0 l/ha | 12.0 l/ha | - | 3 days before harvest | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | Revus | 02789 | Syngenta UK Ltd | Fungicide | | | 0.6 l/ha | 2.4 l/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Mandipropamid</i> | <i>250 g/l</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 0.5 l/ha | 1.0 l/ha | 2 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | 3 | 1 day before harvest | 1 day |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|--------------------------------------|-----------------|---------------------------|-----------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Baby leaf production</i> | | | | | | | | | | | |
| | Acrobat MZ | 04059 | BASF plc | Fungicide | | | 2.0 kg/ha | 6.0 kg/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Dimethomorph</i> | <i>75 g/kg</i> | | | | | |
| | | | | | <i>Mancozeb</i> | <i>667 g/kg</i> | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Bonalan | 05763 | Gowan Crop Protection Ltd | Herbicide | | | 8.0 l/ha | 8.0 l/ha | - | Incorporation pre sowing/planting | |
| | | | | | <i>Benfluralin</i> | <i>180 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | CIPC | 03066 | Arysta LifeScience Benelux SPRL | Herbicide | | | 1.9 l/ha | 3.8 l/ha | 2 per crop | 2 weeks after planting out | |
| | | | | | <i>Chlorpropham</i> | <i>400 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 | 3 days pre harvest | 3 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | Before GS 49 | 14 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days pre harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Kerb Flo | 02830 | Dow AgroSciences Ltd | Herbicide | | | 1.875 l/ha | 1.875 l/ha | 1 | 24 days before harvest | 24 days |
| | | | | | <i>Propyzamide</i> | <i>400 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|---------------------------|-----------------|---|-----------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Baby leaf production</i> | | | | | | | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Paraat | 03614 | BASF plc | Fungicide | | | 0.36 kg/ha | 0.72 kg/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Dimethomorph</i> | <i>500 g/kg</i> | | | | | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | | | 4.5 l/ha | 4.5 l/ha | 1 | 6 weeks pre harvest | |
| | | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| | Pendifin 400 SC | 05731 | Finchimica S.p.A | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 per crop | 6 weeks before harvest | 6 weeks |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kgs/ha | 1.2 kgs / ha | 3 | 7 days | |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 3 mls per sq m | 6 mls per sq m | - | Immediately post drilling | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|---------------------------------|-----------------|--|----------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Baby leaf production</i> | | | | | | | | | | | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | 1.9 kg/ha | 5.7 kg/ha* | | 35 days before harvest | 35 days |
| | | | | | <i>Metalaxyl M</i> | 4.0 %w/w | | | | | |
| | | | | | <i>Mancozeb</i> | 64 %w/w | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | 1015.1 g/l | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | - | 14 days before harvest | |
| | | | | | <i>Pyraclostrobin</i> | 6.7 % w/w | | | | | |
| | | | | | <i>Boscalid</i> | 26.7 % w/w | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | 100 g/l | | | | | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 / crop | 6 weeks pre harvest | 6 weeks |
| | | | | | <i>Pendimethalin</i> | 400 g/l | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 3.3 l/ha | 3.3 l/ha | 1 / crop | 6 weeks pre harvest | |
| | | | | | <i>Pendimethalin</i> | 455 g/l | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | | | | | <i>Cycloxydim</i> | 100 g/l | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 0.7 kg/ha | 1.4 kg/ha | 2 | 7 days before harvest | |
| | | | | | <i>Cyprodinil</i> | 37.5 % w/w | | | | | |
| | | | | | <i>Fludioxonil</i> | 25.0 % w/w | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | 12% w/w | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | 12% w/w | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | 2 | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | 50 %w/w | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 ml/ha | 600 ml/ha | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Spinosad</i> | 480 g/l | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | 250 g/l | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------|---------------------|---------------|---------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|----------------------------------|-------------|
| <i>Basil</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| <i>Beans - broad</i> | | | | | | | | | | | |
| | Claw 100 | 05874 | DHM Agrochemicals Ltd., | Herbicide | | | 1.5 l/ha | 1.5 l/ha | 1 per crop | Before flower buds visible stage | 7 weeks |
| | | | | | <i>Propaquizafop</i> | <i>100 g/l</i> | | | | | |
| | Farmco Zealot | 05299 | Farmco Agritrading Ltd | Herbicide | | | 1.5 l/ha | 1.5 l/ha | 1 per crop | Before flower buds visible stage | 7 weeks |
| | | | | | <i>Propaquizafop</i> | <i>100 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------|---------------------|--------------------------------------|---------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Beans - field</i> | | | | | | | | | | | |
| Anthem | 04761 | Adama Agricultural Solutions UK Ltd | Herbicide | | | | 3.3 l/ha | 3.3 l/ha | | Pre crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| Defy | 04277 | Syngenta UK Ltd | Herbicide | | | | 5.0 l/ha | 5.0 l/ha | 1 per crop | Pre-emergence | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| Dithane 945 | 03705 | Indofil Industries (Netherlands) B.V | Fungicide | | | | 2.0 kg/ha | 2.0 kg/ha | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>80 % w/w</i> | | | | | |
| Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | | 2.1 kg/ha | 2.1 kg/ha | 1 per crop | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| PDM 330 EC | 03582 | BASF plc | Herbicide | | | | 4.0 l/ha | 4.0 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| Pendifin 400 SC | 05731 | Finchimica S.p.A | Herbicide | | | | 3.3 l/ha | 3.3 l/ha | 1 per crop | Pre-crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | | 2.0 kg/ha | 2.0 kg/ha | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>64 %w/w</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | | | | | |
| Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | | 5.0 l/ha | 5.0 l/ha | 1 per crop | Pre-emergence | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| Stomp 400 SC | 02879 | BASF plc | Herbicide | | | | 3.3 l/ha | 3.3 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| Stomp Aqua | 03750 | BASF plc | Herbicide | | | | 2.9 l/ha | 2.9 l/ha | 1 / crop | Pre crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | | | | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------|---------------------|---------------|---------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|----------------------------------|-------------|
| <i>Beans - field</i> | | | | | | | | | | | |
| | Wakil XL | 05471 | Syngenta UK Ltd | Seed Dressing | | | 2.0 kg/tonne | 2.0 kg/tonne | 1 per batch | Before drilling | |
| | | | | | <i>Cymoxanil</i> | <i>100 g/kg</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>50 g/kg</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>175 g/kg</i> | | | | | |
| <i>Beans - french</i> | | | | | | | | | | | |
| | Claw 100 | 05874 | DHM Agrochemicals Ltd., | Herbicide | | | 1.5 l/ha | 1.5 l/ha | 1 per crop | Before flower buds visible stage | 7 weeks |
| | | | | | <i>Propaquizafop</i> | <i>100 g/l</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | Immediately after sowing | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Farmco Zealot | 05299 | Farmco Agritrading Ltd | Herbicide | | | 1.5 l/ha | 1.5 l/ha | 1 per crop | Before flower buds visible stage | 7 weeks |
| | | | | | <i>Propaquizafop</i> | <i>100 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Beans - navy</i> | | | | | | | | | | | |
| | Claw 100 | 05874 | DHM Agrochemicals Ltd., | Herbicide | | | 1.5 l/ha | 1.5 l/ha | 1 per crop | Before flower buds visible stage | 7 weeks |
| | | | | | <i>Propaquizafop</i> | <i>100 g/l</i> | | | | | |
| | Farmco Zealot | 05299 | Farmco Agritrading Ltd | Herbicide | | | 1.5 l/ha | 1.5 l/ha | 1 per crop | Before flower buds visible stage | 7 weeks |
| | | | | | <i>Propaquizafop</i> | <i>100 g/l</i> | | | | | |
| <i>Beans - runner</i> | | | | | | | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | Immediately after sowing | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------|---------------------|---------------|---------------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Beet - fodder</i> | | | | | | | | | | | |
| | Deca 2.5% EC | 04952 | Sharda Europe b.v.b.a | Insecticide | | | 300 mls/ha | 300 mls/ha | - | 30 days before harvest | 30 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 300 mls/ha | | 30 days before harvest | 30 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Fortress | 02364 | Dow AgroSciences Ltd | Fungicide | | | 0.3 l/ha | 0.4 l/ha | 2 | 28 days before harvest | 28 days |
| | | | | | <i>Quinoxifen</i> | <i>500 g/l</i> | | | | | |
| | GAT Decline 2.5 E | 04422 | FMC Chemical s.p.r.l. | Insecticide | | | 300 mls/ha | 300 mls/ha | - | 30 days before harvest | 30 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | NEMGuard DE | 05368 | ECOspray Ltd | Nematicide | | | 10 kg/ha | 10 kg/ha | 1 | At drilling | |
| | | | | | <i>Garlic Extract</i> | <i>450 g/kg</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 0.5 l/ha | | 28 days before harvest | 28 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Beet - red</i> | | | | | | | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | NEMGuard DE | 05368 | ECOspray Ltd | Nematicide | | | 10 kg/ha | 10 kg/ha | 1 | At drilling | |
| | | | | | <i>Garlic Extract</i> | <i>450 g/kg</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Beet - sugar</i> | | | | | | | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 0.5 l/ha | | 28 days before harvest | 28 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|--------------------------------------|-----------------|-----------------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Beetroot</i> | | | | | | | | | | | |
| | Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 30 days before harvest | 30 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Escolta | 05681 | Bayer CropScience Limited | Fungicide | | | 0.35 l/ha | 0.7 l/ha | 2 per year | 21 days before harvest | 21 days |
| | | | | | <i>Trifloxystrobin</i> | <i>375 g/l</i> | | | | | |
| | | | | | <i>Cyproconazole</i> | <i>160 g/l</i> | | | | | |
| | Fortress | 02364 | Dow AgroSciences Ltd | Fungicide | | | 0.3 l/ha | 0.4 l/ha | 2 | 28 days pre harvest | 28 days |
| | | | | | <i>Quinoxifen</i> | <i>500 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 8 weeks before harvest | 56 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|---------------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Beetroot</i> | | | | | | | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | 10 kg/ha | 20 kgs/ha | 2 | Last application before September | - |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 0.5 l/ha | | 28 days before harvest | 28 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | Venzar 500 SC | 04805 | DuPont (UK) Ltd | Herbicide | | | 0.4 l/ha | 0.4 l/ha | - | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>500 g/l</i> | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | | | | | <i>Oxamyl</i> | <i>100 g/kg</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|-------------------------------------|-----------------|---|--|--------------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Blackberries - field</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 7 days before harvest | 7 days |
| | Apollo 50SC | 04610 | Adama Agricultural Solutions UK Ltd | Acaricide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 0.4 l/ha | 0.4 l/ha | 1 per year | 9 days before harvest | 9 days |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | <i>Clofentazine</i> | <i>500 g/l</i> | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | <i>Tebuconazole</i> | <i>250 g/l</i> | 3.0 l/ha | 3.0 l/ha | 1 | Before flowering or after harvest | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Masai | 03585 | BASF plc | Insecticide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 375 gms/ha | 375 g/ha | 1 / year | 21 days before harvest | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | 0.4 kgs/ha | 1.2 kgs/ha | 3 | 28 days before harvest | 28 days |
| | Scala | 03916 | BASF plc | Fungicide | <i>Pymetrozine</i> | <i>50 %w/w</i> | 2.0 l/ha | 4.0 l/ha | 2 per crop | 7 days before harvest | 7 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Pyrimethanil</i> | <i>400 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.8 l/ha | 1.6 l/ha | - | 21 days before harvest | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | <i>Carfentrazone-ethyl</i> | <i>60 g/l</i> | 1.0 kg/ha | 3.0 kg/ha | 3 | 10 days before harvest | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | <i>Fludioxonil</i> <i>Cyprodinil</i> | <i>25.0 % w/w</i> <i>37.5 % w/w</i> | 200 ml/ha in 1000L water | 400 ml/ha | 2 per year | 1 day before harvest | 1 day |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------------|---------------------|---------------|-------------------------------------|-----------------|---|-------------------|--------------------------|-------------------------|----------------------------|--|-------------|
| <i>Blackberries - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 10 days before harvest | 10 days |
| | Apollo 50SC | 04610 | Adama Agricultural Solutions UK Ltd | Acaricide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 0.4 l/ha | 0.4 l/ha | 1 per year | 9 days before harvest | 9 days |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | <i>Clofentezine</i> | <i>500 g/l</i> | 0.25 l/ha | 0.75 l/ha | 3 | 3 days before harvest | 3 days |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | <i>Thiacloprid</i> | <i>480 g/l</i> | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Tebuconazole</i> | <i>250 g/l</i> | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Masai | 03585 | BASF plc | Insecticide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 375 gms/ha | 375 g/ha | 1 / year | 21 days before harvest | |
| | Scala | 03916 | BASF plc | Fungicide | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | 2.0 l/ha | 4.0 l/ha | 2 per crop | 7 days before harvest | 7 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Pyrimethanil</i> | <i>400 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.8 l/ha | 1.6 l/ha | - | 21 days before harvest | 21 days |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | <i>Carfentrazone-ethyl</i> | <i>60 g/l</i> | 1.0 kg/ha | 3.0 kg/ha | 3 | 14 days before harvest | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 ml/ha in 1000L water | 600 ml/ha | 3 per year | 1 day before harvest | 1 day |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|---------------------|-----------------------------|---------------------|-----------------|---|-----------------|------------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Blackcurrants - field</i> | | | | | | | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | | 0.25 l/ha | 0.75 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Chikara Weed Cont | 04203 | ISK Biosciences Europe N.V. | Herbicide | | | | 120 g/ha | 120 g/ha | 1 per year | Before bud burst | |
| | | | | | <i>Flazasulfuron</i> | <i>25 % w/w</i> | | | | | |
| Corbel | 90438 | BASF plc | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | 3 | 2 weeks before harvest | 2 weeks |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| Manfil 75 WG | 03598 | Indofil Industries Limited | Fungicide | | | | 1.5 kg per 1000 litres water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75 % w/w</i> | | | | | |
| Manfil 80 WP | 03563 | Indofil Industries Limited | Fungicide | | | | 1.4 kg per 1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 1.5 kgs/ha | 1.5 kgs/ha | - | 7 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Penncozeb 80 WP | 03609 | UPL Europe Ltd | Fungicide | | | | 1.4 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Penncozeb WDG | 03608 | UPL Europe Ltd | Fungicide | | | | 1.5 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>750 g/kg</i> | | | | | |
| Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | | 0.4 kgs/ha | 1.2 kgs/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| Scala | 03916 | BASF plc | Fungicide | | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------------|---------------------|---------------|------------------------------|-----------------|--|-------------------|-------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Blackcurrants - field</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | 200 ml/ha in 250L water | 400 ml/ha | 2 per year | 7 days before harvest | 7 days |
| <i>Blackcurrants - protected</i> | | | | | | | | | | | |
| | AQ 10 | 05801 | CBC (Europe) S.r.l. | Fungicide | | | 70 g/ha | - | 12 | - | |
| | Scala | 03916 | BASF plc | Fungicide | <i>Ampelomyces Quisqualis</i> | <i>58 %w/w</i> | | | | | |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|-----------------------------|---------------------|-----------------|---|-------------------|------------------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Blueberries - field</i> | | | | | | | | | | | |
| Batavia | 05905 | Bayer CropScience Limited | Insecticide | | | | 0.75 l/ha | 1.5 l/ha | 2 | Post harvest application | |
| | | | | | <i>Spirotetramat</i> | <i>100 g/l</i> | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | | 0.25 l/ha | 0.75 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Chikara Weed Cont | 04203 | ISK Biosciences Europe N.V. | Herbicide | | | | 120 g/ha | 120 g/ha | 1 per year | Before bud burst | |
| | | | | | <i>Flazasulfuron</i> | <i>25 % w/w</i> | | | | | |
| Corbel | 90438 | BASF plc | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | 3 | 2 weeks before harvest | 2 weeks |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | | 3.0 l/ha | 3.0 l/ha | 1 | Before flowering or after harvest | |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| Manfil 75 WG | 03598 | Indofil Industries Limited | Fungicide | | | | 1.5 kg per 1000 litres water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75 % w/w</i> | | | | | |
| Manfil 80 WP | 03563 | Indofil Industries Limited | Fungicide | | | | 1.4 kg per 1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 1.5 kgs/ha | 1.5 kgs/ha | - | 7 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Penncozeb 80 WP | 03609 | UPL Europe Ltd | Fungicide | | | | 1.4 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Penncozeb WDG | 03608 | UPL Europe Ltd | Fungicide | | | | 1.5 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>750 g/kg</i> | | | | | |
| Scala | 03916 | BASF plc | Fungicide | | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------------|---------------------|---------------|------------------------------|-----------------|------------------------------------|-------------------|-----------------------|---|----------------------------|---|-------------|
| <i>Blueberries - field</i> | | | | | | | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest) | - | 3 days before harvest, N/A for post harvest | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| <i>Blueberries - protected</i> | | | | | | | | | | | |
| | AQ 10 | 05801 | CBC (Europe) S.r.l. | Fungicide | | | 70 g/ha | - | 12 | - | |
| | | | | | <i>Ampelomyces</i> | <i>58 %w/w</i> | | | | | |
| | | | | | <i>Quisqualis</i> | | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis</i> | <i>375 g/kg</i> | | | | | |
| | | | | | <i>subspecies kurstaki EG 2348</i> | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|-----------------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Broccoli</i> | | | | | | | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.2 l/ha | 0.2 l/ha | 1 / crop | 7 days before harvest | 7 days |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | <i>Thiacloprid</i> | <i>480 g/l</i> | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | <i>S-Metolachlor</i> | <i>960 g/l</i> | 1.0 l/ha | 2.25 l/ha | - | 21 days before harvest | 21 days |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | <i>Tebuconazole</i> | <i>250 g/l</i> | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | <i>Clomazone</i> | <i>360 g/l</i> | 1.6 l/ha | 3.2 l/ha | 2 per crop* | Before GS 49 | 14 days |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | <i>Fluopicolide</i> | <i>62.5 g/l</i> | 2.0 kg/ha | 2.0 kg/ha | 1 | 42 days before harvest | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | <i>Propamocarb</i> | <i>625 g/l</i> | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | <i>Pyridate</i> | <i>45 % w/w</i> | 2.0 kgs/ha | 4.0 kgs/ha | | 30 days pre harvest | 30 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Mancozeb</i> | <i>64 %w/w</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.5 g/m ² * | 0.5 g/m ² | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|---------------------|-----------------|-----------------------|------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Broccoli</i> | | | | | | | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | - | 7 days after transplanting | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------|---------------------|---------------|-----------------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Brussel sprouts</i> | | | | | | | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 6.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.2 l/ha | 0.2 l/ha | 1 / crop | 7 days before harvest | 7 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.l. | Herbicide | | | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| | | | | | <i>Clomazone</i> | <i>360 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 3.2 l/ha | 2 per crop* | Before GS 49 | 14 days |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | Insyst | 03249 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 250 g/ha | 1 | 21 days before harvest | 21 days |
| | | | | | <i>Acetamiprid</i> | <i>20 %w/w</i> | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 6.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Steward | 02603 | DuPont (UK) Ltd | Insecticide | | | 85 g/ha | 255 g/ha | 3 / crop | 1 day before harvest | |
| | | | | | <i>Indoxacarb</i> | <i>300 g/kg</i> | | | | | |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | | | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------------|---------------------|---------------|-----------------------------------|-----------------|--------------------------------|------------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Brussel sprouts</i> | | | | | | | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | - | 7 days after transplanting | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |
| <i>Bush/Cane fruit plantations</i> | | | | | | | | | | | |
| | Digital | 03147 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | | 365 days before first harvest | |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Guillotine | 03146 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 365 days before first harvest | |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Most Micro | 04837 | Sipcam U.K. Ltd | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 per year | N/A | |
| | | | | | <i>Pendimethalin</i> | <i>365 g/l</i> | | | | | |
| | Roundup Biactive | 02313 | Monsanto UK Ltd | Herbicide | | | 4 l/ha | 10 l/ha | | 28 days pre harvest | 28 days |
| | | | | | <i>Glyphosate</i> | <i>360 g/l</i> | | | | | |
| | Sumimax | 02764 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 365 days before first harvest | |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------------|---------------|-----------------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Cabbage - chinese</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 | 14 days | |
| | Butisan S | 04631 | BASF plc | Herbicide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.0 l/ha | 1.0 l/ha | - | Post emergence GS 16-18. 6-8 days after planting | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | <i>Metazachlor</i> | <i>500 g/l</i> | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | <i>S-Metolachlor</i> | <i>960 g/l</i> | 1.0 l/ha | 2.25 l/ha | - | 21 days before harvest | 21 days |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.l. | Herbicide | <i>Tebuconazole</i> | <i>250 g/l</i> | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | <i>Clomazone</i> | <i>360 g/l</i> | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | Movento | 04101 | Bayer CropScience Limited | Insecticide | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | 0.5 l/ha | 1.0 l/ha | 2 | 7 days before harvest | 7 days |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | <i>Spirotetramat</i> | <i>150 g/l</i> | 0.4 kgs/ha | 1.2 kgs/ha | 3 | 14 days before harvest | 14 days |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | <i>Pymetrozine</i> | <i>50 %w/w</i> | 0.5 l/ha | 1.0 l/ha | | 21 days before harvest | 21 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Difenoconazole</i> | <i>250 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------------|---------------|---------------------------------|-----------------|--------------------------------|----------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Cabbage - chinese</i> | | | | | | | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------|-----------------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Cabbage - head</i> | | | | | | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.2 l/ha | 0.2 l/ha | 1 / crop | 7 days before harvest | 7 days |
| Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | <i>Thiacloprid</i> | <i>480 g/l</i> | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | <i>S-Metolachlor</i> | <i>960 g/l</i> | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| Globazar SC | 04859 | Globachem nv | Fungicide | <i>Clomazone</i> | <i>360 g/l</i> | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| Infinito | 02532 | Bayer CropScience Limited | Fungicide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.6 l/ha | 3.2 l/ha | 2 per crop* | Before GS 49 | 14 days |
| Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | 0.5 kgs/m ³ | - | - | Pre-planting | |
| Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | 1.5 kgs/ha | 4.5 kgs/ha | | 14 days pre harvest | 14 days |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | <i>Mancozeb</i> | <i>64 %w/w</i> | | | | | |
| Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Esfenvalerate</i> | <i>25 g/l</i> | 10 g/m ³ | 10 g/m ³ | 1 | Pre planting / sowing | |
| | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| Wing P | 03590 | BASF plc | Herbicide | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | 4.0 l/ha | 4.0 l/ha | - | Before planting | |
| | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------|---------------------|---------------|-----------------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Calabrese</i> | | | | | | | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.2 l/ha | 0.2 l/ha | 1 / crop | 7 days before harvest | 7 days |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | <i>Thiacloprid</i> | <i>480 g/l</i> | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | <i>S-Metolachlor</i> | <i>960 g/l</i> | 1.0 l/ha | 2.25 l/ha | | 21 days before harvest | 21 days |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | <i>Tebuconazole</i> | <i>250 g/l</i> | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | <i>Clomazone</i> | <i>360 g/l</i> | 1.6 l/ha | 3.2 l/ha | 2 per crop* | Before GS 49 | 14 days |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | <i>Propamocarb</i> | <i>625 g/l</i> | 2.0 kg/ha | 2.0 kg/ha | 1 | 42 days before harvest | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | <i>Fluopicolide</i> | <i>62.5 g/l</i> | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | <i>Pyridate</i> | <i>45 % w/w</i> | 2.0 kgs/ha | 4.0 kgs/ha | | 30 days pre harvest | 30 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Mancozeb</i> | <i>64 %w/w</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.5 g/m ² * | 0.5 g/m ² | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------|---------------------|---------------|---------------------|-----------------|-----------------------|------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Calabrese</i> | | | | | | | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | - | 7 days after transplanting | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|--------------------------------------|------------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Carrots</i> | | | | | | | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Digital | 03147 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | | 28 days before harvest | 28 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 30 days before harvest | 30 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Fazor Star | 03729 | Arysta LifeScience Great Britain Ltd | Growth Regulator | | | 5.0 kg/ha | 5.0 kg/ha | 1 per crop | 21 days before harvest | |
| | | | | | <i>Maleic hydrazide</i> | <i>600 g/kg</i> | | | | | |
| | Guillotine | 03146 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 28 days before harvest | 28 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 40 days before harvest | 40 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Sencorex Flow | 04361 | Bayer CropScience Limited | Herbicide | | | 0.255 l/ha | 0.51 l/ha | 3 | Before GS 16 | 28 days |
| | | | | | <i>Metribuzin</i> | <i>600 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sumimax | 02764 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 28 days before harvest | 28 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|---------------------|-----------------|--------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Carrots</i> | | | | | | | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Fludioxonil</i> | 25.0 % w/w | | | | | |
| | | | | | <i>Cyprodinil</i> | 37.5 % w/w | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------|---------------------|---------------|----------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Cauliflowers</i> | | | | | | | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.2 l/ha | 0.2 l/ha | 1 / crop | 7 days before harvest | 7 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 1.0 l/ha | 2.25 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | | | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| | | | | | <i>Clomazone</i> | <i>360 g/l</i> | | | | | |
| | Globaztar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 3.2 l/ha | 2 per crop* | Before GS 49 | 14 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | | | 2.0 kg/ha | 2.0 kg/ha | 1 | 42 days before harvest | |
| | | | | | <i>Pyridate</i> | <i>45 % w/w</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | 2.0 kgs/ha | 4.0 kgs/ha | | 30 days pre harvest | 30 days |
| | | | | | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | | | | | |
| | | | | | <i>Mancozeb</i> | <i>64 %w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------|---------------------|---------------|-----------------------------------|-----------------|--|------------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Cauliflowers</i> | | | | | | | | | | | |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | | | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Esfenvalerate</i> | <i>25 g/l</i> | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | - | 7 days after transplanting | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|--------------------------------------|-----------------|-----------------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Celeriac</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 30 days before harvest | 30 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 8 weeks before harvest | 56 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Globaztar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 per year | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|---------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Celeriac</i> | | | | | | | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kg/ha | 1.2 kg/ha | 3 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Sencorex Flow | 04361 | Bayer CropScience Limited | Herbicide | | | 0.255 l/ha | 0.51 l/ha | 3 | Before GS 16 | 28 days |
| | | | | | <i>Metribuzin</i> | <i>600 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.0 kg/ha | 2.0 kg/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Venzar 500 SC | 04805 | DuPont (UK) Ltd | Herbicide | | | 0.4 l/ha | 0.4 l/ha | - | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>500 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|---------------------------------|-----------------|---------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Celeriac</i> | | | | | | | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | <i>Oxamyl</i> | <i>100 g/kg</i> | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------|---------------------|---------------|-------------------------|-----------------|---------------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Celery - field</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | 1 per crop | Before GS11 | 50 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 50 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Globaztar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 50 mls/ha | 150 mls/ha | 4 | 7 days pre harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 50 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 50 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 50 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 50 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 50 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kg/ha | 1.2 kg/ha | 3 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 50 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 50 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------|---------------------|---------------|---------------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Celery - field</i> | | | | | | | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 50 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| <i>Celery - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Globaztar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | - | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|--------------------------------------|-----------------|---|---------------------------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Cherries</i> | | | | | | | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.312 l/ha | 0.625 l/ha | - | 14 days before harvest | 14 days |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | <i>Thiacloprid</i> | <i>480 g/l</i> | 2.0 kg/ha | 8.0 kg/ha | 4 per year | 30 days before harvest | 30 days |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Mancozeb</i> | <i>75% w/w</i> | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Signum | 03799 | BASF plc | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.75 kg/ha | 2.25 kg/ha | | 7 days before harvest | 7 days |
| | Systhane 20 EW | 04944 | Dow AgroSciences Ltd | Fungicide | <i>Boscalid</i> <i>Pyraclostrobin</i> | <i>26.7 % w/w</i> <i>6.7 % w/w</i> | 0.45 l/ha | 0.9 l/ha | 2 | 14 days before harvest | 14 days |
| | | | | | <i>Myclobutanil</i> | <i>200 g/l</i> | | | | | |
| <i>Chicory</i> | | | | | | | | | | | |
| | Flexidor | 04963 | Dow AgroSciences Ltd | Herbicide | | | 0.2 l/ha | 0.2 l/ha | - | Before GS 16 | |
| | | | | | <i>Isoxaben</i> | <i>500 g/l</i> | | | | | |
| <i>Chives</i> | | | | | | | | | | | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | | 14 days pre harvest | 14 days |
| | | | | | <i>Metalaxyl M</i> <i>Mancozeb</i> | <i>4.0 %w/w</i> <i>64 %w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | <i>Cycloxydim</i> | <i>100 g/l</i> | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | <i>Fenhexamid</i> | <i>50 %w/w</i> | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | <i>Spinosad</i> | <i>480 g/l</i> | 200 ml/ha | 800 ml/ha | 4 per crop | 7 days before harvest | 7 days |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------|---------------------|---------------|------------------------------|-----------------|---|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Collards</i> | | | | | | | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | <i>S-Metolachlor</i> | <i>960 g/l</i> | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | <i>Clomazone</i> | <i>360 g/l</i> | 1.6 l/ha | 3.2 l/ha | 2 per crop* | Before GS 49 | 14 days |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Propamocarb</i> | <i>625 g/l</i> | 2.0 kg/ha | 2.0 kg/ha | 1 | 42 days before harvest | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | <i>Pyridate</i> | <i>45 % w/w</i> | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | 10 l/ha | 60 l/ha | 6 | 0 days | |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 10 g/m ³ | 10 g/m ³ | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| <i>Coriander</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | 0 days | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------|---------------------|--------------------------------------|---------------------|-----------------|-----------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Courgettes - field</i> | | | | | | | | | | | |
| Chamane | 05002 | UPL Europe Ltd | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | | | | 0.25 l/ha | 0.25 l/ha | - | Before 2 true leaf stage | |
| | | | | | <i>Clomazone</i> | <i>360 g/l</i> | | | | | |
| Globaztar SC | 05529 | Globachem nv | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Revus | 02789 | Syngenta UK Ltd | Fungicide | | | | 0.6 l/ha | 1.2 l/ha | 2 | 7 days pre harvest | 7 days |
| | | | | | <i>Mandipropamid</i> | <i>250 g/l</i> | | | | | |
| Signum | 03799 | BASF plc | Fungicide | | | | 1.5 kg/ha | 3.0 kg/ha | 2 per crop | 1 day before harvest | 1 day |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-------------------------------|---------------------|---------------|------------------------------|-----------------|---|-------------------|-------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Courgettes - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 80 mls/100 litres water | - | 3 | 3 days before harvest | 3 days |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 0.45 l/ha | 0.9 l/ha | 2 | 3 days before harvest | 3 days |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | <i>Thiacloprid</i> | <i>480 g/l</i> | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 0.4 kgs / ha | 1.2 kgs/ha | 3 | 3 days before harvest | 3 days |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | <i>Pymetrozine</i> | <i>50 %w/w</i> | 3.0 l/ha | 6.0 l/ha | - | 3 days before harvest | 3 days |
| | Revus | 02789 | Syngenta UK Ltd | Fungicide | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | 0.6 l/ha | 2.4 l/ha | 4 | 3 days pre harvest | 3 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Propamocarb</i> | <i>530 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | <i>Propamocarb</i> | <i>530 g/l</i> | 0.8 kg/ha | 2.4 kg/ha | 3 | 3 days before harvest | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-------------------------------|---------------------|---------------|---------------------------------|-----------------|---|-------------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Courgettes - protected</i> | | | | | | | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | 3 / crop | 1 day before harvest | 1 day |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| | Vivando | 05685 | BASF plc | Fungicide | | | 0.2 l/ha | 0.4 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Metrafenone</i> | <i>500 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| <i>Cranberries - field</i> | | | | | | | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Masai | 03585 | BASF plc | Insecticide | | | 1.5 kgs/ha | 1.5 kgs/ha | - | 7 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------------|---------------------|---------------|------------------------------|-----------------|---|-------------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Cranberries - protected</i> | | | | | | | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kgs/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | | | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m ³ | 10 g/m ³ | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|----------------------|---------------|-------------------------------------|-----------------|--|-------------------|---------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Cucumbers - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 80 mls / 100 litres water | - | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Apollo 50SC | 04610 | Adama Agricultural Solutions UK Ltd | Acaricide | | | 0.4 l/ha | 0.4 l/ha | 1 per year | 9 days before harvest | 9 days |
| | | | | | <i>Clofentezine</i> | <i>500 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.45 l/ha | 0.9 l/ha | 2 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Life Scientific Azox | 04405 | Life Scientific Limited | Fungicide | | | 80ml per 100L water | | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | 0.01kgs/5 litres water | - | - | - | 0 days |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kg/ha | 1.2 kgs/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Revus | 02789 | Syngenta UK Ltd | Fungicide | | | 0.6 l/ha | 2.4 l/ha | 4 | 3 days pre harvest | 3 days |
| | | | | | <i>Mandipropamid</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | 3 / crop | 1 day before harvest | 1 day |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|---------------------|---------------|---------------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Cucumbers - protected</i> | | | | | | | | | | | |
| | Vivando | 05685 | BASF plc | Fungicide | | | 0.2 l/ha | 0.4 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Metrafenone</i> | <i>500 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| <i>Damsons</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Dill</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| <i>Elderberries</i> | | | | | | | | | | | |
| | Batavia | 05905 | Bayer CropScience Limited | Insecticide | | | 0.75 l/ha | 1.5 l/ha | 2 | Post harvest application | |
| | | | | | <i>Spirotetramat</i> | <i>100 g/l</i> | | | | | |
| <i>Endives</i> | | | | | | | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 | 3 days pre harvest | 3 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------|---------------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Fennel</i> | | | | | | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|--------------------------------------|-----------------|-----------------------------|-----------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Garlic</i> | | | | | | | | | | | |
| | Amistar Opti | 03408 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | CIPC | 03066 | Arysta LifeScience Benelux SPRL | Herbicide | | | 1.8 l/ha | 5.4 l/ha | 3 per crop | GS 14 | |
| | | | | | <i>Chlorpropham</i> | <i>400 g/l</i> | | | | | |
| | Curator | 04504 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before 5th leaf visible | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 true leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Flexidor | 04963 | Dow AgroSciences Ltd | Herbicide | | | 0.2 l/ha | 0.2 l/ha | 1 per year | After transplanting or beginning of root formation | |
| | | | | | <i>Isoxaben</i> | <i>500 g/l</i> | | | | | |
| | Folio Gold | 02363 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 6.0 l/ha | 3 | 14 days before harvest | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>37.5 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 4 weeks before harvest | 28 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | 7 days before harvest | 7 days |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|----------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Garlic</i> | | | | | | | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | <i>Chlorothalonil</i> | <i>500 g/l</i> | 75 mls/ha | 225 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | 2.0 kg/ha | 2.0 kg/ha | 1 | Before 7 leaf stage | |
| | NEMGuard DE | 05368 | ECOSpray Ltd | Nematicide | <i>Pyridate</i> | <i>45 % w/w</i> | 20 kg/ha | 20 kg/ha | 1 | At drilling | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | <i>Garlic Extract</i> | <i>450 g/kg</i> | 75 mls/ha | 225 mls/ha | 3 | 21 days before harvest | 21 days |
| | Pyramin DF | 04264 | BASF plc | Herbicide | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | 1.4 kg/ha | 1.4 kg/ha | - | Before GS 13 | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | <i>Chloridazon</i> | <i>65% w/w</i> | 5.0 l/ha | 5.0 l/ha | - | Before GS 15 | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Prosulfocarb</i> | <i>800 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Starane Hi-Load H | 04549 | Dow AgroSciences Ltd | Herbicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.3 l/ha | 0.3 l/ha | | Before GS 15 | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | <i>Fluroxypyr</i> | <i>333 g/l</i> | 3.3 l/ha | 3.3 l/ha | 1 | Before crop emergence | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | <i>Pendimethalin</i> | <i>400 g/l</i> | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | Vertik | 04030 | Syngenta UK Ltd | Fungicide | <i>Pendimethalin</i> | <i>455 g/l</i> | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | Wing P | 03590 | BASF plc | Herbicide | <i>Chlorothalonil</i> | <i>500 g/l</i> | 4.0 l/ha | 4.0 l/ha | - | Pre-emergence / Pre-transplanting | |
| | | | | | <i>Azoxystrobin</i> | <i>100 g/l</i> | | | | | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product</i> | | | | | <i>Max Ind.</i> | <i>Max. Total</i> | <i>Max No of</i> | <i>Latest Time</i> | |
|----------------------------|----------------|---------------------------------|-----------------|--------------------------------|----------------|-----------------|-------------------|------------------|--|-------------|
| <i>Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Dose:</i> | <i>Dose:</i> | <i>Applic's:</i> | <i>of Applic:</i> | <i>PHI:</i> |
| <i>Gherkin - protected</i> | | | | | | | | | | |
| Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 3.0 l/ha | 6.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | 3 | 1 day before harvest | 1 day |
| | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| Vivando | 05685 | BASF plc | Fungicide | | | 0.2 l/ha | 0.4 l/ha | - | 3 days before harvest | 3 days |
| | | | | <i>Metrafenone</i> | <i>500 g/l</i> | | | | | |
| Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|-----------------------------|---------------------|-----------------|---|-----------------|------------------------------|-------------------------|----------------------------|---------------------------------|-------------|
| <i>Gooseberries - field</i> | | | | | | | | | | | |
| Batavia | 05905 | Bayer CropScience Limited | Insecticide | | | | 0.75 l/ha | 1.5 l/ha | 2 | Post harvest application | |
| | | | | | <i>Spirotetramat</i> | <i>100 g/l</i> | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | | 0.25 l/ha | 0.75 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Chikara Weed Cont | 04203 | ISK Biosciences Europe N.V. | Herbicide | | | | 120 g/ha | 120 g/ha | 1 per year | Before bud burst | |
| | | | | | <i>Flazasulfuron</i> | <i>25 % w/w</i> | | | | | |
| Corbel | 90438 | BASF plc | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | 3 | 2 weeks before harvest | 2 weeks |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| Manfil 75 WG | 03598 | Indofil Industries Limited | Fungicide | | | | 1.5 kg per 1000 litres water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75 % w/w</i> | | | | | |
| Manfil 80 WP | 03563 | Indofil Industries Limited | Fungicide | | | | 1.4 kg per 1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 1.5 kgs/ha | 1.5 kgs/ha | - | 7 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | | 0.25 kgs/100 litres water | - | 3 | Up to and including fruit swell | |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |
| Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Penncozeb 80 WP | 03609 | UPL Europe Ltd | Fungicide | | | | 1.4 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Penncozeb WDG | 03608 | UPL Europe Ltd | Fungicide | | | | 1.5 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>750 g/kg</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|---------------------------------|-----------------|--------------------------|-------------------|-------------------------|---|----------------------------|---|-------------|
| <i>Gooseberries - field</i> | | | | | | | | | | | |
| | Radspor 400 | 01797 | Arysta LifeScience Benelux SPRL | Fungicide | | | 73 mls/100 litres water | - | 5 | | |
| | Scala | 03916 | BASF plc | Fungicide | <i>Dodine</i> | <i>40 %w/w</i> | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Pyrimethanil</i> | <i>400 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Signum | 03799 | BASF plc | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 1.5 kgs/ha | 4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest) | - | 3 days before harvest, N/A for post harvest | |
| | Stroby WG | 05092 | BASF plc | Fungicide | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | 0.2 kg/ha | 0.6 kg/ha | - | 14 days before harvest | 14 days |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | <i>Boscalid</i> | <i>26.7 % w/w</i> | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Kresoxim Methyl</i> | <i>50% w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------------|---------------------|------------------------------|---------------------|--|--------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Gooseberries - protected</i> | | | | | | | | | | | |
| AQ 10 | 05801 | CBC (Europe) S.r.l. | Fungicide | <i>Ampelomyces Quisqualis</i> | 58 %w/w | 70 g/ha | - | 12 | - | | |
| Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | 375 g/kg | 1.0 kgs/ha | 3.0 kgs/ha | - | N/A | | |
| Scala | 03916 | BASF plc | Fungicide | <i>Pyrimethanil</i> | 400 g/l | 2.0 l/ha | 4.0 l/ha | 2 | 21 days before harvest | 21 days | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus subtilis</i> | 1015.1 g/l | 10 l/ha | 60 l/ha | 6 | | 0 days | |
| T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Trichoderma Asperellum (strain T34)</i> <i>Trichoderma Asperellum (strain T34)</i> | 12% w/w 12% w/w | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------|--------------------------------------|-----------------|-----------------------------|-----------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Herbs - field</i> | | | | | | | | | | |
| Acrobat MZ | 04059 | BASF plc | Fungicide | <i>Dimethomorph</i> | <i>75 g/kg</i> | 2.0 kg/ha | 6.0 kg/ha | - | 21 days before harvest | 21 days |
| | | | | <i>Mancozeb</i> | <i>667 g/kg</i> | | | | | |
| Amistar | 01351 | Syngenta UK Ltd | Fungicide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.0 l/ha | 4.0 l/ha | 4 | 14 days before harvest | 14 days |
| Bonalan | 05763 | Gowan Crop Protection Ltd | Herbicide | <i>Benfluralin</i> | <i>180 g/l</i> | 8.0 l/ha | 8.0 l/ha | - | Incorporation pre sowing/planting | |
| Chamane | 05002 | UPL Europe Ltd | Fungicide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| CIPC | 03066 | Arysta LifeScience Benelux SPRL | Herbicide | <i>Chlorpropham</i> | <i>400 g/l</i> | 1.9 l/ha | 3.8 l/ha | 2 per crop | 2 weeks after planting out | |
| Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | <i>Mancozeb</i> | <i>75% w/w</i> | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 28 days before harvest | 28 days |
| Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| Fenomenal | 04956 | Bayer CropScience Limited | Fungicide | <i>Fosetyl (-aluminium)</i> | <i>600 g/kg</i> | 2.5 kg/ha | 5.0 kg/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | <i>Fenamidone</i> | <i>60 g/kg</i> | | | | | |
| Globaztar SC | 05529 | Globachem nv | Fungicide | <i>Azoxystrobin</i> | <i>250 g/l</i> | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| Infinito | 02532 | Bayer CropScience Limited | Fungicide | <i>Propamocarb</i> | <i>625 g/l</i> | 1.6 l/ha | 1.6 l/ha | 1 per crop | Before GS 49 | 14 days |
| | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | 75 mls/ha | 150 mls/ha | 4 | 7 days pre harvest | 7 days |
| Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------|---------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Herbs - field</i> | | | | | | | | | | |
| Kerb Flo | 02830 | Dow AgroSciences Ltd | Herbicide | | | 1.875 l/ha | 1.875 l/ha | 1 | 24 days before harvest | 24 days |
| | | | | <i>Propyzamide</i> | <i>400 g/l</i> | | | | | |
| | | | | <i>Propyzamide</i> | <i>400 g/l</i> | | | | | |
| Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| PDM 330 EC | 03582 | BASF plc | Herbicide | | | 4.5 l/ha | 4.5 l/ha | 1 | 6 weeks pre harvest | |
| | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| Pendifin 400 SC | 05731 | Finchimica S.p.A | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 per crop | 6 weeks before harvest | 6 weeks |
| | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kgs/ha | 1.2 kgs/ha | 3 | 14 days before harvest | 14 days |
| | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| Scala | 03916 | BASF plc | Fungicide | | | 1.33 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| Signum | 03799 | BASF plc | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | - | 14 days before harvest | |
| | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------|---------------------|---------------|---------------------------------|-----------------|---------------------------|------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Herbs - field</i> | | | | | | | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 / crop | 6 weeks pre harvest | 6 weeks |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 3.3 l/ha | 3.3 l/ha | 1 / crop | 6 weeks pre harvest | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | 2 | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 ml/ha | 600 ml/ha | 3 per year | 3 days before harvest | 3 days |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 1.25 l/ha | 1.25 l/ha | - | Before transplanting | 35 days |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------------|---------------|---------------------------|----------------------|--|-----------------|---------------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Herbs - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | AQ 10 | 05801 | CBC (Europe) S.r.l. | Fungicide | | | 70 g/ha | - | 12 | - | |
| | | | | | <i>Ampelomyces Quisqualis</i> | <i>58 %w/w</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Dynamec | 02640 | Syngenta UK Ltd | Acaricide/Insecticid | | | 50 mls/product/100 litres water | - | 4 | 14 days before harvest | 14 days |
| | | | | | <i>Abamectin</i> | <i>18 g/l</i> | | | | | |
| | Fenomenal | 04956 | Bayer CropScience Limited | Fungicide | | | 2.5 kg/ha | 5.0 kg/ha | 2 per crop | BBCH 42 | |
| | | | | | <i>Fenamidone</i> | <i>60 g/kg</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>600 g/kg</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Kerb Flo | 02830 | Dow AgroSciences Ltd | Herbicide | | | 1.875 l/ha | 1.875 l/ha | 1 | 24 days pre harvest | 24 days |
| | | | | | <i>Propyzamide</i> | <i>400 g/l</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | 0.01kgs/5 litres water | - | 2 | - | 0 days |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |
| | Paraat | 03614 | BASF plc | Fungicide | | | 0.36 kg/ha | 0.72 kg/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Dimethomorph</i> | <i>500 g/kg</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kg/ha | 1.2 kg/ha | 3 | 14 days before harvest | 14 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------------|---------------|---------------------------------|-----------------|--------------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Herbs - protected</i> | | | | | | | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | 1.9 kg/ha | 5.7 kg/ha* | | 35 days before harvest | 35 days |
| | | | | | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | | | | | |
| | | | | | <i>Mancozeb</i> | <i>64 %w/w</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 1.33 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | - | 14 days before harvest | |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 ml/ha | 600 ml/ha | 3 per year | 3 days before harvest | 3 days |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------|---------------------|---------------|--------------------------------------|-----------------|-----------------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Horseradish</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| | Butisan S | 04631 | BASF plc | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | Pre-emergence | |
| | | | | | <i>Metazachlor</i> | <i>500 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 30 days before harvest | 30 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 8 weeks before harvest | 56 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Goltix 70 SC | 04668 | Adama Agricultural Solutions UK Ltd | Herbicide | | | 2.0 l/ha | 5.0 l/ha | - | Pre crop emergence | |
| | | | | | <i>Metamitron</i> | <i>700 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------|---------------------|---------------|---------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Horseradish</i> | | | | | | | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Rapsan 500SC | 05886 | Globachem nv | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | Pre-emergence | |
| | | | | | <i>Metazachlor</i> | <i>500 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.0 kg/ha | 2.0 kgs/ha | - | 14 days before harvest | |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | SL567A | 03106 | Syngenta UK Ltd | Fungicide | | | 1.3 l/ha | 1.3 l/ha | 1 | 30 days before harvest | 30 days |
| | | | | | <i>Metalaxyl M</i> | <i>465.2 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------|---------------------|---------------|---------------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Horseradish</i> | | | | | | | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | | | 3.3 l/ha | 3.3 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | Venzar 500 SC | 04805 | DuPont (UK) Ltd | Herbicide | | | 0.4 l/ha | 0.4 l/ha | - | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>500 g/l</i> | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | | | | | <i>Oxamyl</i> | <i>100 g/kg</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|---------------------------|---------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Jerusalem Artichoke</i> | | | | | | | | | | | |
| Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| Corbel | 90438 | BASF plc | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| Decis | 02693 | Bayer CropScience Limited | Insecticide | | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Markate 50 | 05001 | Globachem nv | Insecticide | | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| Ravane | 05536 | Globachem nv | Insecticide | | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Stratos Ultra | 04936 | BASF plc | Herbicide | | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|---------------|---------------------|-----------------|------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Jerusalem Artichoke</i> | | | | | | | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | | | | | <i>Oxamyl</i> | <i>100 g/kg</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-------------|---------------------|---------------|-------------------------------------|-----------------|--|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Kale</i> | | | | | | | | | | | |
| | Anthem | 04761 | Adama Agricultural Solutions UK Ltd | Herbicide | | | 3.3 l/ha | 3.3 l/ha | - | | |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.2 l/ha | 0.2 l/ha | 1 / crop | 7 days before harvest | 7 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Drilled crops- pre-emergence. Or within 5 days transplanting | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | | | 0.25 l/ha | 0.25 l/ha | - | Before 6 true leaf stage | |
| | | | | | <i>Clomazone</i> | <i>360 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 3.2 l/ha | 2 per crop* | Before GS 49 | 14 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | | | 2.0 kg/ha | 2.0 kg/ha | 1 | 42 days before harvest | |
| | | | | | <i>Pyridate</i> | <i>45 % w/w</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kg/ha | 1.2 kg/ha | 3 | 14 days before harvest | 14 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | | | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------|---------------------|---------------|-----------------------------------|-----------------|---|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Kale</i> | | | | | | | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 12 mls/product/plant | - | 1 | Pre planting - 6 leaf stage | |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| <i>Kohl-rabi</i> | | | | | | | | | | | |
| | Butisan S | 04631 | BASF plc | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | Pre-emergence | |
| | | | | | <i>Metazachlor</i> | <i>500 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Movento | 04101 | Bayer CropScience Limited | Insecticide | | | 0.5 l/ha | 1.0 l/ha | 2 | 7 days before harvest | 7 days |
| | | | | | <i>Spirotetramat</i> | <i>150 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | | | 165 mls/ha | 330 mls/ha | 2 | 7 days pre harvest | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------|---------------------|---------------|----------------------------|-----------------|---------------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Leeks</i> | | | | | | | | | | | |
| | Acrobat MZ | 04059 | BASF plc | Fungicide | | | 2.0 kg/ha | 6.0 kg/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Mancozeb</i> | <i>667 g/kg</i> | | | | | |
| | | | | | <i>Dimethomorph</i> | <i>75 g/kg</i> | | | | | |
| | Basagran | 02341 | BASF plc | Herbicide | | | 0.75 l/ha | 1.5 l/ha | - | Before 3 true leaf stage | |
| | | | | | <i>Bentazone</i> | <i>480 g/l</i> | | | | | |
| | Basagran SG | 05192 | BASF plc | Herbicide | | | 0.4 kg/ha | 0.8 kg/ha | - | Before 3 true leaf stage | |
| | | | | | <i>Bentazone</i> | <i>870 g/kg</i> | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Butisan S | 04631 | BASF plc | Herbicide | | | 1.5 l/ha | 1.5 l/ha | - | 5-21 days after planting | |
| | | | | | <i>Metazachlor</i> | <i>500 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 250 mls/ha | 750 mls/ha | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before 5th leaf visible | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Flexidor | 04963 | Dow AgroSciences Ltd | Herbicide | | | 0.3 l/ha | 0.3 l/ha | 1 per year | After transplanting | |
| | | | | | <i>Isoxaben</i> | <i>500 g/l</i> | | | | | |
| | Folio Gold | 02363 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 6.0 l/ha | 3 | 14 days before harvest | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>37.5 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 100 mls/ha | 200 mls/ha | 2 per crop | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | | | 2.0 kg/ha | 2.0 kg/ha | 1 | 28 days before harvest | |
| | | | | | <i>Pyridate</i> | <i>45 % w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------|---------------------|---------------|---------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Leeks</i> | | | | | | | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | NEMGuard DE | 05368 | ECOspray Ltd | Nematicide | | | 20 kg/ha | 20 kg/ha | 1 | At drilling | |
| | | | | | <i>Garlic Extract</i> | <i>450 g/kg</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 100 mls/ha | 200 mls/ha | 2 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 15 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Starane Hi-Load H | 04549 | Dow AgroSciences Ltd | Herbicide | | | 0.3 l/ha | 0.3 l/ha | | 28 days before harvest | 28 days |
| | | | | | <i>Fluroxypyr</i> | <i>333 g/l</i> | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | - | Pre-emergence / Pre-transplanting | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------|----------------------|---------------|--------------------------------------|-----------------|------------------------------------|-----------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Lettuce - field</i> | | | | | | | | | | | |
| | Acrobat MZ | 04059 | BASF plc | Fungicide | | | 2.0 kg/ha | 6.0 kg/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Dimethomorph</i> | <i>75 g/kg</i> | | | | | |
| | | | | | <i>Mancozeb</i> | <i>667 g/kg</i> | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Fenomenal | 04956 | Bayer CropScience Limited | Fungicide | | | 2.5 kg/ha | 5.0 kg/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Fenamidone</i> | <i>60 g/kg</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>600 g/kg</i> | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 | 3 days pre harvest | 3 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Globazar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 1.0 l/ha | 1 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | Before GS 49 | 14 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Kerb Flo | 02830 | Dow AgroSciences Ltd | Herbicide | | | 1.875 l/ha | 1.875 l/ha | 1 | 24 days before harvest | 24 days |
| | | | | | <i>Propyzamide</i> | <i>400 g/l</i> | | | | | |
| | Life Scientific Azox | 04405 | Life Scientific Limited | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST</i> | <i>2% w/w</i> | | | | | |
| | | | | | <i>BIPESCO 5/F52</i> | | | | | | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | | | 4.5 l/ha | 4.5 l/ha | 1 | 6 weeks pre harvest | |
| | | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| | Pendifin 400 SC | 05731 | Finchimica S.p.A | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 per crop | 6 weeks before harvest | 6 weeks |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------|---------------------|---------------|---------------------------|-----------------|-----------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Lettuce - field</i> | | | | | | | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.4 kg/ha | 1.2 kg/ha | 3 | 7 days before harvest | 7 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 1.33 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 / crop | 6 weeks pre harvest | 6 weeks |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 3.3 l/ha | 3.3 l/ha | 1 / crop | 6 weeks pre harvest | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 0.7 kg/ha | 1.4 kg/ha | 2 | 7 days before harvest | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | Systhane 20 EW | 04944 | Dow AgroSciences Ltd | Fungicide | | | 0.3 l/ha | 0.6 l/ha | 2 | 14 days before harvest | 14 days |
| | | | | | <i>Myclobutanil</i> | <i>200 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | 2 / crop | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 ml/ha | 600 ml/ha | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 1.25 l/ha | 1.25 l/ha | - | Before transplanting | 35 days |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|----------------------------|----------------------|-----------------|--|-----------------|---------------------------------------|-------------------------|----------------------------|-------------------------------|-----------------|
| <i>Lettuce - protected</i> | | | | | | | | | | | |
| Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | | 1.0 l/ha | 1.0 l/ha | 1 | 7 days post transplanting | 7 days post tra |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Dynamec | 02640 | Syngenta UK Ltd | Acaricide/Insecticid | | | | 50 mls/product/100 litres water | - | 4 | 14 days before harvest | 14 days |
| | | | | | <i>Abamectin</i> | <i>18 g/l</i> | | | | | |
| Fenomenal | 04956 | Bayer CropScience Limited | Fungicide | | | | 2.5 kg/ha | 5.0 kg/ha | 2 per crop | BBCH 42 | |
| | | | | | <i>Fenamidone</i> | <i>60 g/kg</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>600 g/kg</i> | | | | | |
| Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | | 250 g/ha | 500 g/ha | 2 | 3 days pre harvest | 3 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| Globazar SC | 04859 | Globachem nv | Fungicide | | | | 1.0 l/ha | 1.0 l/ha | 1 per crop | 14 days after transplanting | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Kerb Flo | 02830 | Dow AgroSciences Ltd | Herbicide | | | | 1.875 l/ha | 1.875 l/ha | 1 | 24 days pre harvest | 24 days |
| | | | | | <i>Propyzamide</i> | <i>400 g/l</i> | | | | | |
| Life Scientific Azox | 04405 | Life Scientific Limited | Fungicide | | | | 1.0 l/ha | 1.0 l/ha | 1 per crop | 7 days after transplanting | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| Paraat | 03614 | BASF plc | Fungicide | | | | 0.36 kg/ha | 0.72 kg/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Dimethomorph</i> | <i>500 g/kg</i> | | | | | |
| Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | | 0.4 kg/ha | 1.2 kgs/ha | 3 | 14 days before harvest | 14 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | | 1.9 kg/ha | 5.7 kg/ha* | | 35 days before harvest | 35 days |
| | | | | | <i>Mancozeb</i> | <i>64 %w/w</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | | | | | |
| Scala | 03916 | BASF plc | Fungicide | | | | 1.33 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|---------------|------------------------------|-----------------|--|--|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Lettuce - protected</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.7 kg/ha | 1.4 kg/ha | 2 | 7 days before harvest | 7 days |
| | Systhane 20 EW | 04944 | Dow AgroSciences Ltd | Fungicide | <i>Fludioxonil</i> <i>Cyprodinil</i> | <i>25.0 % w/w</i> <i>37.5 % w/w</i> | 0.3 l/ha | 0.6 l/ha | 2 | 14 days before harvest | 14 days |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Myclobutanil</i> | <i>200 g/l</i> | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | <i>Trichoderma Asperellum (strain T34)</i> <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> <i>12% w/w</i> | 1.5 kgs/ha | 3.0 kgs/ha | 2 / crop | 3 days before harvest | 3 days |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | <i>Fenhexamid</i> <i>Spinosad</i> | <i>50 %w/w</i> <i>480 g/l</i> | 200 ml/ha | 600 ml/ha | 3 per crop | 3 days before harvest | 3 days |
| <i>Linseed</i> | | | | | | | | | | | |
| | Butisan S | 04631 | BASF plc | Herbicide | <i>Metazachlor</i> | <i>500 g/l</i> | 1.5 l/ha | 1.5 l/ha | - | Before GS 19 | |
| | Devrinol | 91627 | UPL Europe Ltd | Herbicide | <i>Napropamide</i> | <i>450 g/l</i> | 2.8 l/ha | 2.8 l/ha | - | Pre-emergence | |
| | Rapsan 500SC | 05886 | Globachem nv | Herbicide | <i>Metazachlor</i> | <i>500 g/l</i> | 1.5 l/ha | 1.5 l/ha | - | GS 19 | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|---------------------------|-----------------|---|-------------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Loganberries - field</i> | | | | | | | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Masai | 03585 | BASF plc | Insecticide | | | 375 gms/ha | 375 g/ha | 1 / year | 21 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | | | 0.8 l/ha | 1.6 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Carfentrazone-ethyl</i> | <i>60 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------------|---------------------|---------------|-------------------------------------|-----------------|---|----------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Loganberries - protected</i> | | | | | | | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Tebuconazole</i> | 250 g/l | 1.0 l/ha | 3.0 l/ha | - | N/A | |
| | Masai | 03585 | BASF plc | Insecticide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | 375 g/kg | 375 gms/ha | 375 gms/ha | 1 / year | 21 days before harvest | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | <i>Tebufenpyrad</i> | 200 g/kg | 0.5 kgs/m ³ | | | pre-planting | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | 2% w/w | 10 l/ha | 60 l/ha | 6 | - | 0 days |
| | Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | <i>Bacillus subtilis</i> | 1015.1 g/l | 0.8 l/ha | 1.6 l/ha | - | 21 days before harvest | 21 days |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | <i>Carfentrazone-ethyl</i> | 60 g/l | 10 g/m ³ | 10 g/m ³ | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | 12% w/w | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | 12% w/w | | | | | |
| <i>Lupins</i> | | | | | | | | | | | |
| | Anthem | 04761 | Adama Agricultural Solutions UK Ltd | Herbicide | | | 3.75 l/ha | 3.75 l/ha | | Pre crop emergence | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | <i>Pendimethalin</i> | 400 g/l | 4.5 l/ha | 4.5 l/ha | 1 | Before crop emergence | |
| | Pendifin 400 SC | 05731 | Finchimica S.p.A | Herbicide | <i>Pendimethalin</i> | 330 g/l | 3.75 l/ha | 3.75 l/ha | 1 per crop | Pre-crop emergence | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Pendimethalin</i> | 400 g/l | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | <i>Bacillus subtilis</i> | 1015.1 g/l | 3.3 l/ha | 3.3 l/ha | 1 / crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | 455 g/l | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product</i> | | | | | | <i>Max Ind.</i> | <i>Max. Total</i> | <i>Max No of</i> | <i>Latest Time</i> | |
|---------------------------------------|---------------|---------------------------|-----------------|--------------------------|-------------------|-----------------|-------------------|------------------|--|-------------|
| <i>Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Dose:</i> | <i>Dose:</i> | <i>Applic's:</i> | <i>of Applic:</i> | <i>PHI:</i> |
| <i>Maize - Forage - open ground</i> | | | | | | | | | | |
| Comet | 02354 | BASF plc | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 | Tasel/silk visible stage | |
| | | | | <i>Pyraclostrobin</i> | <i>250 g/l</i> | | | | | |
| Comet 200 | 03696 | BASF plc | Fungicide | | | 1.25 l/ha | 2.5 l/ha | 2 | Tasel/silk visible stage (BBCH 51-59) | |
| | | | | <i>Pyraclostrobin</i> | <i>200 g/l</i> | | | | | |
| Modem | 02355 | BASF plc | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | Tasel/silk visible stage | |
| | | | | <i>Pyraclostrobin</i> | <i>250 g/l</i> | | | | | |
| Modem 200 | 03695 | BASF plc | Fungicide | | | 1.25 l/ha | 2.5 l/ha | 2 | Tasel/silk visible stage (BBCH 51-59) | |
| | | | | <i>Pyraclostrobin</i> | <i>200 g/l</i> | | | | | |
| <i>Maize - Forage - under plastic</i> | | | | | | | | | | |
| Comet 200 | 03696 | BASF plc | Fungicide | | | 1.25 l/ha | 2.5 l/ha | 2 | Tasel/silk visible stage (BBCH 51-59) | |
| | | | | <i>Pyraclostrobin</i> | <i>200 g/l</i> | | | | | |
| Modem | 02355 | BASF plc | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | Tasel/silk visible stage | |
| | | | | <i>Pyraclostrobin</i> | <i>250 g/l</i> | | | | | |
| Modem 200 | 03695 | BASF plc | Fungicide | | | 1.25 l/ha | 2.5 l/ha | 2 per crop | Tassel/silk visible stage (BBCH 51-59) | |
| | | | | <i>Pyraclostrobin</i> | <i>200 g/l</i> | | | | | |
| <i>Maize - Grain - open ground</i> | | | | | | | | | | |
| Comet | 02354 | BASF plc | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 | Tasel/silk visible stage | |
| | | | | <i>Pyraclostrobin</i> | <i>250 g/l</i> | | | | | |
| <i>Mange tout peas</i> | | | | | | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Mangolds</i> | | | | | | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------------|---------------|------------------------------|-----------------|--------------------------------|-------------------|--------------------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Marrows</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 3 mls per sq m | 6 mls per sq m | - | Before transplanting | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Melon - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| <i>Mint</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| <i>Mushrooms</i> | | | | | | | | | | | |
| | Pyrethrum 5 EC | 92317 | Agropharm Ltd | Insecticide | | | 0.1 ml product per Sq m growing area | N/A | N/A | 1 day before harvest | 1 day |
| | | | | | <i>Pyrethrins</i> | <i>50 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|---------------------|---------------|---------------------------|-----------------|------------------------|-------------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Oilseed Rape - spring</i> | | | | | | | | | | | |
| | Galera | 05188 | Dow AgroSciences Ltd | Herbicide | | | 0.35 l/ha | 0.35 l/ha | - | Before flower buds visible above crop canopy | |
| | | | | | <i>Picloram</i> | <i>67 g/l</i> | | | | | |
| | | | | | <i>Clopyralid</i> | <i>267 g/l</i> | | | | | |
| | Proline | 03786 | Bayer CropScience Limited | Fungicide | | | 0.7 l/ha | 1.4 l/ha | - | 56 days before harvest | 56 days |
| | | | | | <i>Prothioconazole</i> | <i>250 g/l</i> | | | | | |
| | Prosaro | 02106 | Bayer CropScience Limited | Fungicide | | | 1.0 l/ha | 2.0 l/ha | | 56 days before harvest | 56 days |
| | | | | | <i>Prothioconazole</i> | <i>12.8 % w/w</i> | | | | | |
| | | | | | <i>Tebuconazole</i> | <i>12.8 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|-----------------------------------|-----------------|-----------------------|-----------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Onions</i> | | | | | | | | | | | |
| | Amistar Opti | 03408 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Curator | 04504 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before 5th leaf visible | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Digital | 03147 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | - | 42 days before harvest | 42 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 true leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Flexidor | 04963 | Dow AgroSciences Ltd | Herbicide | | | 0.2 l/ha | 0.2 l/ha | 1 per year | After transplanting or beginning of root formation | |
| | | | | | <i>Isoxaben</i> | <i>500 g/l</i> | | | | | |
| | Folio Gold | 02363 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 6.0 l/ha | 3 | 14 days before harvest | |
| | | | | | <i>Metalaxyl M</i> | <i>37.5 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Guillotine | 03146 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 42 days before harvest | 42 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | 7 days before harvest | 7 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|-----------------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Onions</i> | | | | | | | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 225 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | NEMGuard DE | 05368 | ECOSpray Ltd | Nematicide | | | 20 kg/ha | 20 kg/ha | 1 | At drilling | |
| | | | | | <i>Garlic Extract</i> | <i>450 g/kg</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 225 mls/ha | 3 | 21 days before harvest | 21 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Pyramin DF | 04264 | BASF plc | Herbicide | | | 1.4 kg/ha | 1.4 kg/ha | - | Before GS 13 | |
| | | | | | <i>Chloridazon</i> | <i>65% w/w</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 15 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Starane Hi-Load H | 04549 | Dow AgroSciences Ltd | Herbicide | | | 0.3 l/ha | 0.3 l/ha | | Before GS 15 | |
| | | | | | <i>Fluroxypyr</i> | <i>333 g/l</i> | | | | | |
| | Sumimax | 02764 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 42 days before harvest | 42 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Valbon | 03411 | Certis Europe BV | Fungicide | | | 1.6 kg/ha | 9.6 kg/ha | 6 per crop | 28 days pre-harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>700 g/kg</i> | | | | | |
| | | | | | <i>Benthiavalicarb</i> | <i>17.5 g/kg</i> | | | | | |
| | Vertik | 04030 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>100 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | - | Pre-emergence / Pre-transplanting | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------|---------------------|---------------|---------------------------|-----------------|-----------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Onions - salad</i> | | | | | | | | | | | |
| | Acrobat MZ | 04059 | BASF plc | Fungicide | | | 2.0 kg/ha | 10.0 kg/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Mancozeb</i> | <i>667 g/kg</i> | | | | | |
| | | | | | <i>Dimethomorph</i> | <i>75 g/kg</i> | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 21 days before harvest | 21 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Amistar Opti | 03408 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Basagran | 02341 | BASF plc | Herbicide | | | 1.0 l/ha | 2.0 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Bentazone</i> | <i>480 g/l</i> | | | | | |
| | Basagran SG | 05192 | BASF plc | Herbicide | | | 0.55 kg/ha | 1.1 kg/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Bentazone</i> | <i>870 g/kg</i> | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Curator | 04504 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | | Before 3rd true leaf | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 21 days | |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |
| | Folio Gold | 02363 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 6.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>37.5 g/l</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------|---------------------|---------------|----------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Onions - salad</i> | | | | | | | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 150 mls/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Manfil 75 WG | 03598 | Indofil Industries Limited | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75 % w/w</i> | | | | | |
| | Manfil 80 WP | 03563 | Indofil Industries Limited | Fungicide | | | 2.0 kg/ha | 8.0 kg/ha | 4 per crop | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 150 mls/ha | 2 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Penncozeb 80 WP | 03609 | UPL Europe Ltd | Fungicide | | | 2.0 kg/ha | 8.0 kg/ha | 4 per crop | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| | Penncozeb WDG | 03608 | UPL Europe Ltd | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>750 g/kg</i> | | | | | |
| | Pyramin DF | 04264 | BASF plc | Herbicide | | | 1.4 kg/ha | 1.4 kg/ha | - | Before GS 13 | |
| | | | | | <i>Chloridazon</i> | <i>65% w/w</i> | | | | | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | | 14 days pre harvest | 14 days |
| | | | | | <i>Mancozeb</i> | <i>64 %w/w</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 13 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------|---------------------|---------------|---------------------------------|-----------------|-----------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Onions - salad</i> | | | | | | | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.5 kg/ha | 3.0 kg/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 2.9 l/ha | 2.9 l/ha | 1 per crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 21 days before harvest | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | Vertik | 04030 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>100 g/l</i> | | | | | |
| | Wing P | 03590 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | - | Pre-emergence | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------|-----------------------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Orchards - top fruit</i> | | | | | | | | | | |
| Digital | 03147 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | | 365 days before first harvest | |
| | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| Dow Shield 100 | 03861 | Dow AgroSciences Ltd | Herbicide | | | 2.0 l/ha | 2.0 l/ha | - | - | |
| | | | | <i>Clopyralid</i> | <i>100 g/l</i> | | | | | |
| Guillotine | 03146 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 365 days before first harvest | |
| | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| Most Micro | 04837 | Sipcam U.K. Ltd | Herbicide | | | 3.75 l/ha | 3.75 l/ha | 1 per year | N/A | |
| | | | | <i>Pendimethalin</i> | <i>365 g/l</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| Sumimax | 02764 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 365 days before first harvest | |
| | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| Vivendi 200 | 05122 | UPL Europe Ltd | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | - | |
| | | | | <i>Clopyralid</i> | <i>200 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|---------------|---------------------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Parsley - field</i> | | | | | | | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 | 3 days pre harvest | 3 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Revus | 02789 | Syngenta UK Ltd | Fungicide | | | 0.6 l/ha | 1.2 l/ha | 2 | 7 days pre harvest | 7 days |
| | | | | | <i>Mandipropamid</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| <i>Parsley - protected</i> | | | | | | | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Revus | 02789 | Syngenta UK Ltd | Fungicide | | | 0.6 l/ha | 0.6 l/ha | 1 | 7 days pre harvest | 7 days |
| | | | | | <i>Mandipropamid</i> | <i>250 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------|---------------------|---------------|--------------------------------------|-----------------|-----------------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Parsley Root</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 30 days before harvest | 30 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 8 weeks before harvest | 56 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Globazar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------|---------------------|---------------|---------------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Parsley Root</i> | | | | | | | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | | | 3.3 l/ha | 3.3 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | | | | | <i>Oxamyl</i> | <i>100 g/kg</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|--------------------------------------|------------------|-----------------------------|-----------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Parsnips</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Digital | 03147 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | - | 28 days before harvest | 28 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 30 days before harvest | 30 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Fazor Star | 03729 | Arysta LifeScience Great Britain Ltd | Growth Regulator | | | 5.0 kg/ha | 5.0 kg/ha | 1 per crop | 21 days before harvest | |
| | | | | | <i>Maleic hydrazide</i> | <i>600 g/kg</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3 l/ha | 3 l/ha | 1 | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Globazar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 per year | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Goltix 70 SC | 04668 | Adama Agricultural Solutions UK Ltd | Herbicide | | | 2.0 l/ha | 5.0 l/ha | - | Pre crop emergence | |
| | | | | | <i>Metamitron</i> | <i>700 g/l</i> | | | | | |
| | Guillotine | 03146 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 28 days before harvest | 28 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | 0.01kgs/5 litres water | - | 2 | - | 0 days |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|-----------------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|--------------------------------------|-------------|
| <i>Parsnips</i> | | | | | | | | | | | |
| | Movento | 04101 | Bayer CropScience Limited | Insecticide | | | 0.3 l/ha | 0.6 l/ha | 2 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Spirotetramat</i> | <i>150 g/l</i> | | | | | |
| | Nativo 75 WG | 03226 | Bayer CropScience Limited | Fungicide | | | 0.3 kg/ha | 0.9 kg/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Trifloxystrobin</i> | <i>250 g/kg</i> | | | | | |
| | | | | | <i>Tebuconazole</i> | <i>500 g/kg</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.0 kg/ha | 2.0 kgs/ha | - | 14 days before harvest | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | SL567A | 03106 | Syngenta UK Ltd | Fungicide | | | 1.3 l/ha | 1.3 l/ha | 1 | no later than 6 weeks after drilling | 30 days |
| | | | | | <i>Metalaxyl M</i> | <i>465.2 g/l</i> | | | | | |
| | Sumimax | 02764 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 28 days before harvest | 28 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxyastrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-------------------------|---------------------|---------------|--|------------------|---------------------------|-------------------|--------------------------|-------------------------|----------------------------|--|-------------|
| <i>Pears</i> | | | | | | | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.375 l/ha | 0.75 l/ha | - | 14 days before harvest | 14 days |
| | Curenox 50 WP | 05829 | Industrias Quimicas Del Valles S.A (IQV, S.A.) | Fungicide | <i>Thiacloprid</i> | <i>480 g/l</i> | 2.0 kg/ha | 8.0 kg/ha | - | Before GS 07 | |
| | Exilis | 04702 | Fine Agrochemicals Ltd | Growth Regulator | <i>Copper oxychloride</i> | <i>87.8% w/w</i> | 7.5 l/ha * | 7.5 l/ha | 1 per year | Up to fruitlet diameter of 15 mm | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | <i>6-benzyladenine</i> | <i>20.0 g/l</i> | | | | | |
| | | | | | <i>6-benzyladenine</i> | <i>20.0 g/l</i> | 0.6 l/ha | 1.8 l/ha | 3 | After final harvest but before final leaf fall | |
| | Masai | 03585 | BASF plc | Insecticide | <i>Tebuconazole</i> | <i>250 g/l</i> | 125 gms/1 m crown height | | 1 / year | 21 days before harvest | |
| | Scala | 03916 | BASF plc | Fungicide | <i>Tebuconazole</i> | <i>250 g/l</i> | 1.125 l/ha | 4.5 l/ha | 4 per year | 56 days before harvest | 56 days |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | <i>Pyrimethanil</i> | <i>400 g/l</i> | 0.25 l/ha | 1.0 l/ha | | 28 days before harvest | 28 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Difenoconazole</i> | <i>250 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.8 kg/ha | 2.4 kg/ha | 3 | 3 days before harvest | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | Talius | 04604 | DuPont (UK) Ltd | Fungicide | <i>Proquinazid</i> | <i>200 g/l</i> | 0.25 l/ha | 0.5 l/ha | 2 per year | 49 days before harvest | 49 days |
| <i>Peas - combining</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | <i>Esfenvalerate</i> | <i>25 g/l</i> | 330 mls/ha | 330 mls/ha | 2 | 14 days pre harvest | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-------------------------------------|---------------------|---------------|-----------------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Peas - vining</i> | | | | | | | | | | | |
| | Digital | 03147 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | - | 42 days before harvest | 42 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Guillotine | 03146 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 42 days before harvest | 42 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Stallion Sync Tech | 05483 | FMC Chemical s.p.r.l. | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 per crop | Pre-crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>333 g/l</i> | | | | | |
| | | | | | <i>Clomazone</i> | <i>30 g/l</i> | | | | | |
| | Sumi-Alpha | 02334 | Sumitomo Chemical Agro Europe SAS | Insecticide | | | 330 mls/ha | 330 mls/ha | 2 | 14 days pre harvest | |
| | | | | | <i>Esfenvalerate</i> | <i>25 g/l</i> | | | | | |
| | Sumimax | 02764 | Sumitomo Chemical Agro Europe SAS | Herbicide | | | 100 mls/ha | 100 mls/ha | 1 | 42 days before harvest | 42 days |
| | | | | | <i>Flumioxazin</i> | <i>300 g/l</i> | | | | | |
| <i>Peppers - Chilli - protected</i> | | | | | | | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|---------------|---------------------------|----------------------|-----------------------------|-------------------|---------------------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Peppers - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.45 l/ha | 0.9 l/ha | 2 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Dynamec | 02640 | Syngenta UK Ltd | Acaricide/Insecticid | | | 50 mls/product/100 litres water | - | 5 | 3 days before harvest | 3 days |
| | | | | | <i>Abamectin</i> | <i>18 g/l</i> | | | | | |
| | Globaztar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | 0.01kgs/5 litres water | - | - | - | 0 days |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.04 kgs/100 l water | - | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 3.0 l/ha | 12.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | - | 7 days before harvest | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | 3 / crop | 1 day before harvest | 1 day |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|---------------------|---------------|--|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Peppers - protected</i> | | | | | | | | | | | |
| | Vivando | 05685 | BASF plc | Fungicide | | | 0.3 l/ha | 0.6 l/ha | - | 3 days before harvest | 3 days |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | <i>Metrafenone</i> | <i>500 g/l</i> | 1.0 l/ha | 2.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| <i>Plums</i> | | | | | | | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Signum | 03799 | BASF plc | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.75 kg/ha | 2.25 kg/ha | - | 7 days before harvest | 7 days |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| <i>Potatoes - seed crops</i> | | | | | | | | | | | |
| | Amistar Opti | 03408 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 7 days before harvest | 7 days |
| | Curator | 04504 | Syngenta UK Ltd | Fungicide | <i>Chlorothalonil</i> | <i>400 g/l</i> | 1.0 l/ha | 3.0 l/ha | - | 7 days before harvest | 7 days |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Curenox 50 WP | 05829 | Industrias Quimicas Del Valles S.A (IQV, S.A.) | Fungicide | <i>Azoxystrobin</i> | <i>80 g/l</i> | 1.0 kg/ha | 6.0 kg/ha | - | 8 days before harvest | 8 days |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Copper oxychloride</i> | <i>87.8% w/w</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Vertik | 04030 | Syngenta UK Ltd | Fungicide | | | 0.8 l/ha | 2.4 l/ha | - | 7 days pre-harvest | 7 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|---------------------|---------------|--|-----------------|--------------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Potatoes - ware crops</i> | | | | | | | | | | | |
| | Amistar Opti | 03408 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 7 days before harvest | 7 days |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | Curator | 04504 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 7 days before harvest | 7 days |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | Curenox 50 WP | 05829 | Industrias Quimicas Del Valles S.A (IQV, S.A.) | Fungicide | | | 1.0 kg/ha | 6.0 kg/ha | - | 8 days before harvest | 8 days |
| | | | | | <i>Copper oxychloride</i> | <i>87.8% w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Vertik | 04030 | Syngenta UK Ltd | Fungicide | | | 0.8 l/ha | 2.4 l/ha | - | 7 days pre-harvest | 7 days |
| | | | | | <i>Azoxystrobin</i> | <i>100 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| <i>Pumpkins - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 3 mls per sq m | 6 mls per sq m | - | Before transplanting | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | - | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------|----------------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Radicchio</i> | | | | | | | | | | |
| Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 | 3 days pre harvest | 3 days |
| | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|---------------------------|-----------------|-----------------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Radish</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Butisan S | 04631 | BASF plc | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | Pre-emergence | |
| | | | | | <i>Metazachlor</i> | <i>500 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 8 weeks before harvest | 56 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Globaztar SC | 04859 | Globachem nv | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 per year | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Rapsan 500SC | 05886 | Globachem nv | Herbicide | | | 1.0 l/ha | 1.0 l/ha | - | | |
| | | | | | <i>Metazachlor</i> | <i>500 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------|---------------------|---------------|---------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Radish</i> | | | | | | | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | SL567A | 03106 | Syngenta UK Ltd | Fungicide | | | 1.3 l/ha | 1.3 l/ha | 1 | 30 days before harvest | 30 days |
| | | | | | <i>Metalaxyl M</i> | <i>465.2 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | | | | | <i>Oxamyl</i> | <i>100 g/kg</i> | | | | | |
| <i>Rape - fodder</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|-------------------------------------|---------------------|-----------------|----------------------------|-------------------|-------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Raspberries - field</i> | | | | | | | | | | | |
| Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 7 days before harvest | 7 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Amistrobin | 04879 | Servem Ltd | Fungicide | | | | 1.0 l/ha | 2.0 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Apollo 50SC | 04610 | Adama Agricultural Solutions UK Ltd | Acaricide | | | | 0.4 l/ha | 0.4 l/ha | 1 per year | 9 days before harvest | 9 days |
| | | | | | <i>Clofentezine</i> | <i>500 g/l</i> | | | | | |
| Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 375 gms/ha | 375 g/ha | 1 / year | 21 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | | 0.4 kgs/ha | 1.2 kgs/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| Pyrimala | 05429 | Servem Ltd | Fungicide | | | | 2.0 l/ha | 4.0 l/ha | 2 | 7 days before harvest | 7 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| Scala | 03916 | BASF plc | Fungicide | | | | 2.0l/ha | 4.0 l/ha | 2 per crop | 7 days before harvest | 7 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | | | | 0.8 l/ha | 1.6 l/ha | - | 21 days before harvest | |
| | | | | | <i>Carfentrazone-ethyl</i> | <i>60 g/l</i> | | | | | |
| Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | | 200ml/ha in 1000L water | 400 ml/ha | 2 per year | 1 day before harvest | 1 day |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------------|---------------------|---------------|-------------------------------------|-----------------|--------------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Raspberries - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Amistrobin | 04879 | Servem Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 | 10 days before harvest | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Apollo 50SC | 04610 | Adama Agricultural Solutions UK Ltd | Acaricide | | | 0.4 l/ha | 0.4 l/ha | 1 per year | 9 days before harvest | 9 days |
| | | | | | <i>Clofentezine</i> | <i>500 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.25 l/ha | 0.75 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |
| | Masai | 03585 | BASF plc | Insecticide | | | 375 gms/ha | 375 g/ha | 1 / year | 21 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| | Pyrimala | 05429 | Servem Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | 2 | 7 days before harvest | 7 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Rana | 05283 | Servem Ltd | Insecticide | | | 0.25 l/ha | - | 0.75 l/ha | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 7 days before harvest | 7 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | | | 0.8 l/ha | 1.6 l/ha | | 21 days before harvest | 21 days |
| | | | | | <i>Carfentrazone-ethyl</i> | <i>60 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 14 days before harvest | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product</i> | | | | | <i>Max Ind.</i> | <i>Max. Total</i> | <i>Max No of</i> | <i>Latest Time</i> | |
|--------------------------------|----------------|----------------------|-----------------|------------------|----------------|-----------------------------|-------------------|------------------|----------------------|-------------|
| <i>Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Dose:</i> | <i>Dose:</i> | <i>Applic's:</i> | <i>of Applic:</i> | <i>PHI:</i> |
| <i>Raspberries - protected</i> | | | | | | | | | | |
| Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 ml/ha in 1000L water | 600 ml/ha | 3 per year | 1 day before harvest | 1 day |
| | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------|---------------------|-----------------------------|---------------------|-----------------|---|-----------------|------------------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Redcurrants - field</i> | | | | | | | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | | 0.25 l/ha | 0.75 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Chikara Weed Cont | 04203 | ISK Biosciences Europe N.V. | Herbicide | | | | 120 g/ha | 120 g/ha | 1 per year | Before bud burst | |
| | | | | | <i>Flazasulfuron</i> | <i>25 % w/w</i> | | | | | |
| Corbel | 90438 | BASF plc | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | 2 | 2 weeks before harvest | 2 weeks |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | | 3.0 l/ha | 3.0 l/ha | 1 | Before flowering or after harvest | |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| Manfil 75 WG | 03598 | Indofil Industries Limited | Fungicide | | | | 1.5 kg per 1000 litres water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75 % w/w</i> | | | | | |
| Manfil 80 WP | 03563 | Indofil Industries Limited | Fungicide | | | | 1.4 kg per 1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 1.5 kgs/ha | 1.5 kgs/ha | - | 7 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Penncozeb 80 WP | 03609 | UPL Europe Ltd | Fungicide | | | | 1.4 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Penncozeb WDG | 03608 | UPL Europe Ltd | Fungicide | | | | 1.5 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>750 g/kg</i> | | | | | |
| Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | | 0.4 kgs/ha | 1.2 kgs/ha | 3 | 28 days pre harvest | 28 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------------|---------------------|---------------|------------------------------|-----------------|--|-------------------|-------------------------|---|----------------------------|---|-------------|
| <i>Redcurrants - field</i> | | | | | | | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Signum | 03799 | BASF plc | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest) | - | 3 days before harvest, N/A for post harvest | |
| | | | | | <i>Boscalid</i> | <i>26.7 % w/w</i> | | | | | |
| | | | | | <i>Pyraclostrobin</i> | <i>6.7 % w/w</i> | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 ml/ha in 250L water | 400 ml/ha | 2 per year | 7 days before harvest | 7 days |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| <i>Redcurrants - protected</i> | | | | | | | | | | | |
| | AQ 10 | 05801 | CBC (Europe) S.r.l. | Fungicide | | | 70 g/ha | - | 12 | - | |
| | | | | | <i>Ampelomyces Quisqualis</i> | <i>58 %w/w</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------|---------------------|---------------|----------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Rhubarb</i> | | | | | | | | | | | |
| | Flexidor | 04963 | Dow AgroSciences Ltd | Herbicide | | | 0.4 l/ha | 0.4 l/ha | 1 per year | Dormant stage | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Isoxaben</i> | <i>500 g/l</i> | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 4.0 l/ha | 4.0 l/ha | 1 | Before crop emergence | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Pendimethalin</i> | <i>330 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| <i>Rocket</i> | | | | | | | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 | 3 days pre harvest | 3 days |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | <i>Acetamiprid</i> | <i>20 % w/w</i> | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Rosemary</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | <i>Cycloxydim</i> | <i>100 g/l</i> | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|---------------------------|-----------------|---|-------------------|------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Rubus Hybrid - field</i> | | | | | | | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Masai | 03585 | BASF plc | Insecticide | | | 375 g/ha | 375 g/ha | 1 per year | 21 days before harvest | 21 days |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | | | 0.8 l/ha | 1.6 l/ha | - | 21 days before harvest | |
| | | | | | <i>Carfentrazone-ethyl</i> | <i>60 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------------|---------------------|---------------|------------------------------|-----------------|---|-------------------|------------------------|-------------------------|----------------------------|--|-------------|
| <i>Rubus Hybrid - protected</i> | | | | | | | | | | | |
| | Folicur 250 EW | 00468 | Bayer CropScience Limited | Fungicide | | | 0.8 l/ha | 2.4 l/ha | 3 per year | 14 days before harvest | 14 days |
| | | | | | <i>Tebuconazole</i> | <i>250 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Masai | 03585 | BASF plc | Insecticide | | | 375 g/ha | 375 g/ha | 1 per year | 21 days before harvest | 21 days |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Spotlight Plus | 03294 | FMC Chemical s.p.r.l. | Herbicide | | | 0.8 l/ha | 1.6 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Carfentrazone-ethyl</i> | <i>60 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m ² * | 0.5 g/m ² | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------|---------------------|---------------|---------------------------|-----------------|---------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Rye - spring</i> | | | | | | | | | | | |
| | Compitox Plus | 02843 | Nufarm UK Limited | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 250 mls/ha | 750 mls/ha | | Before GS 85 | |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 per crop | Before GS 22 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Duplosan KV | 02842 | Nufarm UK Limited | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | Optica | 02661 | Nufarm UK Limited | Herbicide | | | 2.0 l/ha | 2.0 l/ha | - | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | Twinprop | 04427 | DHM Agrochemicals Ltd., | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| <i>Rye - winter</i> | | | | | | | | | | | |
| | Compitox Plus | 02843 | Nufarm UK Limited | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | Deca 2.5% EC | 04952 | Sharda Europe b.v.b.a | Insecticide | | | 250 mls/ha | 750 mls/ha | - | Before GS 85 | |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 250 mls/ha | 750 mls/ha | - | Before GS 85 | |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 per crop | Before GS 22 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Duplosan KV | 02842 | Nufarm UK Limited | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | GAT Decline 2.5 E | 04422 | FMC Chemical s.p.r.l. | Insecticide | | | 250 mls/ha | 750 mls/ha | - | Before GS 85 | |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Optica | 02661 | Nufarm UK Limited | Herbicide | | | 2.0 l/ha | 2.0 l/ha | - | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | Twinprop | 04427 | DHM Agrochemicals Ltd., | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-------------|---------------------|---------------|---------------------------|-----------------|--------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Sage</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|--------------------------------------|-----------------|-----------------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Salsify</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 300 mls/ha | 900 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 8.4 kg/ha | 4 per crop | 30 days before harvest | 30 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 8 weeks before harvest | 56 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Globazar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|---------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Salsify</i> | | | | | | | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | PDM 330 EC | 03582 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 | | |
| | | | | | <i>Pendimethalin</i> | <i>330 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before GS 11 | 80 days |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | | 14 days before harvest | 14 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | | | 3.3 l/ha | 3.3 l/ha | 1 | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>400 g/l</i> | | | | | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | | | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | | | | | <i>Pendimethalin</i> | <i>455 g/l</i> | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Cycloxydim</i> | <i>100 g/l</i> | | | | | |
| | Switch | 03761 | Syngenta UK Ltd | Fungicide | | | 1.0 kg/ha | 3.0 kg/ha | 3 | 7 days before harvest | |
| | | | | | <i>Cyprodinil</i> | <i>37.5 % w/w</i> | | | | | |
| | | | | | <i>Fludioxonil</i> | <i>25.0 % w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|---------------------------------|-----------------|---------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Salsify</i> | | | | | | | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | <i>Oxamyl</i> | <i>100 g/kg</i> | 1.0 l/ha | 3.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|----------------------------|-----------------|-----------------------------|-----------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Shallot</i> | | | | | | | | | | | |
| | Amistar Opti | 03408 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Bravo 500 | 03452 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Curator | 04504 | Syngenta UK Ltd | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>400 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>80 g/l</i> | | | | | |
| | Daconil | 05748 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 5.0 l/ha | 5.0 l/ha | - | Before 5th leaf visible | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 true leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Flexidor | 04963 | Dow AgroSciences Ltd | Herbicide | | | 0.2 l/ha | 0.2 l/ha | 1 per year | After transplanting or beginning of root formation | |
| | | | | | <i>Isoxaben</i> | <i>500 g/l</i> | | | | | |
| | Folio Gold | 02363 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 6.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Metalaxyl M</i> | <i>37.5 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | 1 | 4 weeks before harvest | 28 days |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | 7 days before harvest | 7 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Jupital | 04503 | Syngenta UK Ltd | Fungicide | | | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 225 mls/ha | 3 per crop | 21 days before harvest | 21 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Lentagran WP | 03082 | Belchim Crop Protection NV | Herbicide | | | 2.0 kg/ha | 2.0 kg/ha | 1 | Before 7 leaf stage | |
| | | | | | <i>Pyridate</i> | <i>45 % w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|---------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Shallot</i> | | | | | | | | | | | |
| | NEMGuard DE | 05368 | ECOSpray Ltd | Nematicide | | | 20 kg/ha | 20 kg/ha | 1 | At drilling | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | <i>Garlic Extract</i> | <i>450 g/kg</i> | 75 mls/ha | 225 mls/ha | 4 | 21 days before harvest | 21 days |
| | Pyramin DF | 04264 | BASF plc | Herbicide | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | 1.4 kg/ha | 1.4 kg/ha | - | Before GS 13 | |
| | Roxy 800 EC | 05739 | Globachem nv | Herbicide | <i>Chloridazon</i> | <i>65% w/w</i> | 5.0 l/ha | 5.0 l/ha | - | Before GS 15 | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Prosulfocarb</i> | <i>800 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Starane Hi-Load H | 04549 | Dow AgroSciences Ltd | Herbicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 0.3 l/ha | 0.3 l/ha | | Before GS 15 | |
| | Stomp 400 SC | 02879 | BASF plc | Herbicide | <i>Fluroxypyr</i> | <i>333 g/l</i> | 3.3 l/ha | 3.3 l/ha | 1 | Before crop emergence | |
| | Stomp Aqua | 03750 | BASF plc | Herbicide | <i>Pendimethalin</i> | <i>400 g/l</i> | 2.9 l/ha | 2.9 l/ha | 1 / crop | Before crop emergence | |
| | Valbon | 03411 | Certis Europe BV | Fungicide | <i>Pendimethalin</i> | <i>455 g/l</i> | 1.6 kg/ha | 9.6 kg/ha | 6 per crop | 28 days pre-harvest | 28 days |
| | Vertik | 04030 | Syngenta UK Ltd | Fungicide | <i>Mancozeb</i> | <i>700 g/kg</i> | 2.0 l/ha | 4.0 l/ha | - | 14 days before harvest | 14 days |
| | Wing P | 03590 | BASF plc | Herbicide | <i>Benthiavalicarb</i> | <i>17.5 g/kg</i> | 4.0 l/ha | 4.0 l/ha | - | Pre-emergence / Pre-transplanting | |
| | | | | | <i>Azoxystrobin</i> | <i>100 g/l</i> | | | | | |
| | | | | | <i>Chlorothalonil</i> | <i>500 g/l</i> | | | | | |
| | | | | | <i>Dimethenamid-P</i> | <i>212.5 g/l</i> | | | | | |
| | | | | | <i>Pendimethalin</i> | <i>250g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|---------------------------|-----------------|------------------------------------|-----------------|------------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Spinach</i> | | | | | | | | | | | |
| | Bonalan | 05763 | Gowan Crop Protection Ltd | Herbicide | | | 8.0 l/ha | 8.0 l/ha | - | Incorporation pre sowing/planting | |
| | | | | | <i>Benfluralin</i> | <i>180 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fenomenal | 04956 | Bayer CropScience Limited | Fungicide | | | 2.5 kg/ha | 5.0 kg/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>600 g/kg</i> | | | | | |
| | | | | | <i>Fenamidone</i> | <i>60 g/kg</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Infinito | 02532 | Bayer CropScience Limited | Fungicide | | | 1.6 l/ha | 1.6 l/ha | 1 per crop | Before GS 49 | 14 days |
| | | | | | <i>Fluopicolide</i> | <i>62.5 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>625 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days pre harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST</i> | <i>2% w/w</i> | | | | | |
| | | | | | <i>BIPESCO 5/F52</i> | | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------|---------------------|---------------|---------------------------------|-----------------|--------------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Spinach</i> | | | | | | | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 2.5 l/ha | 5.0 l/ha | - | 21 days before harvest | 21 days |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 7 days before harvest | 7 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 10 g/m3 | 10 g/m3 | 1 | Pre planting / sowing | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 mls/ha | 600 mls/ha | 3 | | 3 days |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| | Venzar 500 SC | 04805 | DuPont (UK) Ltd | Herbicide | | | 0.8 l/ha | 0.8 l/ha | - | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>500 g/l</i> | | | | | |
| | Venzar Flowable | 91516 | DuPont (UK) Ltd | Herbicide | | | 5 l/ha | 5 l/ha | 1 | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>440 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------|---------------------|---------------|---------------------------------|-----------------|--|-----------------|------------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Spinach beet</i> | | | | | | | | | | | |
| | Bonalan | 05763 | Gowan Crop Protection Ltd | Herbicide | | | 8.0 l/ha | 8.0 l/ha | - | Incorporation pre sowing/planting | |
| | | | | | <i>Benfluralin</i> | <i>180 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Fenomenal | 04956 | Bayer CropScience Limited | Fungicide | | | 2.5 kg/ha | 5.0 kg/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>600 g/kg</i> | | | | | |
| | | | | | <i>Fenamidone</i> | <i>60 g/kg</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Met 52 Granular | 04811 | Novozymes France S.A.S. | Insecticide | | | 0.5 kgs/m ³ | - | - | Pre-planting | |
| | | | | | <i>Metarhizium anisoplice - ST BIPESCO 5/F52</i> | <i>2% w/w</i> | | | | | |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | | | 200 mls/ha | 600 mls/ha | 3 | | 3 days |
| | | | | | <i>Spinosad</i> | <i>480 g/l</i> | | | | | |
| | Venzar 500 SC | 04805 | DuPont (UK) Ltd | Herbicide | | | 0.8 l/ha | 0.8 l/ha | - | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>500 g/l</i> | | | | | |
| | Venzar Flowable | 91516 | DuPont (UK) Ltd | Herbicide | | | 5 l/ha | 5 l/ha | 1 | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>440 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------|---------------------|---------------|------------------------------|-----------------|--------------------------------|----------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Squash - protected</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Previcur Energy | 04041 | Bayer CropScience Limited | Fungicide | | | 3 mls per sq m | 6 mls per sq m | - | Before transplanting | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Fosetyl (-aluminium)</i> | <i>310 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | | | | | <i>Propamocarb</i> | <i>530 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | | | | | <i>Trichoderma</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Asperellum (strain T34)</i> | | | | | | |
| | Vivando | 05685 | BASF plc | Fungicide | | | 0.2 l/ha | 0.4 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Metrafenone</i> | <i>500 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|-------------------------------------|---------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Strawberries - field</i> | | | | | | | | | | | |
| Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 | 3 days pre-harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Amistrobin | 04879 | Servem Ltd | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 | 3 days before harvest | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Apollo 50SC | 04610 | Adama Agricultural Solutions UK Ltd | Acaricide | | | | 0.4 l/ha | 0.4 l/ha | 1 per year | 9 days before harvest | 9 days |
| | | | | | <i>Clofentezine</i> | <i>500 g/l</i> | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | | 0.25 l/ha | 0.50 l/ha | 2 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Chess WG | 03365 | Syngenta UK Ltd | Insecticide | | | | 0.4 kg/ha | 1.2 kg/ha | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | | 0.7 l/ha | 1.3 l/ha | - | 30 days before harvest | 30 days |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| Fortress | 02364 | Dow AgroSciences Ltd | Fungicide | | | | 0.25 l/ha | 0.5 l/ha | 2 | 14 days before harvest | 14 days |
| | | | | | <i>Quinoxifen</i> | <i>500 g/l</i> | | | | | |
| Globaztar SC | 04859 | Globachem nv | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | | 125 ml/ha | 250 ml/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Life Scientific Azox | 04405 | Life Scientific Limited | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 375 gms/ha | 375 g/ha | 1 / year | 21 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Nimrod | 90789 | Adama Agricultural Solutions UK Ltd | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Bupirimate</i> | <i>250 g/l</i> | | | | | |
| Rana | 05283 | Servem Ltd | Insecticide | | | | 0.25 l/ha | - | 0.5 l/ha | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------------|---------------------|-------------------------------------|----------------------|-----------------|---------------------------|-----------------|---|---|----------------------------|-------------------------------|-------------|
| <i>Strawberries - protected</i> | | | | | | | | | | | |
| Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 | 3 days pre-harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Amistrobin | 04879 | Servem Ltd | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 | 3 days before harvest | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Apollo 50SC | 04610 | Adama Agricultural Solutions UK Ltd | Acaricide | | | | 0.4 l/ha | 0.4 l/ha | 1 per year | 9 days before harvest | 9 days |
| | | | | | <i>Clofentezine</i> | <i>500 g/l</i> | | | | | |
| Borneo | 04235 | Sumitomo Chemical Agro Europe SAS | Acaricide | | | | 500mls/ha - max conc. 35 mls/100 litres water | 500mls/ha - max conc. 35 mls/100 litres water | 1/crop | 3 days before harvest | 3 days |
| | | | | | <i>Etoxazole</i> | <i>110 g/l</i> | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | | 0.25 l/ha | 0.50 l/ha | 2 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Chess WG | 03365 | Syngenta UK Ltd | Insecticide | | | | 0.4 kg/ha | 1.2 kg/ha | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| Dynamec | 02640 | Syngenta UK Ltd | Acaricide/Insecticid | | | | 50 mls/product/100 litres water | - | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Abamectin</i> | <i>18 g/l</i> | | | | | |
| Fortress | 02364 | Dow AgroSciences Ltd | Fungicide | | | | 0.25 l/ha | 0.5 l/ha | 2 | 14 days before harvest | 14 days |
| | | | | | <i>Quinoxifen</i> | <i>500 g/l</i> | | | | | |
| Globazar SC | 04859 | Globachem nv | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | | 75 ml/ha | 150 ml/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Life Scientific Azox | 04405 | Life Scientific Limited | Fungicide | | | | 1.0 l/ha | 4.0 l/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 375 gms/ha | 375 g/ha | 1 / year | 21 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | | | | | | |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------------------------|---------------------|---------------|-------------------------------------|----------------------|--|----------------|-----------------------|-------------------------|----------------------------|--|-------------|
| <i>Strawberries - protected</i> | | | | | | | | | | | |
| | Nimrod | 90789 | Adama Agricultural Solutions UK Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Bupirimate</i> | <i>250 g/l</i> | | | | | |
| | Rana | 05283 | Servem Ltd | Insecticide | | | 0.25 l/ha | - | 0.5 l/ha | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Sabik-1.8% | 04907 | Servem Ltd | Acaricide/Insecticid | | | 50 mls / 100 l water | 4.0 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Abamectin</i> | <i>18 g/l</i> | | | | | |
| | Sequoia | 05915 | Dow AgroSciences Ltd | Insecticide | | | 200 mls/ha | 400 mls/ha | 2 | 1 day before harvest | 1 day |
| | | | | | <i>Sulfoxaflor</i> | <i>120 g/l</i> | | | | | |
| | T34 Biocontrol | 05620 | Biocontrol Technologies S.L. | Fungicide | | | 0.5 g/m2* | 0.5 g/m2 | 1 | Immediately after sowing seeds or rooting cuttings | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | | | | | <i>Trichoderma Asperellum (strain T34)</i> | <i>12% w/w</i> | | | | | |
| | Talius | 04604 | DuPont (UK) Ltd | Fungicide | | | 0.19 l/ha | 0.19 l/ha | 1 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Proquinazid</i> | <i>200 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|--|-----------------|---------------------------|------------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Swedes</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 10 days before harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Benevia 10 OD | 05354 | DuPont (UK) Ltd | Insecticide | | | 0.75 l/ha | 1.5 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Cyantraniliprole</i> | <i>100 g/l</i> | | | | | |
| | Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| | Bonalan | 05763 | Gowan Crop Protection Ltd | Herbicide | | | 8.0 l/ha | 8.0 l/ha | - | Incorporation pre sowing/planting | |
| | | | | | <i>Benfluralin</i> | <i>180 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Curenox 50 WP | 05829 | Industrias Quimicas Del Valles S.A (IQV, S.A.) | Fungicide | | | 1 kg/ha | 6 kg/ha | - | 8 days before harvest | 8 days |
| | | | | | <i>Copper oxychloride</i> | <i>87.8% w/w</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fortress | 02364 | Dow AgroSciences Ltd | Fungicide | | | 0.3 l/ha | 0.4 l/ha | 2 | 28 days pre harvest | 28 days |
| | | | | | <i>Quinoxifen</i> | <i>500 g/l</i> | | | | | |
| | Gamit 36 CS | 05357 | FMC Chemical s.p.r.l. | Herbicide | | | 0.25 l/ha | 0.25 l/ha | - | pre-emergence | |
| | | | | | <i>Clomazone</i> | <i>360 g/l</i> | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|---------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Swedes</i> | | | | | | | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | | N/A | |
| | | | | | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Movento | 04101 | Bayer CropScience Limited | Insecticide | | | 0.3 l/ha | 0.6 l/ha | 2 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Spirotetramat</i> | <i>150 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 0.5 l/ha | | 28 days before harvest | 28 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Venzar 500 SC | 04805 | DuPont (UK) Ltd | Herbicide | | | 0.4 l/ha | 0.4 l/ha | - | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>500 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------|---------------------|---------------|---------------------------------|-----------------|---|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Swedes</i> | | | | | | | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | <i>Oxamyl</i> | <i>100 g/kg</i> | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| <i>Sweetcorn</i> | | | | | | | | | | | |
| | Lepinox Plus | 05254 | CBC (Europe) S.r.l. | Insecticide | | | 1.0 kg/ha | 3.0 kg/ha | - | N/A | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus thuringiensis subspecies kurstaki EG 2348</i> | <i>375 g/kg</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| <i>Thyme</i> | | | | | | | | | | | |
| | Stratos Ultra | 04936 | BASF plc | Herbicide | | | 4.0 l/ha | 4.0 l/ha | 1 per crop | 4 weeks before harvest | 4 weeks |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | <i>Cycloxydim</i> | <i>100 g/l</i> | 1.5 kgs/ha | 3.0 kgs/ha | | 3 days before harvest | 3 days |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|--------------------------------------|------------------|----------------------------|------------------|---------------------------------------|---------------------------------------|----------------------------|-------------------------------|-------------|
| <i>Tomatoes - protected</i> | | | | | | | | | | | |
| | Acrobat MZ | 04059 | BASF plc | Fungicide | | | 2.4 kg/ha | 7.2 kg/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Mancozeb</i> | <i>667 g/kg</i> | | | | | |
| | | | | | <i>Dimethomorph</i> | <i>75 g/kg</i> | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 100 mls / 100 litres water | - | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | 0.45 l/ha | 0.9 l/ha | 2/crop | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| | Cerone | 03933 | Bayer CropScience Limited | Growth Regulator | | | 100 ml in 100 litre water. To run off | 100 ml in 100 litre water. To run off | 1 per crop | 5 days before harvest | 5 days |
| | | | | | <i>Ethephon</i> | <i>480 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Coragen | 05295 | DuPont (UK) Ltd | Insecticide | | | 250 ml/ha | 500 ml/ha | - | 1 day before harvest | 1 day |
| | | | | | <i>Chlorantraniliprole</i> | <i>200 g/l</i> | | | | | |
| | Curzate M WG | 04723 | DuPont (UK) Ltd | Fungicide | | | 2.3 kg/ha | 6.9 kg/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Cymoxanil</i> | <i>4.5 % w/w</i> | | | | | |
| | | | | | <i>Mancozeb</i> | <i>68 % w/w</i> | | | | | |
| | Dithane 945 | 03705 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.0 kg/ha | 6.0 kg/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Mancozeb</i> | <i>80 % w/w</i> | | | | | |
| | Dithane Dry Flowa | 03707 | Indofil Industries (Netherlands) B.V | Fungicide | | | 2.1 kg/ha | 6.3 kg/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Mancozeb</i> | <i>75% w/w</i> | | | | | |
| | Electis 75 WG | 01821 | Gowan Crop Protection Ltd | Fungicide | | | 1.8 kg/ha | 5.4 kg/ha | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Zoxamide</i> | <i>8.3 %w/w</i> | | | | | |
| | | | | | <i>Mancozeb</i> | <i>66.7 %w/w</i> | | | | | |
| | Globaztar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|----------------------|---------------|-------------------------------------|-----------------|--------------------------|-------------------|---------------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Tomatoes - protected</i> | | | | | | | | | | | |
| | Life Scientific Azox | 04405 | Life Scientific Limited | Fungicide | | | 100 ml per 100L water | | 3 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Manfil 75 WG | 03598 | Indofil Industries Limited | Fungicide | | | 2.1 kg/ha | 6.3 kg/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Mancozeb</i> | <i>75 % w/w</i> | | | | | |
| | Manfil 80 WP | 03563 | Indofil Industries Limited | Fungicide | | | 2.0 kg/ha | 4.0 kg/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| | Microthiol Special | 90548 | UPL Europe Ltd | Fungicide | | | 0.01kgs/5 litres water | - | - | - | 0 days |
| | | | | | <i>Sulphur</i> | <i>80 %w/w</i> | | | | | |
| | Nimrod | 90789 | Adama Agricultural Solutions UK Ltd | Fungicide | | | 0.2L per 100L water | - | 6 per crop | 2 days before harvest | 2 days |
| | | | | | <i>Bupirimate</i> | <i>250 g/l</i> | | | | | |
| | Option | 04722 | DuPont (UK) Ltd | Fungicide | | | 0.4 kg/ha | 1.6 kg/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Cymoxanil</i> | <i>600 g/kg</i> | | | | | |
| | Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | 0.06 kgs/100 litres water | - | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| | Revus | 02789 | Syngenta UK Ltd | Fungicide | | | 0.6 l/ha | 2.4 l/ha | 4 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Mandipropamid</i> | <i>250 g/l</i> | | | | | |
| | Ridomil Gold MZ 6 | 01516 | Syngenta UK Ltd | Fungicide | | | 2.5 kg/ha | 7.5 kg/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Mancozeb</i> | <i>64 %w/w</i> | | | | | |
| | | | | | <i>Metalaxyl M</i> | <i>4.0 %w/w</i> | | | | | |
| | Scala | 03916 | BASF plc | Fungicide | | | 0.5 l/ha | 1.0 l/ha | 2 per crop | 3 days before harvest | 3 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | SL567A | 03106 | Syngenta UK Ltd | Fungicide | | | 0.3 l/ha | 0.9 l/ha | 3 per crop | | 3 days |
| | | | | | <i>Metalaxyl M</i> | <i>465.2 g/l</i> | | | | | |
| | Teldor | 02003 | Bayer CropScience Limited | Fungicide | | | 1.5 kgs/ha | 4.5 kgs/ha | 3 / crop | 1 day before harvest | 1 day |
| | | | | | <i>Fenhexamid</i> | <i>50 %w/w</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|-----------------------------|---------------------|---------------|---------------------------------|-----------------|---------------------|----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Tomatoes - protected</i> | | | | | | | | | | | |
| | Vivando | 05685 | BASF plc | Fungicide | | | 0.45 l/ha | 0.9 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Metrafenone</i> | <i>500 g/l</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 3.0 l/ha | - | 3 days before harvest | 3 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------|---------------------|---------------|-------------------------------------|-----------------|------------------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Triticale</i> | | | | | | | | | | | |
| | Barclay Bolt XL | 04102 | Barclay Chemicals Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | 2 per crop | Before GS 71 | |
| | | | | | <i>Propiconazole</i> | <i>250 g/l</i> | | | | | |
| | Barclay Propizole | 04099 | Barclay Chemicals Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | 2 per crop | Before GS 71 | |
| | | | | | <i>Propiconazole</i> | <i>250 g/l</i> | | | | | |
| | Bumper | 03292 | Adama Agricultural Solutions UK Ltd | Fungicide | | | 0.5 l/ha | 1.0 l/ha | 2 per crop | Before GS 71 | 35 days |
| | | | | | <i>Propiconazole</i> | <i>250 g/l</i> | | | | | |
| | Cliophar 600 SL | 04864 | Arysta LifeScience Benelux SPRL | Herbicide | | | 0.12 l/ha | 0.12 l/ha | - | Before GS 33 | |
| | | | | | <i>Clopyralid</i> | <i>600 g/l</i> | | | | | |
| | Compitox Plus | 02843 | Nufarm UK Limited | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | Deca 2.5% EC | 04952 | Sharda Europe b.v.b.a | Insecticide | | | 250 mls/ha | 750 mls/ha | - | Before GS 85 | |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Decis | 02693 | Bayer CropScience Limited | Insecticide | | | 250 mls/ha | 750 mls/ha | | before GS 85 | |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Defy | 04277 | Syngenta UK Ltd | Herbicide | | | 3.0 l/ha | 3.0 l/ha | - | Before GS22 | |
| | | | | | <i>Prosulfocarb</i> | <i>800 g/l</i> | | | | | |
| | Duplosan KV | 02842 | Nufarm UK Limited | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| | Empire Supreme | 05061 | Croplink (Ireland) Ltd | Herbicide | | | 100 g/ha | 100 g/ha | 1 per crop | Before GS 39 | |
| | | | | | <i>Thifensulfuron methyl</i> | <i>400 g/kg</i> | | | | | |
| | | | | | <i>Metsulfuron (-methyl)</i> | <i>40 g/kg</i> | | | | | |
| | Finish SX | 02329 | DuPont (UK) Ltd | Herbicide | | | 75 g/ha | 75 g/ha | 1 | Before GS 39 | |
| | | | | | <i>Metsulfuron (-methyl)</i> | <i>67 g/kg</i> | | | | | |
| | | | | | <i>Thifensulfuron methyl</i> | <i>333 g/kg</i> | | | | | |
| | GAT Decline 2.5 E | 04422 | FMC Chemical s.p.r.l. | Insecticide | | | 250 mls/ha | 750 mls/ha | - | Before GS 85 | |
| | | | | | <i>Deltamethrin</i> | <i>25 g/l</i> | | | | | |
| | Harmony M SX | 02330 | DuPont (UK) Ltd | Herbicide | | | 100 g/ha | 100 g/ha | 1 | Before GS 39 | |
| | | | | | <i>Thifensulfuron methyl</i> | <i>400 g/kg</i> | | | | | |
| | | | | | <i>Metsulfuron (-methyl)</i> | <i>40 g/kg</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|--------------------------|---------------|-------------------------|-----------------|------------------------------|-----------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Triticale</i> | | | | | | | | | | |
| Optica | 02661 | Nufarm UK Limited | Herbicide | | | 2.0 l/ha | 2.0 l/ha | - | Before GS 33 | |
| | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |
| Presite SX | 02888 | DuPont (UK) Ltd | Herbicide | | | 75 g/ha | 75 g/ha | 1 | Before GS 39 | |
| | | | | <i>Thifensulfuron methyl</i> | <i>333 g/kg</i> | | | | | |
| | | | | <i>Metsulfuron (-methyl)</i> | <i>67 g/kg</i> | | | | | |
| Propi 25 EC | 04205 | Sharda Europe b.v.b.a | Fungicide | | | 0.5 l/ha | 1.0 l/ha | 2 per crop | Before GS 71 | |
| | | | | <i>Propiconazole</i> | <i>250 g/l</i> | | | | | |
| Twinprop | 04427 | DHM Agrochemicals Ltd., | Herbicide | | | 2.3 l/ha | 2.3 l/ha | 1 per crop | Before GS 33 | |
| | | | | <i>Mecoprop-P</i> | <i>600 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|--|-----------------|---------------------------|------------------|-----------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Turnip</i> | | | | | | | | | | | |
| | Amistar | 01351 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 4.0 l/ha | 4 | 10 days pre harvest | 10 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Benevia 10 OD | 05354 | DuPont (UK) Ltd | Insecticide | | | 0.75 l/ha | 1.5 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Cyantraniliprole</i> | <i>100 g/l</i> | | | | | |
| | Biscaya | 03602 | Bayer CropScience Limited | Insecticide | | | 0.4 l/ha | 0.8 l/ha | 2 | 7 days before harvest | |
| | | | | | <i>Thiacloprid</i> | <i>240 g/l</i> | | | | | |
| | Bonalan | 05763 | Gowan Crop Protection Ltd | Herbicide | | | 8.0 l/ha | 8.0 l/ha | - | Incorporation pre sowing/planting | |
| | | | | | <i>Benfluralin</i> | <i>180 g/l</i> | | | | | |
| | Chamane | 05002 | UPL Europe Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Corbel | 90438 | BASF plc | Fungicide | | | 1.0 l/ha | 3.0 l/ha | 3 | 28 days before harvest | 28 days |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| | Curenox 50 WP | 05829 | Industrias Quimicas Del Valles S.A (IQV, S.A.) | Fungicide | | | 1 kg/ha | 6 kg/ha | - | 8 days before harvest | 8 days |
| | | | | | <i>Copper oxychloride</i> | <i>87.8% w/w</i> | | | | | |
| | Dual Gold | 04192 | Syngenta UK Ltd | Herbicide | | | 1.4 l/ha | 1.4 l/ha | - | Up to 4 leaf stage | |
| | | | | | <i>S-Metolachlor</i> | <i>960 g/l</i> | | | | | |
| | Eminentos 10 CS | 04933 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Fortress | 02364 | Dow AgroSciences Ltd | Fungicide | | | 0.3 l/ha | 0.4 l/ha | 2 | 28 days pre harvest | 28 days |
| | | | | | <i>Quinoxifen</i> | <i>500 g/l</i> | | | | | |
| | Gazelle | 03248 | Nisso Chemical Europe GmbH | Insecticide | | | 250 g/ha | 500 g/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Acetamiprid</i> | <i>20 % w/w</i> | | | | | |
| | Globazar SC | 05529 | Globachem nv | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |
| | Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Karis 10 CS | 04454 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|---------------|---------------------|---------------|---------------------------------|-----------------|---------------------------|-------------------|-----------------------|-------------------------|----------------------------|-------------------------------|-------------|
| <i>Turnip</i> | | | | | | | | | | | |
| | Laidir 10 CS | 04932 | FMC Chemical s.p.r.l. | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Life Scientific Lam | 04868 | Life Scientific Limited | Insecticide | | | 75 mls/ha | 150 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Markate 50 | 05001 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Movento | 04101 | Bayer CropScience Limited | Insecticide | | | 0.3 l/ha | 0.6 l/ha | 2 per crop | 8 weeks before harvest | 8 weeks |
| | | | | | <i>Spirotetramat</i> | <i>150 g/l</i> | | | | | |
| | Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | 150 mls/ha | 450 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Pilot Ultra | 04955 | Nissan Chemical Europe | Herbicide | | | 2.5 l/ha | 2.5 l/ha | - | 60 days before harvest | 60 days |
| | | | | | <i>Quizalofop-P-ethyl</i> | <i>50 g/l</i> | | | | | |
| | Ravane | 05536 | Globachem nv | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 per crop | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>50 g/l</i> | | | | | |
| | Reflect | 05462 | Syngenta UK Ltd | Fungicide | | | 1.0 l/ha | 2.0 l/ha | 2 per crop | 14 days before harvest | 14 days |
| | | | | | <i>Isopyrazam</i> | <i>125 g/l</i> | | | | | |
| | Score 250 EC | 04566 | Syngenta UK Ltd | Fungicide | | | 0.5 l/ha | 0.5 l/ha | | 28 days before harvest | 28 days |
| | | | | | <i>Difenoconazole</i> | <i>250 g/l</i> | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | | | | | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | | | | | |
| | Sparviero | 04469 | Oxon Italia SpA | Insecticide | | | 75 ml/ha | 150 ml/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| | Venzar 500 SC | 04805 | DuPont (UK) Ltd | Herbicide | | | 0.4 l/ha | 0.4 l/ha | - | Pre-emergence | |
| | | | | | <i>Lenacil</i> | <i>500 g/l</i> | | | | | |
| | Vydate 10G | 03698 | DuPont (UK) Ltd | Nematicide | | | 0.09 kg/100 m row | 0.09 kg/100 m row | 1 per crop | At drilling | |
| | | | | | <i>Oxamyl</i> | <i>100 g/kg</i> | | | | | |
| | Zoxis | 05348 | Arysta LifeScience Benelux SPRL | Fungicide | | | 1.0 l/ha | 2.0 l/ha | - | 14 days before harvest | 14 days |
| | | | | | <i>Azoxystrobin</i> | <i>250 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|------------------------------|---------------------|-----------------------------|---------------------|-----------------|-----------------------------|-----------------|------------------------------|-------------------------|----------------------------|-----------------------------------|-------------|
| <i>Whitecurrants - field</i> | | | | | | | | | | | |
| CaLypso | 02441 | Bayer CropScience Limited | Insecticide | | | | 0.25 l/ha | 0.75 l/ha | 3 | 3 days before harvest | 3 days |
| | | | | | <i>Thiacloprid</i> | <i>480 g/l</i> | | | | | |
| Chikara Weed Cont | 04203 | ISK Biosciences Europe N.V. | Herbicide | | | | 120 g/ha | 120 g/ha | 1 per year | Before bud burst | |
| | | | | | <i>Flazasulfuron</i> | <i>25 % w/w</i> | | | | | |
| Corbel | 90438 | BASF plc | Fungicide | | | | 1.0 l/ha | 3.0 l/ha | 3 | 2 weeks before harvest | 2 weeks |
| | | | | | <i>Fenpropimorph</i> | <i>750 g/l</i> | | | | | |
| Fusilade Max | 01472 | Syngenta UK Ltd | Herbicide | | | | 3.0 l/ha | 3.0 l/ha | 1 | Before flowering or after harvest | |
| | | | | | <i>Fluazifop-P (-butyl)</i> | <i>125 g/l</i> | | | | | |
| Karate Zeon | 04084 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days pre harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Manfil 75 WG | 03598 | Indofil Industries Limited | Fungicide | | | | 1.5 kg per 1000 litres water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>75 % w/w</i> | | | | | |
| Manfil 80 WP | 03563 | Indofil Industries Limited | Fungicide | | | | 1.4 kg per 1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Masai | 03585 | BASF plc | Insecticide | | | | 1.5 kgs/ha | 1.5 kgs/ha | - | 7 days before harvest | |
| | | | | | <i>Tebufenpyrad</i> | <i>200 g/kg</i> | | | | | |
| Ninja 10CS | 05178 | Syngenta UK Ltd | Insecticide | | | | 100 mls/ha | 200 mls/ha | 4 | 14 days before harvest | 14 days |
| | | | | | <i>lambda-Cyhalothrin</i> | <i>100 g/l</i> | | | | | |
| Penncozeb 80 WP | 03609 | UPL Europe Ltd | Fungicide | | | | 1.4 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>800 g/kg</i> | | | | | |
| Penncozeb WDG | 03608 | UPL Europe Ltd | Fungicide | | | | 1.5 kg/1000L water | - | - | 28 days before harvest | 28 days |
| | | | | | <i>Mancozeb</i> | <i>750 g/kg</i> | | | | | |
| Plenum | 02238 | Syngenta UK Ltd | Insecticide | | | | 0.4 kgs/ha | 1.2 kgs/ha | 3 | 28 days pre harvest | 28 days |
| | | | | | <i>Pymetrozine</i> | <i>50 %w/w</i> | | | | | |
| Scala | 03916 | BASF plc | Fungicide | | | | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | | | | | <i>Pyrimethanil</i> | <i>400 g/l</i> | | | | | |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.

| <i>Crop</i> | <i>Product Name</i> | <i>PCS No</i> | <i>Company Name</i> | <i>Function</i> | <i>Substance</i> | <i>Content</i> | <i>Max Ind. Dose:</i> | <i>Max. Total Dose:</i> | <i>Max No of Applic's:</i> | <i>Latest Time of Applic:</i> | <i>PHI:</i> |
|----------------------------------|---------------------|---------------|---------------------------|-----------------|---|---------------------------------------|-------------------------|---|----------------------------|---|-------------|
| <i>Whitecurrants - field</i> | | | | | | | | | | | |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | | | 10 l/ha | 60 l/ha | 6 | | 0 days |
| | Signum | 03799 | BASF plc | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 1.5 kgs/ha | 4.5 kgs/ha (3 kgs pre , 1.5 kgs post harvest) | - | 3 days before harvest, N/A for post harvest | |
| | Stroby WG | 05092 | BASF plc | Fungicide | <i>Pyraclostrobin</i> <i>Boscalid</i> | <i>6.7 % w/w</i> <i>26.7 % w/w</i> | 0.2 kg/ha | 0.6 kg/ha | - | 14 days before harvest | 14 days |
| | Tracer | 02649 | Dow AgroSciences Ltd | Insecticide | <i>Kresoxim Methyl</i> <i>Spinosad</i> | <i>50% w/w</i> <i>480 g/l</i> | 200 ml/ha in 250L water | 400 ml/ha | 2 per year | 7 days before harvest | 7 days |
| <i>Whitecurrants - protected</i> | | | | | | | | | | | |
| | AQ 10 | 05801 | CBC (Europe) S.r.l. | Fungicide | <i>Ampelomyces Quisqualis</i> | <i>58 %w/w</i> | 70 g/ha | - | 12 | - | |
| | Scala | 03916 | BASF plc | Fungicide | <i>Pyrimethanil</i> | <i>400 g/l</i> | 2.0 l/ha | 4.0 l/ha | 2 per crop | 21 days before harvest | 21 days |
| | Serenade ASO | 03847 | Bayer CropScience Limited | Fungicide | <i>Bacillus subtilis</i> | <i>1015.1 g/l</i> | 10 l/ha | 60 l/ha | 6 | | 0 days |

Warning Off-Label approvals:

The use of any product in accordance with an off-label approval is entirely at the risk of the user. The directions for use on the instruction sheet must be followed – it is an offence to use products in a manner that has not been approved.