

750ml

approx. measure

500ml

250ml



Tern® 750EC

GROUP 5 FUNGICIDE

TERN® 750EC is an emulsifiable concentrate containing 750 g/l (81.8 %w/w) fenpropidin.

TERN® 750EC provides moderate control of powdery mildew on winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

PLEASE SEE ACCOMPANYING LEAFLET FOR PRODUCT USE DETAILS.

Authorisation Holder	Marketing Company
Syngenta UK Limited Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY Tel +44 (0) 1223 883400	Syngenta Ireland Limited Block 6, Cleaboy Business Park, Old Kilmeaden Road, Waterford, Ireland Tel: (051) 377203

**IN CASE OF TOXIC OR TRANSPORT EMERGENCY RING
+44 (0) 1484 538444 ANYTIME (24HR).**

**PROTECT FROM FROST
SHAKE WELL BEFORE USE**

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

1 Litre

© Syngenta AG, 2025



FOR PROFESSIONAL USE ONLY

To avoid risks to human health and the environment comply with the instructions for use.
TERN® 750EC is an emulsifiable concentrate containing 750 g/l (81.8% w/w) fenpropidin.

Warning

Harmful if swallowed or if inhaled.

Causes serious eye irritation.

May cause respiratory irritation.

May cause damage to organs (Central nervous system) through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects.

Do not breathe the mist or vapours.

Wash skin thoroughly after handling.

Wear eye protection/face protection.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

Call a POISON CENTRE/doctor if you feel unwell.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty triple rinsed clean containers which can be disposed of as non-hazardous waste.

Contains fenpropidin. May produce an allergic reaction.



PCS No. 04305

UFI: PWM0-ROTK-WOON-HF1K

CONDITIONS FOR USE

FOR USE AS AN AGRICULTURAL FUNGICIDE

CROP	Max. single dose (litres/hectare/crop)	Max. no of applications	Max. total dose (litres/hectare/crop)	Latest time of application
Winter wheat, spring wheat, durum wheat, winter barley, spring barley, oats, rye and triticale.	0.5	-	1.0	Before beginning of flowering: first anthers visible (GS 61).

Additional Safety Information

Operator Protection

Wash splashes from skin and eyes immediately.

Wash hands and exposed skin before meals and after work.

Harmful to Livestock. Keep livestock out of treated areas for at least 7 day.

Wear suitable protective clothing, gloves and eye/face protection.

In case of contact with eyes, rinse immediately with plenty of water and seek medical advice.

Do not breathe spray.

Storage and disposal

Keep in original container, tightly closed in a safe place.

Wash out containers thoroughly, preferably using an integrated pressure rinsing device, or manually rinse three times. Add washings to the sprayer at the time of filling. Complete filling to the required volume and continue to agitate throughout the spraying operation.

Restrictions

Do not apply more than 3 times to any one crop.

To protect aquatic organisms, respect an unsprayed buffer zone of 15m to surface water bodies.

TERN 750EC should only be used up to full the ear emergence (GS 59) stage on all crops.

Disease control may be reduced if strains of the pathogen less sensitive to TERN 750EC develop.

L1132307 REL/01D PPE 4225440

L1132307 REL/01D PPE 4225440

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

CROPS

TERN 750EC can be used on all varieties of winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.

DISEASE CONTROL

For moderate control of powdery mildew apply during the early stages of disease development. Best results will be achieved when not more than 5% of the third leaf from the top of the plant is infected. A repeat spray may be necessary during prolonged or severe attacks.

APPLICATION RATE

Apply TERN 750EC at 0.5 litres per hectare.

MIXING AND SPRAYING

Make sure the sprayer is set to give an even application at the correct volume.

Fill the sprayer tank with half the required quantity of clean water and begin agitation. Add the required quantity of TERN 750EC and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying.

Thoroughly wash all spray equipment with water immediately after use.

APPLICATION

Volume

Apply TERN 750EC in a recommended of 200 litres of water per hectare.

COMPATIBILITY

The following products may be tank-mixed with TERN 750EC provided the timing is correct for both TERN 750EC and the partner in the mixture.

The products should be added separately to the bulk of the water in the spray tank. Continuous agitation should be maintained and the products used immediately after mixing.

Herbicides

Approved salt formulations of:

MCPA

Mecoprop

Growth Regulators

Approved formulations containing chlormequat or chlormequat.

For further information on compatibility, contact Syngenta Ireland Ltd.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier