

Fungicide

INFINITO®

Infinito is a protectant/systemic fungicide used to control late blight in potatoes.

**A suspension concentrate containing
62.5 g/L (5.53 % w/w) fluopicolide +
625 g/L (55.3%w/w) propamocarb hydrochloride.**

For Professional use only.

Authorisation holder:
Bayer CropScience Ltd.
PO Box 1582
Cambridge
CB1 0FE
United Kingdom

Marketing company:
Bayer CropScience Ltd
Bayer Ltd
1st Floor, The Grange Offices
The Grange, Brewery Road,
Stillorgan, Co. Dublin
A94 H2K7

Freephone: 1800 818534

**For 24 hour emergency information
contact Bayer CropScience Limited
+44 330 678 3382 (24 hr)**
(charged as a standard international call to the UK)

**National Poisons Information Centre
Dublin: 01 809 2166**

10 L e



Safety information

INFINITO

UFI: 5UQ0-T0CN-300N-SQ2J

Contains 62.5 g/L (5.53 % w/w)
fluopicolide and 625 g/L (55.3 % w/w)
propamocarb hydrochloride.



Danger

**May cause an allergic skin reaction.
Suspected of damaging the unborn
child.**

**Very toxic to aquatic life with long
lasting effects.**

**Can cause long-lasting and diffuse
contamination of water resources.**

Obtain special instructions before
use.

Wear protective gloves/protective
clothing/eye protection/face
protection.

IF exposed or concerned: Call a
POISON CENTER/ doctor/ physician.
Dispose of contents/container to a
licensed hazardous waste disposal
contractor or collection site except
for triple rinsed empty containers
which can be disposed of as non-
hazardous waste.

**To avoid risks to human health and
the environment, comply with the
instructions for use.**

PCS No. 04309

IE85333615d rA6

GROUP 28 | 43 FUNGICIDES



INFINITO®

IE85324721d rA6

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate and when handling contaminated surfaces.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

WASH CONCENTRATE from skin or eyes immediately.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

To protect ground water do not apply this or any other products containing flupicolide at a rate of more than 400 g/ha in any three year period.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

PROTECT FROM FROST

SHAKE WELL BEFORE USE

READ ALL INSTRUCTIONS CAREFULLY BEFORE USE



To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.bayercropscience.ie/sds/infinito.pdf

or alternatively contact your supplier

DIRECTIONS FOR USE

INTRODUCTION

Infito combines the protectant activity of fluopicolide and the systemic and protectant properties of propamocarb hydrochloride.

RATE OF USE

Crop	Maximum individual dose	Maximum number of applications	Latest time of application
Potatoes (Seed and Ware)	1.6 L/ha	4 per season	7 days before harvest

RESTRICTIONS

Apply Infito only to dry foliage. Do not apply Infito if rainfall or irrigation is imminent. Avoid drift to areas outside those being sprayed having due regard to the prevailing weather conditions and the spray quality being used.

CROP SPECIFIC INFORMATION

Infito may be used on all varieties of potato and crops grown for seed

Dose rate

Infito should be applied as a foliar spray at 1.6 litre/ha

DISEASES CONTROLLED

When used as part of a full programme of correctly timed sprays, Infito gives control of late blight of potatoes (*Phytophthora infestans*).

Infito will protect tubers from the risk of late blight infection and reduce tuber blight incidence when used as part of a full blight protection programme.

Resistance management

To reduce the incidence of resistance developing, Infito should be used in single or block applications in alternation with fungicides from a different cross-resistance group. The maximum number of consecutive applications should be limited to three.

TIMING AND DOSE

Applications must be made preventively as part of a blight control programme with regard to Blight Infection Periods or local blight warnings. In the absence of any warnings, applications should commence when the crop meets along the row.

Apply Infito at 1.6 L/ha on a 7 day spray interval. Latest time of application: 7 days before harvest.

Do not apply Infito as a curative treatment when blight is present in the crop.

Tuber blight

Used as part of a full blight protection programme, Infito will protect tubers from the risk of late blight infection and reduce tuber blight incidence when applied at the recommended dose and spray intervals from full canopy development to haulm desiccation.

APPLICATION

Infinito must be applied to dry foliage by tractor mounted/trailed sprayer only.

Water volume

Apply in 200-400 litres of water per hectare

SPRAY QUALITY

Apply as a MEDIUM quality spray. Good cover of the potato plant foliage is essential.

COMPATIBILITY

Infinito may be applied as a tank-mix with a range of products. Contact Bayer CropScience for compatibility information on specific tank-mixes. Full manufacturer's instructions must be followed for each tank-mix component.

CONTAINER DISPOSAL

Empty containers should be disposed of in a safe way.

DISEASE RESISTANCE

Certain diseases may develop resistance to Bayer CropScience Limited's products. Since such circumstances are beyond our control, Bayer CropScience can accept no liability for any loss or damage whatsoever.

® Infinito is a Registered Trade Mark of Bayer

© Bayer CropScience Limited 2026

Specimen
2026 to date