



Cheetah[®] extra

5Le



For use only as an agricultural herbicide for the control of wild oats and grass weeds in winter and spring wheat.

An oil in water emulsion containing 69 g/L (7.14% w/w) fenoxaprop-p-ethyl

For professional use only

Approval Holder

Bayer CropScience Limited
230 Cambridge Science Park
Milton Road, Cambridge
CB4 0WB, UK

Marketing Company

Bayer CropScience Ltd.
Bayer Ltd.
The Atrium, Blackthorn Road
Sandyford, Dublin 18

Freephone: 1800 818534

For 24 hour emergency information contact

Bayer CropScience Limited Telephone: 1 800 409 399

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate. WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WHEN USING, DO NOT EAT, DRINK OR SMOKE. WASH CONCENTRATE from skin or eyes immediately. WASH HANDS AND EXPOSED SKIN before meals and after use.

Avoid contact with skin

If swallowed, seek medical advice immediately and show this container or label.

Environmental Protection

Extreme care must be taken to avoid spray drift onto plants outside of the target area.

Do not contaminate water with the product or its container.

(Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

Storage and Disposal

Keep out of reach of children.

Keep away from food, drink and animal feedingstuffs.

DO NOT RE-USE CONTAINER for any other purpose.

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Store in a safe dry place designated as an agrochemical store.

PROTECT FROM FROST.

READ ALL INSTRUCTIONS CAREFULLY BEFORE USE

Crop	Maximum individual dose	Maximum total dose	Latest time of application
Winter wheat, Spring wheat	1.2 L product per hectare	1.2 L product per hectare	Before flag leaf sheath extending stage (GS 41)

Safety Information

CHEETAH EXTRA

Contains 69 g/L (7.14% w/w) fenoxaprop-P-ethyl.



Warning

May cause an allergic skin reaction

Toxic to aquatic life with long lasting effects

Wear protective gloves/protective clothing/eye protection/face protection. If skin irritation or rash occurs: Get medical advice/attention. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site, except for triple rinsed containers which can be disposed of as non-hazardous waste.

Repeated exposure may cause skin dryness or cracking.

To avoid risks to human health and the environment, comply with the instructions for use.



IE79988516e rA3b

PCS No. 04314

DIRECTIONS FOR USE

RESTRICTIONS

For professional use only.

DO NOT apply Cheetah Extra to barley, durum wheat, undersown crops or those to be undersown.

DO NOT apply Cheetah Extra to wheat beyond the flag leaf ligule visible stage (GS 39) of the crop.

DO NOT roll or harrow within one week of spraying.

DO NOT spray crops under stress, suffering drought, waterlogged, grazed, lacking nutrients or if soil is compacted.

Broadcast crops should be sprayed post-emergence after the plants have a well-established root system and all other conditions on the label and leaflet are met.

Cheetah Extra can be sprayed in frosty weather provided the crop has hardened off.

Cheetah Extra is rainfast one hour after spraying.

DO NOT spray wet foliage liable to run off or leaves covered in ice.

Avoid spraying immediately before or after a sudden or extended drop in temperature or a period of warm days/cold nights.

Dry conditions resulting in moisture stress may reduce effectiveness. Such conditions will tend to occur with late applications to spring crops.

HERBICIDE RESISTANCE MANAGEMENT

This product contains fenoxaprop-P-ethyl which is an ACCase inhibitor, also classified by the Herbicide Resistance Action Committee as 'Group A'. Use only as part of a resistance management strategy that includes cultural methods of control and does not use ACCase inhibitors as the sole chemical method of grass-weed control.

Applying a second product containing an ACCase inhibitor to a crop will increase the risk of resistance development; only use a second ACCase inhibitor to control different weeds at a different timing.

Strains of some annual grasses (e.g. black-grass, wild oats and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced and copies are available from Teagasc, your distributor, crop adviser or product manufacturer.

To reduce the risk of developing resistance, applications should be made to young, actively growing weeds.

Use crop rotation and other cultural control measures to prevent and manage herbicide resistant grass weeds.

Apply when the majority of wild oats have emerged and if possible avoid late applications – beyond tillering and into the stem extension phase of wild oats.

continued →



cheetah extra

To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.bayercropscience.ie/sds/cheetahextra.pdf

or alternatively contact your supplier

IE/79980019f rA3b

Bayer

WEEDS CONTROLLED

Cheetah Extra controls emerged weeds on all soils and tilths and is unaffected by high organic matter or high Kd factor (ie soils with a high adsorptive capacity).

The rates of use of Cheetah Extra depend on weed type, size and time of application.

Apply post-emergence for the control of emerged weeds from the 2 fully expanded leaf stage of the weed up to the weed growth stage stated below:

Weed	Post-emergence control at 0.8 L/ha up to:	Post-emergence control at 1.0 L/ha up to:	Post-emergence control at 1.2 L/ha up to:
Awned canary-grass	GS 23	GS 30	-
Rough-stalked meadow -grass	GS 23	GS 30	-
Wild oats	GS 23	GS 30	GS 39

CROP SPECIFIC INFORMATION

For use on winter wheat and spring wheat.

Apply Cheetah Extra from emergence of the crop up to before flag leaf sheath extending stage (GS 41).

Cheetah Extra is rainfast from one hour after spraying.

Broadcast crops should be sprayed post-emergence after the plants have a well-established root system.

FOLLOWING CROPS and CROP FAILURE

In the event of crop failure, any broad-leaved crop may be sown, or after an interval of 3 weeks, any cereal may be sown. Following normal harvest of a treated crop, any broad-leaved or cereal crop may be sown.

MIXING AND SPRAYING

For use by tractor-mounter/trailed sprayer only.

For optimal results, use application techniques which ensure good weed coverage and crop penetration, using flat fan nozzles.

Conventional Volume

Volume: 200-300 litres of water per hectare

Always use the higher volume when weeds or crop are dense to avoid shading of the weed.

Pressure: 2.5-3.0 bars

Spray Quality: **FINE TO MEDIUM** (BCPC category)

Low Volume: 100 litres of water per hectare

Pressure: 2.5 bars

Spray quality: **FINE TO MEDIUM** (BCPC category)

Cheetah Extra may be applied at low volume providing conditions are optimal and all other recommendations are adhered to. Low volume can be used up to the early tillering stage of the crop (GS 23) after which growth may restrict spray penetration and conventional volumes should be used. Low volume is not advised where weed populations are dense.

SHAKE WELL BEFORE USE.

Add the recommended quantity of Cheetah Extra to the spray tank half-filled with the required quantity of clean water. Add the remainder of the water with the sprayer agitation system in operation. When emptying containers, wash out and add washings to the sprayer. Agitate before and during spraying. Do not leave the sprayer standing with chemical in it. After spraying, thoroughly wash out the spray tank.

COMPATIBILITY

Cheetah Extra may be applied as a tank-mix with a range of products. Contact Bayer CropScience for compatibility information on specific tank-mixes. Full manufacturer's instructions must be followed for each tank-mix component.

DO NOT mix Cheetah Extra with herbicides containing hormones.

Cheetah Extra can be used as part of a spray programme with any other spray chemical provided an interval of at least 7 days elapses between applications.

® Cheetah is a registered Trade Mark of Bayer

© Bayer CropScience 2016

SPECIMEN -
2016 to date