

Sipcam C50WG

A water dispersible granule formulation containing 50% w/w cymoxanil.

A fungicide for use on all varieties of early and main crop potatoes for the control of foliar late blight.

RISK AND SAFETY INFORMATION – 24 HOUR EMERGENCY NUMBER - +44 (0)1763 212100

WARNING



- Harmful if swallowed
- May cause an allergic skin reaction
- May cause damage to organs through prolonged or repeated exposure (blood and thymus)
- Suspected of damaging fertility and the unborn child
- Toxic to aquatic life with long lasting effects

- Wash hands thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Wear protective gloves/clothing and eye/face protection.
- If exposed or concerned get medical advice/attention.
- Collect spillage.
- Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.
- Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

PCS No. 04494

To avoid risks to human health and the environment, comply with the instructions for use.

Batch Number: See package

Registration Holder

SIPCAM OXON S.P.A.
Via Sempione 195,
20016 Pero (MI) Italy

Marketing Company

SIPCAM UK Ltd, 4C Archway House,
The Lanterns, Melbourn Street,
Royston, Herts SG8 7BX

SIPCAM C50WG is a registered trademark and product of SIPCAM OXON S.P.A.



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from sipcamuk.co.uk or contact your supplier.

sipcamuk.co.uk

FUNGICIDE

Net contents:

1 kg e



FOR PROFESSIONAL USE ONLY – DIRECTIONS FOR USE

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

| Crops | Maximum individual dose (product/hectare) | Maximum total dose (product / ha) | Latest time of application |
|--------|---|-----------------------------------|----------------------------|
| Potato | 0.24kg | 0.96kg | 7 days before harvest |

Other Specific Restrictions: A minimum interval of 7 days must be observed between applications.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

SAFETY PRECAUTIONS

Operator Protection

Operators must wear suitable protective clothing (coveralls) and suitable protective gloves when handling the product. Operators must wear suitable protective gloves when handling contaminated surfaces.

WHEN USING, DO NOT DRINK, EAT OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately. AVOID CONTACT WITH SKIN AND EYES.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF SWALLOWED, SEEK MEDICAL ADVICE IMMEDIATELY AND SHOW THIS CONTAINER OR LABEL.

Environmental Protection

DO NOT CONTAMINATE WATER WITH THE PRODUCT OR ITS CONTAINER.

DO NOT CLEAN APPLICATION EQUIPMENT NEAR SURFACE WATER. Avoid contamination via drains from farmyards and roads.

Storage and Disposal

- KEEP IN ORIGINAL CONTAINER tightly closed in a safe place DO NOT RE-USE CONTAINER for any purpose.
- KEEP OUT OF REACH OF CHILDREN.
- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.
- RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.
- Use appropriate containment to avoid environmental contamination.
- This material and its container must be disposed of as hazardous waste.

DIRECTIONS FOR USE

IMPORTANT

This information is authorised as part of the Product Label.

All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

SIPCAM C50WG must always be used in mixture with products containing mancozeb or fluazinam.

Disease Control

Blight in potatoes (*Phytophthora infestans*) is a disease which is variable in its occurrence and severity. SIPCAM C50WG allows the use of a flexible programme to retain continuous healthy foliage free from disease. The recommended mixes combine the local systemic activity of SIPCAM C50WG with protectant activity of MANCOZEB or FLUAZINAM.

The product is a foliar fungicide with a local systemic activity and protective and curative action. It also inhibits sporulation. SIPCAM C50WG will control strains of blight both resistant and sensitive to phenylamides.

Crops

All varieties of early and main crop potatoes may be treated with SIPCAM C50WG. On crops grown for processing consult your processor before use.

Mixing instructions

Spraying equipment should be thoroughly cleaned before spraying. SIPCAM C50WG should be added directly to the spray tank during filling. Continue agitation until spraying is completed.

Wash out sprayer thoroughly after use.

Spray quality

Apply SIPCAM C50WG as a MEDIUM spray.

Resistance

To reduce risk of resistance developing SIPCAM C50WG should be used as part of a spray programme of treatments using fungicides with different modes of action.

SIPCAM C50WG + approved mancozeb containing products

Application rate

Apply 0.24 kg SIPCAM C50WG plus products containing mancozeb in at least 200 litres of water per hectare.

Timing

A maximum of four applications may be applied.

Apply the first spray before the haulm meets across the rows or as soon as blight warning is received, whichever occurs first. Repeat every 10 days. A 7 day period may be used under exceptional circumstances where for instance, blight risk is already high and rain is forecast at the time the next spray is due.

SIPCAM C50WG + approved fluazinam containing products

Application rate

Apply 0.24 kg SIPCAM C50WG plus products containing fluazinam in at least 200 litres of water per hectare.

Timing

A maximum of four applications may be applied.

Apply the first spray before the haulm meets across the rows or as soon as blight warning is received, whichever occurs first.

Apply at 10 day interval in "high risk" situations - following two warm, humid days or irrigated crops.

Apply at 7 day interval in "severe risk" situations - during continue warm, humid weather and when blight is in nearby crops, potato dumps or volunteer plants.

Some reduction in tuber blight may occur when SIPCAM C50WG is used in mixture with fluazinam.

Compatibility

SIPCAM C50WG may be applied to recommended crops in a two-way tank-mix with approved products containing either fluazinam or mancozeb:

Please refer to mixing instructions.

In keeping with good spraying practice, tank mixtures should not be allowed to stand in the tank and agitation be maintained at all times.

Tank-mixed applications must be made within the label instruction of both products.

SIPCAM UK LTD CONDITIONS OF SUPPLY

All goods supplied by us are of high grade, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness of our goods for any purpose, are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.