

750ml

approx. measure



syngenta.

**GROUP 3 FUNGICIDE**


SCORE 250EC® is an emulsifiable concentrate containing 250 g/l difenoconazole (23.3% w/w).

500ml

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

SCORE 250EC controls a wide range of diseases in winter and spring oilseed rape, Brussels sprouts, head cabbage, kale, collard, broccoli, calabrese and cauliflower.

Authorisation Holder	Marketing Company
Syngenta UK Limited Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY Tel +44 (0) 1223 883400	Syngenta Ireland Limited Block 6, Cleaboy Business Park, Old Kilmeaden Road, Waterford. Tel: (051) 377203

In case of toxic or transport emergency ring (01484) 538444 anytime (24hr).

**PLEASE SEE ACCOMPANYING LEAFLET FOR PRODUCT USE DETAILS**

PROTECT FROM FROST  
SHAKE WELL BEFORE USE

© Syngenta AG, 2025

250ml

# 1 Litre

Product names marked ® or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company

**FOR PROFESSIONAL USE ONLY**

To avoid risks to human health and the environment, comply with the instructions for use.  
SCORE 250EC is an emulsifiable concentrate containing 250 g/l (23.3% w/w) difenoconazole.

**Danger.**

May be fatal if swallowed and enters airways.

Causes serious eye irritation.

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Repeated exposure may cause skin dryness or cracking.

Obtain special instructions before use.

Wear protective gloves/ protective clothing/ eye protection/ face protection/ hearing protection.

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/ physician.

IF exposed or concerned: Get medical advice/attention.

Do NOT induce vomiting.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty triple rinsed clean containers which can be disposed of as non-hazardous waste.

PCS No. 04566 UFI: 960R-1360-M00Y-MSPY



**CONDITIONS OF USE**

Crop	Maximum individual dose	Maximum number of applications	Maximum total dose	Latest timing of application
Oilseed rape	0.5l/ha	-	1.0 l/ha	Before end of flowering (GS69)
Brussels sprouts, cabbage, broccoli/ calabrese, cauliflower, kale and collard	0.5l/ha	-	1.0 l/ha	21 days before harvest

**Other specific restrictions:**

A minimum interval of 14 days must be observed between applications to brassica crops.

Applications to brassica must only be made to developed canopy and not before growth stage BBCH 41 as described below:

- Cabbage: heads begin to form, the 2 youngest leaves do not unfold.
- Kale, collard: 10% of the leaf mass typical for the variety reached.
- Brussels sprouts: lateral buds begin to develop.
- Cauliflower, broccoli, calabrese: heads begin to form, width of growing tip >1cm.

**ADDITIONAL SAFETY INFORMATION.**

**a) Operator Protection**

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces during spray application.

WASH SPLASHES from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.  
FOR USE VIA TRACTOR MOUNTED/TRAILED SPRAYERS ONLY.

**b) Environmental Protection**

To protect aquatic organisms, respect an unsprayed buffer zone of 5m to surface waters.

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

**c) Storage and disposal**

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place.  
RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

An emulsifiable concentrate containing 250 g/l (23.4% w/w) difenoconazole. SCORE 250EC is a DMI fungicide with contact and systemic activity against a wide range of diseases.

## DISEASES CONTROLLED

On oilseed rape, SCORE 250EC controls light leaf spot, leaf and pod spot and stem canker.

On Brussels sprouts, cabbage, broccoli, calabrese, kale, collard and cauliflower, SCORE 250EC controls leaf spot and ring spot.

## CROP SPECIFIC INFORMATION

### WINTER AND SPRING SOWN OILSEED RAPE

SCORE 250EC controls light leaf spot, leaf and pod spot (*Alternaria spp*) and stem canker (*Phoma lingam*).

#### Light Leaf Spot

Spray protectively with 0.5 litres SCORE 250EC per ha from the 4 expanded true leaf stage in the autumn where there is a high risk of disease. A repeat spray may be made in the spring at the beginning of stem extension if visual symptoms develop.

In areas of low disease risk, spraying may be delayed until the spring when crops should be sprayed at the first sign of disease.

For moderate control of light leaf spot only, a 2 spray programme of 0.25 litres per ha may be used, starting in the autumn.

#### Leaf and Pod Spot (*Alternaria spp*)

For effective control spray 0.5 litres SCORE 250EC per ha at the end of flowering.

#### Stem Canker (*Phoma lingam*)

Spray 0.5 litres SCORE 250EC per ha in the autumn after the 4 expanded true leaf stage when disease symptoms first occur. Repeat with an additional application of 0.5 litres per ha if foliar disease symptoms redevelop in late winter or early spring with a maximum total dose of no more than 1.0 litre per ha.

## ALTERNATIVELY

A 2 spray programme of 0.25 litres SCORE 250EC per ha may be used, starting in the autumn when disease symptoms first occur, with the second application 4-6 weeks after the initial application. Repeat with an additional application of 0.5 litres per ha if foliar disease symptoms redevelop in late winter or early spring with a maximum total dose of no more than 1.0 litre per ha.

## BRUSSELS SPROUTS, CABBAGE, BROCCOLI, CALABRESE, KALE, COLLARD AND CAULIFLOWER

SCORE 250EC controls leaf spot (*Alternaria spp*) and ring spot. A 2 spray programme should be used commencing at the first sign of disease but not before the canopy has developed to growth stage BBCH 41 as described below.

- Cabbage: heads begin to form, the 2 youngest leaves do not unfold
- Kale, collard: 10% of the leaf mass typical for the variety reached
- Brussels sprouts: lateral buds begin to develop
- Cauliflower, broccoli, calabrese: heads begin to form, width of growing tip >1cm

A minimum interval of 14 days must be observed between applications. Best control will result from spraying before disease is established in the crop.

## MIXING AND SPRAYING

### Mixing

Make sure the sprayer is set to give an even application at the correct volume.

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of SCORE 250EC and continue agitation while adding the rest of the water.

Agitate the mixture thoroughly before use and continue agitation during spraying.

Thoroughly wash all spray equipment with water immediately after use.

### Spray Volume

Oilseed rape	200 litres water per hectare
Brassica	400 litres water per hectare

### Spraying

Apply as a MEDIUM quality spray as defined. A spray pressure of 2-3 bars is preferred.

## RESISTANCE MANAGEMENT

SCORE 250EC is a DMI fungicide. Strains of Light Leaf Spot resistant to DMI fungicides are known to exist. To avoid development of resistance, apply product protectively in response to disease forecasts. Where possible, when Light Leaf Spot is present, avoid the use of DMI based fungicides when targeting other diseases such as Sclerotinia at mid flowering.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier