

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the product. WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces. WASH SPLASHES from skin immediately. WASH HANDS AND EXPOSED SKIN before eating and drinking or smoking and after work. WHEN USING DO NOT EAT, DRINK OR SMOKE. AVOID CONTACT WITH SKIN. WEAR SUITABLE GLOVES.

Environmental protection

DO NOT CONTAMINATE WATER with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads). Use appropriate containment to avoid environmental contamination.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. EMPTY CONTAINER COMPLETELY and dispose of safely. KEEP OUT OF REACH OF CHILDREN. KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS. THIS MATERIAL AND ITS CONTAINER must be disposed of in a safe way.

RESTRICTIONS

Allow a minimum of 7 days between applications. Only apply CYMBAL® 45 in tank-mixture with other fungicides containing a different active ingredient. The container must not be re-used for any purpose.

RESISTANCE

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programmes should include fungicides with different modes of action.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CROPS

CYMBAL® 45 can be used on all varieties of early and main crop potatoes. On crops grown for processing, consult your processor before use.

TIMING AND WEATHER

Timing of spray applications should take into account the normal prevalence of the disease in the area and conditions in the particular season. The first spray should be applied as soon as there is a risk of blight infection or an official warning has been issued, whichever occurs first. In the absence of a warning, spraying should begin just before the crop meets along the rows.

Applications should be repeated at 10-14 day intervals depending on the prevalence of the disease and favourable conditions for infection. A 7 day spray interval may be used under exceptional circumstances, e.g. blight risk is already high and rain is forecast at the time the next 10 day spray is due. High risk situations include conditions of rapid growth, irrigated crops, areas of intensive potato growing, particularly blight sensitive varieties, when further warnings are issued, or when active blight is in the area. The longer interval is recommended under dry conditions. Irrigation of crops increases the risk of blight and crops should always be sprayed with CYMBAL® 45 as soon as possible after irrigation subject to the minimum spray interval. Irrigated crops should normally be sprayed at 10 day intervals. Maximum 10 applications.

DOSE

CYMBAL® 45 should be applied at 0.25 kg/ha in tank-mixture with one of the following products at the recommended rate of application:

Fluazinam (500 g/L) e.g. Shirilan. This mixture will give a reduction in tuber blight.

OR

Cyazofamid e.g. Ranman Top. This mixture will give a reduction in tuber blight.

All other requirements and restrictions on the partner product label should be adhered to.

VOLUME AND APPLICATION

Apply CYMBAL® 45 in not less than 200 litres of water per hectare using a MEDIUM quality spray (as defined by BCPC).

MIXING

Spraying equipment should be thoroughly cleaned before spraying. CYMBAL® 45 mixes easily with water and should be added directly to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation until spraying is complete.

NOTICE TO BUYER

All goods supplied by us are of high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

CONDITIONS OF SUPPLY

The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade or otherwise and whether as to condition, quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort, for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumer's statutory rights are not affected.

cymbal[®] 45

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



WEBLINK to specific data sheet on manufacturer's website:

www.certisbelchim.co.uk/pdf/MSDS/Cymbal_IE_MSDS.pdf

Alternatively, contact your supplier.



cymbal[®] 45

A fungicide for the control of late potato blight.

A water dispersible granule (WG) formulation containing 45% w/w cymoxanil.

FOR PROFESSIONAL USE ONLY

IMPORTANT INFORMATION
FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (kg product/ha)	Maximum number of applications	Maximum total dose	Latest time of application
Potatoes (seed and ware)	0.25	10 per crop	2.5 kg/ha	7 days before harvest

Specific diseases controlled: Control of late blight of potatoes caused by *Phytophthora infestans* is achieved when used in conjunction with products containing active substances with multi-site activity

Other specific restrictions:

A minimum interval of 7 days must be observed between applications. Do not re-use container for any purpose.

USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE PRINCIPLES OF GOOD PLANT PROTECTION PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

CYMBAL 45 A water dispersible granule (WG) formulation containing 45% w/w cymoxanil.



WARNING

May cause an allergic skin reaction. Causes serious eye irritation. Suspected of damaging fertility. Suspected of damaging the unborn child. May cause damage to organs through prolonged or repeated exposure (blood, thymus, eyes). Very toxic to aquatic life with long lasting effects. Do not breathe spray. Avoid release to the environment.

UFI: 4HH8-J045-J00M-VR9X
RISK AND SAFETY INFORMATION

Wear protective gloves, eye protection. IF exposed or concerned: Get medical advice/attention. Collect spillage. Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for empty clean, triple rinsed containers which can be disposed of as non-hazardous waste. To avoid risks to human health and the environment, comply with the instructions for use.

PCS 04647

FUNGICIDE

Net contents

1kg

PROTECT FROM FROST

Production date / Batch number: see packaging

For advice on medical emergencies, fires, spillages or chemical hazards ONLY – phone Carechem 24 EU: +44 (0)1235 239670

For technical and non-emergency calls phone +44 (0)845 373 0305

Distributed by
Whelehan Crop Protection



Suite 12 Bunkilla Plaza
Bracetown Business Park
Clonee, Co. Meath Ireland
Technical Enquiries:
01 4688900

Authorisation Holder
Certis Belchim BV
Stadsplateau 16,
3521 AZ Utrecht,
PO Box 607
3500 AP Utrecht
Netherlands

© Registered Trademark of Certis Belchim BV



Certis Belchim
GROWING TOGETHER

Marketing Company: Certis Belchim BV
Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridgeshire CB21 6AD
Tel.: +44 (0)1223 652500 | Email: info.uk@certisbelchim.com
www.certisbelchim.co.uk

cymbal[®] 45