



An emulsifiable concentrate (EC) formulation containing 250 g/l tebuconazole. A fungicide for the control of a range of diseases in winter and spring wheat, winter and spring barley, oats, winter rye, triticale, and winter and spring oilseed rape.

RISK & SAFETY INFORMATION – 24 HOUR EMERGENCY NUMBER: +44(0)1763 212100

In the event of an emergency, call the National Poisons Centre, Beaumont Hospital at 01 809 2566 or 01 837 9964

DANGER



- Causes serious eye damage
- Suspected of damaging the unborn child



- Toxic to aquatic life with long lasting effects



- Wash hands thoroughly after handling.
- Wear protective gloves/clothing and eye/face protection.
- IF ON SKIN: Wash with plenty of soap and water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.
- If skin irritation or rash occurs: Get medical advice/attention.
- Avoid release to the environment.
- Collect spillage.
- Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.
- Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To avoid risks to human health and the environment, comply with the instructions for use.

UFI: 25NA-W6J3-810N-R4R8

PCN No. 04649

Registration Holder

SIPCAM OXON S.P.A.
Via Sempione 195,
20016 Pero (MI) Italy

Marketing Company

SIPCAM UK Ltd, 4C Archway House,
The Lanterns, Melbourn Street,
Royston, Herts SG8 7BX

FEZAN is a registered trademark and product of SIPCAM OXON S.P.A.



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from sipcamuk.co.uk or contact your supplier.

sipcamuk.co.uk

FUNGICIDE

Net contents:

5Le



FOR PROFESSIONAL USE ONLY
FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops	Maximum single dose	Maximum no. of applications at maximum single dose	Maximum total dose	Latest time of application
Winter & Spring Wheat	1 l/ha	-	2 l/ha	Before grain watery ripe stage (GS71)
Winter & Spring Barley	1 l/ha	-	2 l/ha	Before grain watery ripe stage (GS71)
Winter & Spring Oats	1 l/ha	-	2 l/ha	Before grain watery ripe stage (GS71)
Winter & Spring Triticale	1 l/ha	-	2 l/ha	Before grain watery ripe stage (GS71)
Winter Rye	1 l/ha	-	2 l/ha	Before grain watery ripe stage (GS71)
Oilseed Rape	1 l/ha	-	1 l/ha	End of flowering (BBCH69)

A minimum interval of 14 days applies between applications.

For use on oilseed rape a maximum total dose of 0.5L/ha can be applied before GS19 or a maximum total dose of 1.0 L/ha which can be applied between GS 20 and GS69.

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH SPLASHES from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating or drinking and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental Protection

Toxic to fish and other aquatic life. DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container. Use appropriate containment to avoid environmental contamination.

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies

Storage and Disposal

- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
- KEEP OUT OF REACH OF CHILDREN.
- KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place.
- RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or by manually rinsing three times. Add the washings to the sprayer at the time of filling and dispose of safely.
- DO NOT RE-USE CONTAINER for any purpose.
- This material and its container must be disposed of in a safe way.
- SHAKE THE CONTAINER WELL BEFORE OPENING AND USE.
- PROTECT FROM FROST.

DIRECTIONS FOR USE

Fungicide Resistance

FEZAN is a DMI fungicide. Repeated application of FEZAN alone on the same crop should be avoided. Tank mixture or alternation with fungicides with a different mode of action (e.g. morpholines) have been shown to protect against development of resistant forms of disease.

Resistance to some DMI fungicides has been identified in Septoria leaf blotch (*Mycosphaerella graminicola*) which may seriously affect the performance of some products. For further advice on resistance management in DMIs contact your agronomist or specialist advisor.

RECOMMENDATIONS FOR USE

As a general rule, disease control and yield benefit will be optimal when treatment is made at an early stage of disease development.

Apply FEZAN as a MEDIUM quality spray.

Add the required quantity of FEZAN to a half-filled spray tank with the agitation system in operation and fill to the required level. Maintain agitation whilst spraying and during stoppages. Spray immediately after mixing.

FEZAN is compatible in tank-mix with a range of products. Contact your distributor for compatibility information on specific tank-mixes. Always follow label instructions for each tank-mix component.

SPECIFIC DISEASES CONTROLLED

	Wheat	Barley	Oats	Rye	Triticale
Septoria tritici blotch (<i>Zymoseptoria tritici</i>)	R	-	-	R	R
Septoria glume blotch (<i>Septoria nodorum</i>)	MC	MC	-	MC	-
Yellow Rust (<i>Puccinia striiformis</i>)	C	C	-	-	C
Brown Rust	C	C	-	C	C
Fusarium ear blight	R	R	R	R	R
Sooty moulds	R	-	-	-	-
Powdery Mildew	MC	MC	C	C	MC
Crown rust	-	-	-	-	-
Net Blotch (<i>Pyrenophora teres</i>)	-	R	-	-	-
Leaf Scald (<i>Rhynchosporium secalis</i>)	-	R	-	R	R
C = control, MC = Moderate control, R = Reduction					
Oilseed rape:					
Control: Light leaf spot.					
Reduction: Phoma leaf spot/stem canker, Stem rot (<i>Sclerotinia sclerotiorum</i>)					

CROP RECOMMENDATIONS

May be used on all varieties of winter oilseed rape. If disease pressure remains high application of an alternative effective product may be required to maintain control and to comply with the timing restrictions necessary for FEZAN.

CEREALS

Rate of Application

1.0 litre/ha

Volume of Water

Apply in 200 - 300 litres of water/ha. Use higher water volumes in dense crops.

1. Control of Powdery Mildew (wheat, barley, oats and rye) – Apply FEZAN at the first signs of disease. When disease pressure remains high application may be repeated if necessary, but see considerations under ‘FUNGICIDE RESISTANCE’.

2. Control of *Septoria* Leafspot and *Septoria* Glume Blotch (wheat) – To protect the flag leaf and ear, apply FEZAN from the ‘flag leaf emerging’ stage (GS 37), up to the ‘ear fully emerged’ stage (GS 59).

Earlier application may be necessary if there is a high risk of *Septoria* leaf spot. Apply when disease is active but before it appears on the upper leaves. Applications made when foliar symptoms are already present on the upper leaves will be less effective.

3. Control of Yellow Rust and Brown Rust (wheat, barley and rye) – Apply as soon as the disease is seen. Applications made to established infestations will be less effective. Repeat after 2-3 weeks if re-infection occurs.

4. Control of *Fusarium* Ear Blights and Sooty moulds (wheat) – An application of FEZAN shortly after ear emergence can give a good reduction of *Fusarium* ear blight and a reduction of sooty moulds (*Alternaria* and *Cladosporium*).

5. Control of *Rhynchosporium* Leaf Blotch (barley and rye) – Apply FEZAN at the onset of disease. A second

application may be necessary 2-3 weeks later, but see considerations under ‘FUNGICIDE RESISTANCE’.

6. Control of Net Blotch (barley) – Apply at the first sign of disease in the spring or early summer. If conditions remain favourable for disease development, a second application should be made 2-3 weeks later. If the disease develops after flag leaf emergence, a single application should provide good protection.

7. Control of Crown Rust (oats) – Apply FEZAN on first appearance of the disease. Applications made to control mildew on oats will reduce Crown Rust infections occurring at the same time.

OILSEED RAPE

Rate of Application

0.5 – 1.0 litre/ha (see below)

Volume of Water

Apply in 100 - 400 litres of water/ha. Use higher water volumes in dense crops.

Light Leaf Spot

Apply 500 ml/ha FEZAN in autumn/winter (usually late October to early December). Further spray(s) of 1.0 L/ha may be required in the spring/summer depending on disease development.

When an autumn treatment of FEZAN has not been made and disease develops, apply FEZAN in early spring (March - after start of stem extension).

A later application of 1.0 L/ha of FEZAN pre- or post-flowering may be used to control late development of light leaf spot on pods and leaves.

Phoma Leaf Spot/Stem Canker

Use an autumn/early winter application of 500 ml/ha FEZAN at first signs of disease onset, followed by a further 500 ml - 1.0 litre FEZAN in late winter/early spring.

SIPCAM UK LTD CONDITIONS OF SUPPLY

All goods supplied by us are of high grade, but as we cannot exercise control over their mixing or use, all conditions or warranty, statutory or otherwise, as to the quality or fitness of our goods for any purpose are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

Specimen-
2024 to date