

# SYLLIT® 544 SC

FUNGICIDE

Fungicide against scab in apple and pear and leaf spot in cherry.

Contains 544 g/L dodine formulated as a suspension concentrate (SC)

PROFESSIONAL USE ONLY

	<p><b>Risk &amp; Safety Information</b></p> <p><b>DANGER</b></p> <p><b>Harmful if swallowed</b> <b>Causes serious eye damage</b> <b>Causes skin irritation</b> <b>Harmful if inhaled</b> <b>Very toxic to aquatic life with long lasting effects</b></p> <p>Wear protective gloves/protective clothing/eye protection/face protection IF SWALLOWED: Call a POISON CENTRE or doctor/physician IF SWALLOWED: rinse mouth IF ON SKIN: Wash with plenty of soap and water IF IN EYES: Rinse cautiously for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF IN EYES: Immediately call a POISON CENTRE or doctor/physician. IF INHALED: Call a POISON CENTER or doctor/physician. Collect spillage Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste</p> <p>Contains 1,2-benzisothiazole-3(2H)-one (CAS No. 2634-33-5). May produce an allergic reaction.</p> <p><b>To avoid risks to human health and the environment, comply with the instructions for use.</b></p>	<p><b>IMPORTANT INFORMATION</b></p> <p>FOR PROFESSIONAL USE ONLY AS A HORTICULTURAL FUNGICIDE.</p> <p>Crop: Apples, pears, cherries</p> <p>Maximum Individual dose: 1.25 L/ha</p> <p>Maximum number of treatments: 2 per year</p> <p>Maximum total dose: 2.5 L/ha</p> <p>Latest time of application: 60 days before harvest (apples, pears) 14 days before harvest (cherries)</p>
		<p><b>READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.</b></p> <p><b>SAFETY PRECAUTIONS</b></p> <p><b>Operator protection</b> WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. WASH HANDS AND EXPOSED SKIN before meals and after work. IN CASE OF AN ACCIDENT, OR IF YOU FEEL UNWELL, seek medical advice immediately and show label where possible.</p> <p><b>Environmental Protection</b> To protect aquatic organisms respect an unsprayed buffer zone of 20m to surface water bodies. Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.</p> <p><b>Storage and disposal</b> KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. WASH OUT CONTAINER THOROUGHLY and dispose of safely. DO NOT RE-USE CONTAINER for any other purpose.</p>
		<p>PROTECT FROM FROST</p> <p>Manufacturing date and Batch No.: see packaging</p> <p>Manufacturer, Marketing Company and Approval holder: Aysta LifeScience, Benelux sprl, Rue de Renory 26/1 B-4102 Ougrée, Belgium Tel. 00 32 4 385 9711</p>

UFI: 3FTE-74G5-1SKM-Q9AU

PCS no. 04739

24 hour emergency telephone number: +44 (0) 1235 239 670 (24 hour)

5Le



# SYLLIT® 544 SC

## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

## CROP SPECIFIC INFORMATION

SYLLIT 544 SC is a fungicide for preventative and curative use against *Venturia inaequalis/piri* (scab) on apples and pears and *Blumeriella jaapii* (= *Coccomyces hiemalis*) (leaf spot) on cherries.

Crop	Diseases controlled	Application rate	Pre-Harvest Interval	Application Timing
Apples, pears	<i>Venturia inaequalis</i> , <i>Venturia piri</i> (scab)	1.25 Litres product/ha	60 days	Before flowering (BBCH 53-59) and after flowering (BBCH 71-79). Maximum 2 applications with a spray interval of 7-10 days.
Cherries	<i>Blumeriella jaapii</i> (= <i>Coccomyces hiemalis</i> ) (leaf spot)	1.25 Litres product/ha	14 days	After flowering (BBCH 71-75) and/or after harvest (BBCH 90-97) Maximum 2 applications with a spray interval of 7-10 days.

Where tree height and/or canopy density is reduced, the dose (and water volume) should be adjusted in accordance with an appropriate dose adjustment scheme, for example PACE. Consult your specialist advisor for further information. Further information on the PACE scheme is available from the AHDB Apple Best Practice Guide, or PACE website.

## MIXING AND SPRAYING

To be applied using a conventional orchard sprayer. Half-fill the spray tank with water and add the required amount of product. Agitate and add the remaining quantity of water. Continue agitation during spraying.

## Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.