



Tebuconazole[®] 250 EW



**A broad spectrum systemic fungicide
for use on cereals and oilseed rape**

**Emulsion (oil in water) formulation,
containing 250 g/litre (25.8 % w/w) tebuconazole**

FUNGICIDE

CONTENT: 5 lit. e

! DANGER



RISK AND SAFETY INFORMATION

Causes serious eye damage

May cause respiratory irritation

Suspected of damaging the unborn child

Very toxic to aquatic life with long lasting effects

Keep out of reach of children

Obtain special instructions before use

Do not handle until all safety precautions have been read and understood

Avoid breathing dust/fume/gas/mist/vapours/spray

Wear protective gloves, face protection, eye protection

If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing

If exposed or concerned: Get medical advice/attention

Store in a well-ventilated place. Keep container tightly closed

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use

To protect aquatic organisms respect an unsprayed buffer Zone of 5m to surface water bodies

Do not contaminate water with the product or its container



ET1080A_ED2-V1

**Approval holder
and Marketing company:**

GLOBACHEM NV
Brustem Industriepark
Lichtenberglaan 2019
B-3800 Sint-Truiden, Belgium
Tel: +32 11 78 57 17
globachem@globachem.com
www.globachem.com



PCS no. 04827

FOR PROFESSIONAL USE ONLY

Tebuconazole is a registered trademark for Globachem N.V.

IE

Tebucur[®] 250 EW

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

TEBUCUR[®] 250 EW is recommended for control of a wide range of diseases on winter and spring sown cereals and oilseed rape. For best disease control and yield benefit TEBUCUR[®] 250 EW should be applied at an early stage of disease development, before infection spreads to new crop growth.

CROP SPECIFIC INFORMATION

WINTER AND SPRING WHEAT, WINTER AND SPRING BARLEY, OATS, TRITICALE AND RYE

Dose rate and application timing

TEBUCUR[®] 250 EW may be applied at any time before the grain milky ripe stage (GS 71).

Maximum individual dose: 1.0 litre per hectare

Maximum total dose per crop: 2.0 litres per hectare

Diseases controlled

	Wheat	Barley	Oats	Triticale	Rye
Powdery Mildew	C	C	C	R	C
Fusarium Ear blight	R	R	-	-	-

C: control / R: reduction

WINTER AND SPRING OILSEED RAPE

TEBUCUR[®] 250 EW may be used on all varieties of winter or spring sown oilseed rape.

Dose rate and application timing

TEBUCUR[®] 250 EW may be applied at any time before full flowering (GS 65).

1.0 litre per hectare.

Maximum individual dose: 1.0 litre per hectare

Maximum total dose per crop: 2 litres per hectare

DISEASES CONTROLLED

TEBUCUR[®] 250 EW controls *Alternaria*, *Sclerotinia* and *phoma leaf spot*. It also acts as a growth regulator.

Phoma leaf spot/stem canker

Leaf spot can be found from October onwards and best control of stem canker may be expected from an autumn/early winter application of 500 ml/ha TEBUCUR[®] 250 EW applied at first signs of disease, followed by 500 ml - 1.0 l/ha TEBUCUR[®] 250 EW in late winter/early spring. TEBUCUR 250 EW program applied against phoma leaf spot/stem canker will also give a secondary reduction of oilseed rape growth.

Dark Leafy Spot (Alternaria spp)

Treatment with 1.0 l/ha TEBUCUR[®] 250 EW should begin at the onset of disease.

Sclerotinia Stem Rot

Treatment with 1.0 l/ha TEBUCUR[®] 250 EW should be carried out at early to full flowering.

MIXING, SPRAYING AND TANK CLEANING

Important: Sprayers should be **THOROUGHLY CLEANED** before use and **filters and jets checked for damage and blockages.**

MIXING

Thoroughly shake the pack before use.

Add the required quantity of TEBUCUR[®] 250 EW to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing. Where tank mixes are used TEBUCUR[®] 250 EW should be added to the spray tank last, after first dispersing the other product(s).

SPRAYING

Apply through conventional crop spraying equipment using a medium quality spray (BCPC) at a pressure of at least 2 bar. Apply in at least 100-400 litres of water per hectare. Boom height and water volume should be adjusted to ensure good coverage of the crop, particularly at later growth stages. In dense crops at later growth stages, higher water volumes should be used as recommended.

TANK CLEANING

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at the time of filling and dispose of safely. Thoroughly wash all spraying and

measuring equipment with water and a wetting agent immediately after use. Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads.

RESISTANCE MANAGEMENT

The possible development of diseases resistant to TEBUCUR® 250 EW cannot be excluded or predicted. Where such resistant strains occur, TEBUCUR® 250 EW is unlikely to give satisfactory control. Repeated application of TEBUCUR® 250 EW alone should not be used on the same crop against a high risk pathogen such as cereal powdery mildew. Tank-mixtures or alternation with fungicides having a different mode of action (e.g. morpholines) have been shown to protect against the development of resistant forms of disease.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

SPECIMEN -
2020 to date



Tebucur[®] 250 EW

- ▶ **A broad spectrum systemic fungicide for use on cereals and oilseed rape.**
- ▶ Emulsion (oil in water) formulation, containing 250 g/litre (25.8 % w/w, tebuconazole)
- ▶ **PCS no. 04827**



FUNGICIDE

IMPORTANT INFORMATION
FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

CROPS/ SITUATIONS	MAXIMUM INDIVIDUAL DOSE	MAXIMUM NUMBER OF APPLICATIONS	MAXIMUM TOTAL DOSE	LATEST TIME OF APPLICATION
Winter wheat, spring wheat, winter barley, spring barley, oats, triticale and rye	1.0 l/ha	2	2.0 l/ha	Before grain watery ripe stage (GS 71)
Oilseed rape (winter and spring)	1.0 l/ha	2	2.0 l/ha	Before full flowering (GS 65)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.

PROTECT FROM FROST / SHAKE WELL BEFORE USE
BATCH NUMBER: SEE BOTTLE

FOR PROFESSIONAL USE ONLY

For 24 hour emergency information contact Globachem n.v.
Telephone: +32 11 78 57 17

Tebucur[®] is a registered trademark for Globachem N.V.



5 425030 786543

ET10805_ED2-V1



IE