

# tebusha® 25% EW

Oil in water emulsion containing 250 g/L tebuconazole.

Broad spectrum fungicide for use in winter and spring crops of wheat (excluding durum), barley, oats, rye, triticale and oilseed rape.

# 5L

## RISK AND SAFETY INFORMATION



**DANGER**  
May cause an allergic skin reaction  
Causes serious eye damage  
Suspected of damaging the unborn child  
Toxic to aquatic life with long lasting effects

Keep out of reach of children.

Do not eat, drink or smoke when using this product.

Avoid breathing vapours/spray.

Wear protective gloves/protective clothing/eye protection.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If skin irritation or rash occurs. Get medical advice/attention.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Do not contaminate water with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

**To avoid risks to human health and the environment comply with the instructions for use.**

**PCS No. 04884**

Registration holder: Sharda Europe bvba  
Jozef Mertenstraat 142,m 1702 Dilbeek, Belgium

Distributed by: Sharda Europe bvba  
Jozef Mertenstraat 142,m 1702 Dilbeek, Belgium.Tel.  
+32 349 41 18246 Emergency Tel: +91 22 6678 2800  
(8.00-16.00h)  
Protect from Frost

Batch No.see container

24 hour emergency number -  
National Poisons Centre: +353 (0)1  
809 2566 OR +353 (0)1 837 9964

©Registered trade marks and logos  
of Sharda Worldwide Exports Pvt. Ltd.

  
**Sharda Cropchem**  
SHARDA CROPCHEM LIMITED  
(previously Sharda Worldwide Exports Pvt. Ltd)  
Prime Business Park, Dasturthilal Joshi Road  
Vile Parle (West) Mumbai - 400059, India  
Tel. + 91 22 6678 2800; Fax: + 91 22 6678 2828

UFI:KARA-26QR-U70V-2WDN.



## PRECAUTIONS

If you feel unwell seek medical advice immediately (show the label where possible).

In case of accident, call the National Poisons Information Centre:

Tel: +353 (0)1 809 2566 or +353 (0)1 837 9964

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. Protect from frost.

## READ LABEL CAREFULLY BEFORE USE

IMPORTANT INFORMATION			
FOR USE ONLY AS N AGRICULTURAL FUNGICIDE			
Crops:	Maximum individual dose:	Maximum total dose:	Latest time of application:
Cereals (winter & spring wheat, winter & spring barley, oats, triticale and rye)	1 L/ha	2 L/ha	Before grain watery ripe stage (GS 71)
Oilseed rape (winter and spring)	1 L/ha	2 L/ha	Before full flowering (GS 65)

## INSTRUCTIONS FOR USE

For best disease control and yield, apply the product at an early stage of disease development, before infection spreads to new growth

To avoid the occurrence of resistance, do not apply the product or similar products with the azole-type active substance other than for preventive reasons or as soon as possible after the occurrence of infection. Do not rely on curative potential of this type of active ingredient.

To avoid occurrence of resistance, do not apply the product or similar products based exclusively on azole-type active ingredient in the crop consecutively without interruption by treatment using a different fungicide with a different mechanism of efficiency.

Avoid drift onto plants outside the target area due to risk of damage.

### Diseases Controlled-

**Wheat:** Reduction of *Septoria* leaf blotch, Glume blotch and *Fusarium* ear blight. Control of yellow rust, brown rust and powdery mildew.

**Barley:** Reduction of *Fusarium* ear blight. Control of brown rust and powdery mildew.

**Oat:** Control of powdery mildew and crown rust.

**Triticale:** Reduction of *Septoria* leaf blotch and Glume blotch. Control of yellow rust, brown rust and powdery mildew.

**Rye:** Control of yellow rust, brown rust and powdery mildew.

**Oilseed rape:** Control of *Alternaria*, *Sclerotinia* and *Phoma* leaf spot.

**Tebusha@25% EW** acts as a growth regulator in oilseed rape.

**Crop sensitivity:** For advice on crop sensitivity and replacement after crop failure, consult with distributor.

### Resistance

To prevent resistance always follow label recommendations and monitor

fields regularly for any signs of poor control. Use fungicides with different modes of action.

## CROP SPECIFIC RECOMMENDATIONS

### Wheat, barley, rye, triticale, oats

Apply at first sign of disease up until before grain watery ripe stage of growth. A second application may be required two to three weeks later.

### Water volumes

Most crops: 100 – 200 L/ha water

Dense crops after first node detectable stage: 250 L/ha water

Larger crops and varieties highly susceptible to disease: 300 L/ha water

**Oilseed rape:** Apply in 100 – 400 l/ha water at any time up to and including the full flowering stage.

## MIXING, APPLICATION AND CLEANING

Never prepare more spray solution than required.

Thoroughly shake pack before use.

When preparing the mixture, half fill the tank with clean water and start the agitation.

Add **Tebusha@25% EW** in the tank under agitation and then fill the tank with water. Continue agitation until spraying is completed. Apply using a tractor mounted/ trailed sprayer.

Use a medium quality spray to ensure a good and even spray cover of the target. A spray pressure of 2-3 bar is recommended.

Avoid spillage. In case of spill, remove contaminated clothes and wash skin immediately.

Clean equipment immediately after use with a spray tank cleaner.

Empty container and rinse with an integrated pressure rinsing device or manually rinsing three times.