

GAVEL®

MAPP 17586 / PCS 05089

GROUP 3 FUNGICIDE

A soluble concentrate containing 100 g/litre imazalil (9.3% w/w) for use as a protectant fungicide on potato for the reduction of silver scurf, gangrene, skin spot and *Fusarium* dry rot on seed potatoes.

FUNGICIDE

F

Net content

5L

PROTECT FROM FROST
SHAKE WELL BEFORE USE

Production date /
Batch number: see packaging

GAVEL

A soluble concentrate containing
100 g/litre imazalil (9.3% w/w)

DANGER



CAUSES SERIOUS EYE DAMAGE.
SUSPECTED OF CAUSING CANCER.
**VERY TOXIC TO AQUATIC LIFE WITH
LONG LAST EFFECTS.**

Keep out of reach of children.
Avoid release to the environment.

Wear protective gloves/
protective clothing/
eye protection/
face protection.

IF SWALLOWED:
Call a POISON
CENTRE or
doctor if
you feel unwell.

IF INHALED:
Remove person
to fresh air and keep
comfortable for breathing.

IF IN EYES: Rinse cautiously
with water for several
minutes. Remove contact
lenses, if present and easy
to do. Continue rinsing.
Immediately call a POISON
CENTRE/Doctor.
Dispose of contents/
container to a licensed
hazardous waste disposal
contractor or collection site
except for triple rinsed,
empty clean containers
which can be disposed of
as non-hazardous waste.

**To avoid risks to
human health and the
environment, comply with
the instructions for use.**

For advice on medical
emergencies, fires,
spillages or chemical
hazards ONLY – phone
Carechem 24 EU:
+44 (0)1235 239670

For technical and
non-emergency calls phone
+44 (0)845 373 0305

Authorisation Holder:
Certis Belchim B.V.,
Stadsplateau 16
3521 AZ Utrecht, PO Box 607
3500 Ap Utrecht, Netherlands

® Registered Trademark of
Certis Belchim B.V.

NI UFI: QJ99-F0HR-900G-A6WY IRE UFI: MEW9-T0D2-P00C-UVNK

MAPP 17586/PCS 05089

06012601_2403 • 107044148_UK-IE



Certis Belchim
GROWING TOGETHER

Marketing Company: **Certis Belchim B.V.**
Suite 5, 3 Riverside, Granta Park, Great Abington
Cambridgeshire CB21 6AD
Tel: +44 (0)1223 652500 | Email: info.uk@certisbelchim.com
www.certisbelchim.co.uk

IMPORTANT INFORMATION FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crops	Maximum individual dose	Maximum number of treatments (per batch)	Latest time of application
Potato (seed)	Up to 150 ml of product per tonne of potatoes	One	Post-harvest, prior to chitting

Other specific restrictions

This product must only be applied by conventional canopied hydraulic or spinning disc potato sprayer equipment mounted on a rolling conveyor.

Treated seed potatoes (including oversized tubers) are not to be used for human consumption or animal feed.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

THE CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH (COSHH) REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK – UK only.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective clothing:

OPERATORS MUST WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND EYE PROTECTION (GOGGLES) when handling treated seed and contaminated seed sowing equipment. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection. IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice. AVOID ALL CONTACT WITH SKIN.

TAKE OFF IMMEDIATELY all contaminated clothing. DO NOT BREATHE SPRAY.

DO NOT HANDLE seed unnecessarily.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating, drinking and smoking and after work.

IN CASE OF ACCIDENT OR IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Consumer protection

DO NOT RE-USE SACKS CONTAINING TREATED SEED for food or feed.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

KEEP TREATED SEED SECURE from people, domestic stock/pets and wildlife at all times during storage and use.

BURY OR REMOVE SPILLAGES.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose. SHAKE WELL BEFORE USE.

Store in a suitable pesticide store.

Keep dry and protect from frost.



The following does not form part of the product label under the Plant Protection Products Regulations 2005.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

The following precautions must appear on sack labels for treated seed:

THIS SEED HAS BEEN CHEMICALLY TREATED with GAVEL (contains imazalil).
WEAR SUITABLE PROTECTIVE GLOVES AND EYE PROTECTION (GOGGLES) when handling treated tubers and contaminated equipment during planting operations.
DO NOT HANDLE seed unnecessarily.
DO NOT USE TREATED SEED as food or feed.
DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE BEEN USED FOR TREATED SEED for food or feed.
KEEP TREATED SEED SECURE from people, domestic stock/pets and wildlife at all times during storage and use.
WASH HANDS AND EXPOSED SKIN before meals and after work.
BURY OR REMOVE SPILLAGES TO PROTECT BIRDS/WILD MAMMALS, treated seed should not be left on the soil surface. Bury or remove spillages.
This product is approved under the Plant Protection Products Regulations 2005

CONDITIONS OF SUPPLY: The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade or otherwise and whether as to condition, quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort, for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumers statutory rights are not affected.

GAVEL®

MAPP 17586/PCS 05089

FUNGICIDE

IMPORTANT INFORMATION
FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crops	Maximum individual dose	Maximum number of treatments (per batch)	Latest time of application
Potato (seed)	Up to 150 ml of product per tonne of potatoes	One	Post-harvest, prior to chitting

Other specific restrictions

This product must only be applied by conventional canopied hydraulic or spinning disc potato sprayer equipment mounted on a rolling conveyor. Treated seed potatoes (including oversized tubers) are not to be used for human consumption or animal feed.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

THE CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH (COSHH) REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK – UK only.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective clothing:
OPERATORS MUST WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate, handling contaminated surfaces, handling freshly treated material or applying the product.
WORKERS MUST WEAR SUITABLE PROTECTIVE GLOVES AND EYE PROTECTION (GOGGLES) when handling treated seed and contaminated seed sowing equipment. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.
IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.
AVOID ALL CONTACT WITH SKIN.
TAKE OFF IMMEDIATELY all contaminated clothing.
DO NOT BREATHE SPRAY.
DO NOT HANDLE seed unnecessarily.
WHEN USING DO NOT EAT, DRINK OR SMOKE.
WASH HANDS AND EXPOSED SKIN before eating, drinking and smoking and after work.

IN CASE OF ACCIDENT OR IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads.

Consumer protection

DO NOT RE-USE SACKS CONTAINING TREATED SEED for food or feed.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN. KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
KEEP TREATED SEED SECURE from people, domestic stock/pets and wildlife at all times during storage and use.
BURY OR REMOVE SPILLAGES.
RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.
DO NOT RE-USE CONTAINER for any purpose.
SHAKE WELL BEFORE USE.
Store in a suitable pesticide store.
Keep dry and protect from frost.

06012603-2403 • 107044148:UK-IE



GAVEL

A soluble concentrate containing
100 g/litre imazalil (9.3% w/w)

NI UFI: QJ99-F0HR-900G-A6WY IRE UFI: MEW9-T0D2-P00C-UVNK

DANGER

**CAUSES SERIOUS EYE DAMAGE.
SUSPECTED OF CAUSING CANCER.
VERY TOXIC TO AQUATIC LIFE WITH LONG LAST EFFECTS.**

Keep out of reach of children.

Avoid release to the environment.

Wear protective gloves/protective clothing/eye protection/
face protection.

IF SWALLOWED: Call a POISON CENTRE or doctor if you
feel unwell.

IF INHALED: Remove person to fresh air and keep
comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes.
Remove contact lenses, if present and easy to do.

Continue rinsing.

Immediately call a POISON CENTRE/Doctor.

Dispose of contents/container to a licensed hazardous
waste disposal contractor or collection site except for
triple rinsed, empty clean containers which can be
disposed of as non-hazardous waste.

**To avoid risks to human health and the environment,
comply with the instructions for use.**

MAPP 17586/PCS 05089

DIRECTIONS FOR USE

**IMPORTANT: This information is approved as part
of the Product Label. All instructions within this
section must be read carefully in order to obtain
safe and successful use of this product.**

GAVEL is a seed treatment fungicide for the
reduction of silver scurf, gangrene, skin spot and
Fusarium dry rot on seed potatoes.

Restrictions/Warnings

In situations where soil-borne inoculum is high,
treatment of seed tubers with GAVEL may not
give full protection against infection of stems and
daughter tubers.

In early potato seed crops, GAVEL can cause a
temporary reduction in the number of open eyes
when applied as an out of store treatment.
Treatment should be made before chitting to
minimise mechanical damage.
Reduction of established wound diseases may not
be achieved.

For best results the tubers should be clean and free
from soil (less than 5% of surface). Diseases under
areas of soil/dust will not be exposed to spray and
will not be controlled. The skin can be dry or damp
at treatment but not wet.

Disease Control

GAVEL reduces the following diseases of seed
potatoes, when applied during storage:-

Silver scurf	<i>Helminthosporium solani</i>
Skin spot	<i>Polyscytalum pustulans</i>
Gangrene	<i>Phoma exigua</i>
Dry rot	<i>Fusarium</i> spp

When applied prior to planting of seed potatoes as
a protective treatment, only reduction of skin spot
and silver scurf can be expected.

GAVEL should be used in conjunction with other
disease management measures such as, good
handling, curing and storage practice, good store
hygiene, and the use of resistant varieties.

Crops

GAVEL can be used on seed tubers of all varieties
of early and maincrop potatoes.

For best results the tubers should be clean and free
from soil (less than 5% of surface). Diseases under
areas of soil/dust will not be exposed to spray and
will not be controlled. The skin can be dry or damp
at treatment but not wet.

TIME OF APPLICATION

Either

To achieve the maximum disease reduction in
storage, GAVEL should be applied as soon as
possible after harvest, preferably within 7 to 10
days. To obtain the best reduction of both skin and
wound diseases application must be made before
wounds have begun to close. Later applications
may be less effective as wound infections will no
longer be exposed to treatment.

Or

A pre-planting treatment with GAVEL at final
grading prior to chitting, or prior to planting if
not chitted, will reduce skin spot and silver scurf
in the progeny crop. Treatment should be made
before chitting to minimise mechanical damage.
Reduction of established wound diseases may not
be achieved.

Rate of Application

Use GAVEL at 150 ml per tonne of seed potatoes.

Mixing and Spraying

Apply GAVEL through canopied, hydraulic or
spinning disc equipment (with or without
electrostatics) mounted on a rolling conveyor or
table. It may be applied diluted with water at up
to 2.0 litres of spray solution per tonne of potatoes
ensuring good coverage. Higher volumes within
the equipment's capabilities are preferred, to obtain
maximum cover of the skin, penetrate any wounds
etc and hence provide the best disease control.

Fill the tank with half the required volume of water.
Measure the appropriate amount of GAVEL into
the tank and make up the required volume of
water. Agitate thoroughly before application.

On emptying the container, RINSE CONTAINER
THOROUGHLY by using an integrated pressure
rinsing device or manually mixing three times. Add
washings to sprayer at time of filling and dispose
of container safely.

**To access the Safety Data Sheet
for this product scan the QR code
or use the weblink below:**

WEBLINK to specific data sheet on
manufacturer's website:



[https://certisbelchim.co.uk/pdf/
MSDS/Gavel_MSDS.pdf](https://certisbelchim.co.uk/pdf/MSDS/Gavel_MSDS.pdf)

Alternatively, contact your supplier.