

Fungicide

Skyway[®] Xpro

A fungicide for the control of stem-base, foliar and ear diseases in winter and spring wheat, triticale, winter rye and for disease control in oilseed rape.

An emulsifiable concentrate formulation containing 75 g/L (7.4% w/w) bixafen, 100 g/L (9.9% w/w) prothioconazole and 100 g/L tebuconazole (9.9% w/w).

Professional use only.

For 24 hour emergency information contact Bayer CropScience Limited +44 330 678 3382 (24 hr)
(charged as a standard international call to the UK)

**National Poisons Information Centre
Dublin: 01 809 2166**

5 L e



Safety information

SKYWAYXPRO

UFI: Y9G0-80MQ-5003-KNSV
Contains 75 g/L (7.4% w/w) bixafen, 100 g/L (9.9% w/w) prothioconazole and 100 g/L tebuconazole (9.9% w/w) and N,N-Dimethyl decanamide.



Warning

Harmful if swallowed

May cause an allergic skin reaction

May cause respiratory irritation.

**Suspected of damaging the unborn child
Very toxic to aquatic life with long lasting effects.**

Wear protective gloves/protective clothing/
eye protection/face protection

IF exposed or concerned: Call a POISON
CENTER or doctor/physician

Collect spillage

Protect from sunlight

Dispose of contents/container to a licensed
hazardous waste disposal contractor or
collection site except for triple rinsed empty
containers which can be disposed of as non-
hazardous waste.

**To avoid risks to human health and the
environment, comply with the instructions
for use.**

PCS No. 05259

GROUP 7 | 3 FUNGICIDES

F

Shake Well
Shake Well
Shake Well



SAFETY PRECAUTIONS

Operator Protection

WHEN USING DO NOT EAT,
DRINK OR SMOKE.

WASH ANY CONTAMINATION
from eyes immediately.

WASH HANDS AND
EXPOSED SKIN before meals
and after work.

IF YOU FEEL UNWELL, seek
medical advice (show the
label where possible).

Environmental Protection

Do not contaminate water with
the product or its container.
(Do not clean application
equipment near surface water.
Avoid contamination via drains
from farmyards and roads).

To protect aquatic organisms
respect an unsprayed buffer
zone of 10 m to surface water
bodies. Aim spray away from
water.

Storage and Disposal

Keep out of reach of children.

Keep away from food, drink
and animal feedingstuffs.

KEEP IN ORIGINAL
CONTAINER tightly closed in a
safe place.

RINSE CONTAINER

THOROUGHLY by using an
integrated pressure rinsing
device or manually rinsing
device or manually rinsing
three times. Add washings to
sprayer at time of filling and
dispose of safely.

DO NOT RE-USE CONTAINER
for any purpose.

PROTECT FROM FROST



To access the **Safety Data Sheet** for this
product scan the code or use the link below:

www.bayercropscience.ie/sds/skywayxpro.pdf

or alternatively contact your supplier

Authorisation holder: Bayer CropScience Ltd., PO Box 1582,
Cambridge, CB1 0FE, UK

Marketing company: Bayer CropScience Ltd, Bayer Ltd,
1st Floor, The Grange Offices, The Grange, Brewery Road,
Stillorgan, Co. Dublin A94 H2K7 Freephone: 1800 818534

Bayer

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

SKYWAY XPRO is a mixture of a carboxamide, triazole and triazolinthione fungicide recommended for control of a wide range of diseases on, winter and spring wheat, winter rye, triticale and oilseed rape.

RATE OF USE

| Crop | Maximum individual dose | Maximum total dose | Latest time of application |
|--|---------------------------------|--------------------------------|--|
| Winter and spring wheat, winter rye, triticale | 1.25 litres product per hectare | 2.5 litres product per hectare | Before grain milky ripe stage, (GS 73) |
| Oilseed rape | 1.0 litres product per hectare | 2.0 litres product per hectare | 56 days before harvest |

Other specific restrictions

A maximum of 2 foliar applications of product(s) containing SDHIs can be applied to any cereal crop.

| | |
|----------------------------------|---|
| Method of application: | Tractor mounted sprayer |
| Risk Mitigation measures: | To protect aquatic organisms respect an unsprayed buffer zone of 10 m to surface water bodies |

A spray pressure of 2-3 bar is recommended. Apply SKYWAY XPRO in 100-300 litres water per hectare.

Apply as a medium spray quality.

CROPS

SKYWAY XPRO may be used on all commercial varieties of winter and spring wheat, triticale, winter rye and oilseed rape.

APPLICATION

Water volume

Apply SKYWAY XPRO in 100-300 litres per hectare water. The higher spray volumes are recommended where the crop is dense or disease pressure / risk is high to ensure good penetration to the lower leaves and stem bases. Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve.

A spray pressure of 2-3 bar is recommended.

Spray quality

Apply as a MEDIUM spray quality (as defined by BCPC).

Latest Permitted Timing

In wheat, triticale and rye, SKYWAY XPRO may be applied at any stage before grain milky ripe stage (GS 73).

Oilseed rape up to 56 days before harvest.

Mixing

Thoroughly shake the pack before use. Add the required quantity of SKYWAY XPRO to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

General

Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages.

Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop. In dense crops, at later growth stages, higher water volumes should be used.

DISEASES CONTROLLED:

Cereals

| | Wheat | Triticale | Rye |
|--|-------|-----------|-----|
| Eyespot (<i>Oculimacula</i> spp.) | R | - | - |
| <i>Septoria</i> leaf blotch (<i>Mycosphaerella graminicola</i>) | C | C | C |
| Glume blotch (<i>Septoria nodorum</i>) | C | - | C |
| Powdery mildew (<i>Blumeria graminis</i>) | C | C | C |
| Yellow rust | C | C | - |
| Brown rust | C | C | C |
| Tan spot | C | - | - |
| <i>Fusarium</i> ear blight | MC | MC | - |
| Sooty moulds | R | R | - |
| <i>Rhynchosporium</i> leaf blotch | - | - | C |
| C = Control MC = Moderate control R = Reduction | | | |

Oilseed rape

| | |
|---|---|
| Light leaf spot | C |
| <i>Phoma</i> leaf spot and stem canker | C |
| <i>Sclerotinia</i> stem rot | C |
| C = Control MC = Moderate control R = Reduction | |

APPLICATION TIMING

Cereals

Eyespot (*Oculimacula spp.*)

Spray in the spring at the first sign of disease, from when the leaf sheaths begin to become erect until the 2nd node is detectable (GS 30-32).

Septoria Leaf Blotch (*S. tritici*) and Glume Blotch (*Mycosphaerella graminicola* and *Stagonospora nodorum*)

Apply before disease is established in the crop. To protect the upper leaves and ear apply SKYWAY XPRO at full flag leaf emergence (GS 37) up to mid-flowering (GS 65). Where disease pressure remains high application may be repeated.

Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective.

Powdery Mildew (*Blumeria graminis*)

Apply SKYWAY XPRO at the first signs of disease. Where disease pressure remains high application may be repeated.

Yellow Rust (*Puccinia striiformis*)

Apply SKYWAY XPRO at the first signs of disease. A second application may be made 2-3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

Brown Rust (*Puccinia triticina*, *P. recondita*)

Apply SKYWAY XPRO at the first signs of disease. A second application may be made 2-3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

Tan Spot (*Pyrenophora tritici-repentis*)

Apply SKYWAY XPRO at the first signs of disease in spring or early summer. Where disease pressure remains high, application may be repeated.

Ear Disease Complex

Apply SKYWAY XPRO soon after ear emergence until the end of flowering (GS 59-69) for moderate control of *Fusarium* ear blight in wheat and reduction of sooty moulds. Control of ear diseases can result in cleaner, brighter ears.

Leaf Blotch (*Rhynchosporium secalis*)

Apply SKYWAY XPRO in spring at the first signs of disease. For severe infections a second application may be necessary 2-3 weeks later.

OILSEED RAPE

Depending upon disease pressure and choice of disease management programme one application of SKYWAY XPRO may be made in the autumn followed a second in the spring/summer. Alternatively two applications may be made in the spring/summer.

Light Leaf Spot

Apply SKYWAY XPRO protectively in autumn/winter (usually late October to early December) after the first two leaves have unfolded. Follow up spray(s) may be required in late winter / early spring from the onset of stem elongation, depending on disease development. The second application of SKYWAY XPRO to oilseed rape must not be made prior to BBCH 21.

Phoma Leaf spot/Stem Canker

Apply SKYWAY XPRO in the autumn (after the first two leaves have unfolded) at the first sign of disease. Follow up spray(s) with another effective product may be required if disease symptoms reoccur.

Sclerotinia stem rot (*Sclerotinia sclerotiorum*)

Apply SKYWAY XPRO at early to full flower.

RESISTANCE STRATEGY

No more than two applications of SDH inhibitors must be applied to the same crop.

Isolates of certain cereal diseases with decreased sensitivity to DMI fungicides are known to exist. Where these occur or develop SKYWAY XPRO may not give satisfactory control.

Isolates of septoria leaf blotch with reduced sensitivity to SDHI fungicides have been detected. Where such resistant isolates occur, SDHI's are unlikely to give satisfactory control.

Strains of Light Leaf Spot resistant to azole fungicides are known to exist. To avoid development of resistance apply product protectively in response to disease forecasts. Where possible, when Light Leaf Spot is present, avoid the use of azole based fungicides when targeting other diseases such as Sclerotinia at mid flowering.

Mixtures of two or more SDHI fungicides do not provide an anti-resistance strategy. Each application of such a mixture counts as one SDHI application.

Tank mixtures or alternation with fungicides having a different mode of action against the diseases present have been shown to protect against the development of resistant forms of disease.

Do not tank mix SKYWAY XPRO with other SDHI containing fungicide formulations.

SKYWAY XPRO should be used in accordance with the instructions for use for the target diseases at the specific growth stages indicated.

Use SKYWAY XPRO as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control.

For further advice on resistance management contact your agronomist or specialist advisor and visit the FRAG-UK website.

CAUTION: The possible development of disease strains resistant to SKYWAY XPRO cannot be excluded or predicted. Where such resistant strains occur, SKYWAY XPRO is unlikely to give satisfactory control.

© Skyway and Xpro are Registered Trade Marks of Bayer

© Bayer CropScience Limited 2026