

# Sunfire

PCS 05497

GROUP 15 HERBICIDE

**SUNFIRE is a residual herbicide for use in winter sown cereals for the control of black-grass, annual meadow grass, loose silky bent and other annual grass weeds.**

A suspension concentrate containing 500 g/litre flufenacet.

**FOR PROFESSIONAL USE ONLY**

HERBICIDE

H

Net content

**5L**

**PROTECT FROM FROST  
SHAKE WELL BEFORE USE**

Production date / Batch number:  
see packaging

## SUNFIRE

A suspension concentrate containing 500 g litre flufenacet and 1,2-benzisothiazolin -3-one.

## HARMFUL IF SWALLOWED.

**MAY CAUSE AN ALLERGIC SKIN REACTION.  
MAY CAUSE DAMAGE TO ORGANS  
THROUGH PROLONGED OR  
REPEATED EXPOSURE.**

**VERY TOXIC TO AQUATIC LIFE  
WITH LONG LASTING EFFECTS.**

Avoid breathing vapours and spray.

Wash hands thoroughly after handling.

Wear protective gloves/ protective clothing/ eye protection/ face protection IF SWALLOWED:

Call a POISON CENTRE or a doctor/ physician if you feel unwell. Collect spillage.

Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty, clean containers which can be disposed of as non-hazardous waste.

**Contains flufenacet and 1,2-benzisothiazolin -3-one. May produce an allergic reaction.**

**To avoid risks to human health and the environment, comply with the instructions for use.**

**For advice on medical emergencies, fires, spillages or chemical hazards ONLY – phone Carechem 24 EU: +44 (0)1235 239670**

For technical and non-emergency calls phone +44 (0)845 373 0305

Authorisation Holder:  
**Certis Belchim BV**  
Stadsplateau 16  
3521 AZ Utrecht  
PO Box 607  
3500 Ap Utrecht  
Netherlands

**WARNING**

UFI: 328T-T07A-H002-G231

**PCS 05497**

® Registered Trademark  
of Certis Belchim BV

05015201\_XY • 108044149\_UK



**Certis Belchim**  
GROWING TOGETHER

Marketing company: **Certis Belchim BV**  
Suite 5, 3 Riverside, Granta Park, Great Abington  
Cambridgeshire CB21 6AD  
Tel: +44 (0)1223 652500 | email: info.uk@certisbelchim.com  
[www.certisbelchim.co.uk](http://www.certisbelchim.co.uk)

## IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations	Maximum individual dose (Litres product/ha)	Maximum total dose (Litres product/ha)	Latest time of application
Winter Cereals, wheat, barley, triticale and rye.	0.48	0.48	Up to third tiller detectable stage (GS 23)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

## SAFETY PRECAUTIONS

### Operator Protection

WHEN USING, DO NOT EAT DRINK OR SMOKE. WASH HANDS/EXPOSED SKIN before eating and drinking/smoking/after work.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY WITH PLENTY OF WATER AND SEEK MEDICAL ADVICE.

IF SWALLOWED, SEEK MEDICAL ADVICE IMMEDIATELY AND SHOW THIS CONTAINER OR LABEL.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE

GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

Treated crops should not be re-entered before spray deposits on leaf surfaces have completely dried.

### Environmental Protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

USE APPROPRIATE CONTAINMENT TO AVOID ENVIRONMENTAL CONTAMINATION.

To protect aquatic organisms respect an unsprayed buffer zone of 5 metres to surface water bodies.

### Storage and Disposal

KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

STORE IN ORIGINAL CONTAINER, tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

THIS MATERIAL AND ITS CONTAINER MUST BE DISPOSED OF IN A SAFE WAY.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

Protect against frost. Keep away from heat and direct sunlight.

FOR PROFESSIONAL USE ONLY.



## MIXING AND APPLICATION

Half-fill the spray tank and start the agitation. Shake the product several times before opening to ensure thorough mixing and add the required amount of SUNFIRE to the spray tank while re-circulating. Fill up the tank with water and continue agitation until spraying is complete. When using tank-mix partners, carefully follow the instructions on the partner product. Each product should be added separately to the spray tank and fully dispersed before the addition of any further product.

Ensure that all packaging is rinsed thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to the spray tank at the time of filling and dispose of container safely.

When applying the product, ensure even spray cover of the target weeds by using a suitable nozzle type. Apply SUNFIRE in 200-400 L/ha of water.

For tractor-mounted or trailed sprayers a travel speed of 6-8 km/h is advised to avoid the risk of uneven cover due to excessive boom whip and bounce. The slower speed is recommended where the target is in a thick crop or weed cover and the ground conditions are rough.

## COMPATIBILITY

SUNFIRE may be tank-mixed with a variety of plant protection products providing that the application timing is correct for both SUNFIRE and the partner(s) in the mixture.

When tank-mixing with other products use a sufficient water volume. Any use restrictions of products in the mixture should be observed. For further information on possible tank-mixes, consult your dealer representative.

## RESISTANCE MANAGEMENT

Strains of some annual grasses (e.g. black-grass, wild-oats, and Italian rye-grass) have developed resistance to herbicides, which may lead to poor control. A strategy for preventing and managing such resistance should be adopted.

Populations of black-grass and Italian rye-grass with high levels of enhanced metabolism resistance will not be fully controlled.

Maximise the use of cultural control measures wherever possible (e.g. crop rotation etc.)

Use tank-mixes or sequences of effective herbicides with different modes of action within individual crops, or successive crops.

For the control of herbicide resistant grass weeds, always use SUNFIRE in tank-mix or sequence with other effective graminicides with different modes of action.

Apply pre-emergence of weeds wherever possible. If applications are delayed, apply post-emergence products/mixtures to small actively growing weeds if high levels of resistance are suspected and to reduce the risk of resistance development.

## FOLLOWING CROPS

### After normal harvest:

There are no restrictions on following crops after the use of SUNFIRE. Plough to at least 150 mm before planting of all monocotyledon crop types, with the exception of wheat, barley, triticale, rye or maize.

### In the event of crop failure:

Plough to at least 150 mm. The following crops may then be sown; wheat, barley, triticale or rye with no restriction and any spring planted broad-leaved crop type following a minimum period of 8 weeks after application.

**Please refer to the following crop recommendations of any possible tank-mix or sequence partner products.**

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:

WEBLINK to specific data sheet on manufacturer's website:

[https://certisbelchim.co.uk/pdf/MSDS/Sunfire\\_MSDS.pdf](https://certisbelchim.co.uk/pdf/MSDS/Sunfire_MSDS.pdf)



Alternatively, contact your supplier.

# Sunfire

PCS 05497

HERBICIDE

## IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations	Maximum individual dose (Litres product/ha)	Maximum total dose (Litres product/ha)	Latest time of application
Winter Cereals, wheat, barley, triticale and rye.	0.48	0.48	Up to third tiller detectable stage (GS 23)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

## SAFETY PRECAUTIONS

### Operator Protection

WHEN USING, DO NOT EAT DRINK OR SMOKE. WASH HANDS/EXPOSED SKIN before eating and drinking/smoking/after work.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY WITH PLENTY OF WATER AND SEEK MEDICAL ADVICE.

IF SWALLOWED, SEEK MEDICAL ADVICE IMMEDIATELY AND SHOW THIS CONTAINER OR LABEL.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate. WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

Treated crops should not be re-entered before spray deposits on leaf surfaces have completely dried.

### Environmental Protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

USE APPROPRIATE CONTAINMENT TO AVOID ENVIRONMENTAL CONTAMINATION.

To protect aquatic organisms respect an unsprayed buffer zone of 5 metres to surface water bodies.

### Storage and Disposal

KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

STORE IN ORIGINAL CONTAINER, tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely. THIS MATERIAL AND ITS CONTAINER MUST BE DISPOSED OF IN A SAFE WAY.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE. Protect against frost. Keep away from heat and direct sunlight.

FOR PROFESSIONAL USE ONLY.

05015202\_XY • 108044149\_UK



## SUNFIRE

A suspension concentrate containing  
500 g/litre flufenacet and 1,2-benzisothiazolin-3-one.

UFI 328T-T07A-H002-G231



### WARNING

**HARMFUL IF SWALLOWED.  
MAY CAUSE AN ALLERGIC SKIN REACTION.  
MAY CAUSE DAMAGE TO ORGANS THROUGH  
PROLONGED OR REPEATED EXPOSURE.  
VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS.**

Avoid breathing vapours and spray.  
Wash hands thoroughly after handling.  
Wear protective gloves/ protective clothing/eye protection/  
face protection.  
IF SWALLOWED: Call a POISON CENTRE or a doctor/  
physician if you feel unwell.  
Collect spillage.

Dispose of contents/container to a licensed hazardous waste  
disposal contractor or collection site except for triple rinsed  
empty, clean containers which can be disposed of as  
non-hazardous waste.

**Contains flufenacet and 1,2-benzisothiazolin-3-one. May  
produce an allergic reaction.**

**To avoid risks to human health and the environment,  
comply with the instructions for use.**

## DIRECTIONS FOR USE

**IMPORTANT: This information is approved as part of  
the Product Label. All instructions within this section  
must be read carefully in order to obtain safe and  
successful use of this product.**

SUNFIRE is a herbicide with residual activity for the  
control of black-grass, loose silky bent and annual  
meadow grass in winter sown cereals.

## CROP SPECIFIC INFORMATION

### CEREALS

SUNFIRE can be used on all varieties of winter sown  
wheat, barley, triticale and rye.

Apply from the pre-emergence up to third tiller stage  
(GS 23) of the crop in the Autumn.

## WEEDS CONTROLLED

Susceptibility of weeds to single application of SUNFIRE.

Weeds are controlled both pre-emergence and post-emergence.

Weed susceptibility	L/ha Pre-emergence use	Pre-emergence	L/ha Post-emergence use	Post-emergence
Black-grass	0.48	MS	0.48	MS
Loose Silky Bent	0.36	S	0.36	S
Annual Meadow Grass	0.36	S	0.36	S

S= Susceptible

MS = Moderately susceptible

MT = Moderately tolerant

## RESTRICTIONS

Soil moisture must be present for SUNFIRE to work.  
Best results will be obtained if rainfall occurs within  
7 days of application. Residual control may be reduced  
under prolonged dry conditions, on soils with a high  
kd factor, where organic matter exceeds 6%, and in  
soils where the ash content is high. Do not disturb the  
soil after SUNFIRE has been applied as this will result  
in reduced weed control.

Do not use in windy conditions in order to avoid the  
risk of drift onto other crops.

Avoid spraying during periods of prolonged or severe  
drought or frost.

Where there is very wet weather before and after  
application and particularly where soils become  
water-logged, reduction in crop vigour which may  
result in yield reductions may occur.

## Soil Types

SUNFIRE may be used on all mineral soil types. Do  
not use on soils with more than 10% organic matter.  
On stony or gravelly soils, crop damage could occur,  
particularly if heavy rain follows treatment. Control of  
black-grass may be reduced on soils with a high clay  
content.

Do not use on waterlogged soils or soils prone to  
water logging.

## Seedbed preparation

Trash and straw should be incorporated evenly to a  
depth of 15 cm during seedbed preparation. Seedbed  
must have a fine, firm tilth. Consolidate loose or  
cloddy seedbeds before use.

Following pre-emergence applications, unconsolidated  
clods, if they are larger than 7.5cm in diameter may  
reduce the level of weed control and cause seed to be  
inadequately covered, which could result in crop damage.

## Crop Safety

Extreme caution must be taken to avoid spray drift  
onto non target crops.

Do not use on crops suffering from stress, which  
may be caused for example by pests, disease, water  
logging, poor seedbed preparation or previous  
chemical treatment.

Seed should be covered with a minimum of 3.2 cm of  
settled soil.

Shallow drilled crops should be treated  
post-emergence.

Do not soil incorporate. Do not spray undersown crops.