



# EXTERIS®

## + Stressgard®

F

### Fungicide

✓ Exteris® Stressgard is a systemic and mesostemic fungicide which provides moderate control of Microdochium Patch (*Microdochium nivale*) and control of Dollar Spot (*Sclerotinia homoeocarpa*) when used in preventative or early curative situations on bowling greens, cricket pitches, lawn tennis courts, professional sports grounds/stadiums and golf greens, tees and fairways.

An aqueous suspension concentrate containing 12.5 g/L fluopyram and 12.5 g/L trifloxystrobin

#### For Professional use only

##### Exteris® Stressgard

Contains 12.5 g/L fluopyram and 12.5 g/L trifloxystrobin.

May cause an allergic skin reaction.

May cause harm to breast-fed children.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/protective clothing/eye protection/face protection.

IF ON SKIN: Wash with plenty of water.

If skin irritation or rash occurs: Get medical advice/attention.

Do not breathe the gas/mist/vapours/spray.

Avoid contact during pregnancy/while nursing.

Dispose of contents/container to a licenced hazardous

waste disposal contractor or collection site except

for triple rinsed empty containers which can be

disposed of as non-hazardous waste.

To avoid risks to human health and the

environment, comply with the instructions for use.

Contains trifloxystrobin and 1, 2-benzisothiazol-3(2H)-one.

May produce an allergic skin reaction.

UFI : 1421-POXA-400R-GM84



**WARNING**

GB MAPP 20242 – NI MAPP 20241  
PCS 05719

GROUP 7 | 11 FUNGICIDE



envu™

5 L e

DPM00001298A - A3 - ARTICLE D00001313

Preventative Microdochium  
Patch disease management



# EXTERIS®

## + Stressgard®

**IMPORTANT INFORMATION**  
FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Situation: Managed amenity turf.  
Maximum individual dose: 10 L/ha  
Maximum number of treatments: Two per year

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**



DPM00001299A - A3 - **ARTICLE D00001313**

24 hours emergency information contact: Telephone +44 20 3807 3798

Or

 GB: nearest National Poisons Information Centre

 IE: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.

Further information is available from: 00800 12149451

[www.uk.envu.com](http://www.uk.envu.com) for SDS and larger label.

**Authorisation Holder**

**Great Britain**

2022 Environmental Science FR SAS  
Milton Hall, Ely Rd, Milton, Cambridge, CB24 6WZ

**Ireland and Northern Ireland**

2022 Environmental Science FR SAS  
1 Place Giovanni Da Verrazzano 69009 Lyon, France

Batch Number : see on the pack

Envu, the Envu logo and Exteris® Stressgard® are trademarks owned by Environmental Science U.S. LLC. or one of its affiliates

## SAFETY PRECAUTIONS


### Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment :

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces


WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment

 However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

### Environmental Protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

 To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements. DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water. This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Use appropriate containment to avoid environmental contamination.


This material and its container must be disposed of in a safe way.

 To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

### Storage and Disposal

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

WASH OUT CONTAINER THOROUGHLY and dispose of safely.

 Triple rinsed containers should be punctured to prevent re-use and may be disposed of by an authorised contractor or at a municipal waste recycling site.

**READ ALL INSTRUCTIONS CAREFULLY BEFORE USE.**



### IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Situation: Managed amenity turf.  
Maximum individual dose: 10 L/ha  
Maximum number of treatments: Two per year

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**



### GENERAL INFORMATION:

Exteris® Stressgard® is a systemic and mesostemic fungicide which provides moderate control of Microdochium Patch (*Microdochium nivale*) and control of Dollar Spot (*Sclerotinia homoeocarpa*) when used in preventative or early curative situations on bowling greens, cricket pitches, lawn tennis courts, professional sports grounds/stadiums and golf greens, tees and fairways.



### RESTRICTIONS:

Exteris® Stressgard® is not to be used on commercial, residential or estate lawns.

Do not apply during drought conditions or to frozen turf.

If grass is being mown, Exteris® Stressgard® should be applied after cutting and further mowing should not take place until AT LEAST 24 hours after treatment.

Do not apply more than 2 applications per year.

A minimum of two weeks must be observed between treatments.



### DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

PROTECT FROM FROST



### APPLICATION

For preventative and early curative moderate control of Microdochium Patch (*Microdochium nivale*) and preventative and early curative control of Dollar Spot (*Sclerotinia homoeocarpa*)

The product is to be applied as a foliar spray with a rate of 10 L/ha diluted in 200 to 600 L of water per hectare of turf.

For use with knapsack sprayers or vehicle mounted hydraulic sprayers.

Application Equipment			
Vehicle-mounted Hydraulic Sprayer		Knapsack Sprayer	
Rate per hectare	Water per hectare	Rate per 100 m <sup>2</sup>	Water per 100 m <sup>2</sup>
10 Litres	200-600 litres	100 mls	2 -6 litres
		Rate per 500 m <sup>2</sup>	Water per 500 m <sup>2</sup>
		500 mls	10-30 litres

### **Mixing:**

Sprayers should be THOROUGHLY CLEANED before use and filters and jets checked for damage and blockages. Thoroughly shake the pack before use.

Half fill the sprayer tank with clean water and agitate. Add the required amount of Exteris® Stressgard®.

Mix thoroughly and make up to the required volume of water while still agitating.

Spray immediately after mixing. Continue agitation at all times during spraying and stoppages, until the tank is completely empty.

Ensure an even distribution on the turf to be treated

Wash out all spray equipment thoroughly with water immediately after use.

### **Application Timing**

Exteris® Stressgard® is used as a foliar-applied fungicide and for best results should be used when the grass is actively growing.

#### **Microdochium Patch (*Microdochium nivale*).**

For best results Exteris® Stressgard® should be used preventatively or early curative before or at the first occurrence of symptoms, usually early autumn when weather conditions are conducive to disease development. A second application can be made if necessary, usually 28 days (minimum 14 days interval).

#### **Dollar Spot (*Sclerotinia homoeocarpa*).**

Exteris® Stressgard® can be used preventatively or curatively. For best results apply Exteris® Stressgard® at the first occurrence of symptoms, usually early summer to autumn when weather conditions are conducive to disease development. A second application can be made if necessary, usually 28 days (minimum 14 days interval).

Exteris® Stressgard® can be applied year round.

### **Resistance Management:**

Exteris® Stressgard® contains active ingredients from strobilurin class of chemistry, trifloxystrobin, and the SDHI class of chemistry, fluopyram.

The use of Exteris® Stressgard® should conform to resistance management strategies. Such strategies should include using in rotation with products containing an active substance with a different mode of action which is effective against the target disease, limiting the total number of applications per season, and using non-chemical methods of disease control. Envu recommends responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

Maximum number of treatments: 2 per year.

This strategy should include using in rotation with products that have a different mode of action or different chemical group that are effective against the target disease.