

Zypar[®]
Arylex[®] active

HERBICIDE

™ ® Trademarks of Corteva Agriscience and its affiliated companies

SAFETY INFORMATION

ZYPAR[®]

UFI: 40A7-TODN-S002-N4HU



WARNING

Causes skin irritation
May cause an allergic skin reaction
Causes serious eye irritation
Very toxic to aquatic life with long lasting effects

Wear protective gloves / clothing / eye / face protection.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 05800

Authorisation Holder:

Corteva Agriscience UK Limited

Unit H4, Building H, Melbourn Science Park,
Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB, UK.
Technical enquiries: +44 800 689899 or email: ukhotline@corvea.com
24 Hour Emergency Telephone Number: +353 816 663 627
National Poison Centre (Dublin): 01 8092166

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.
DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking, and after work.

Environmental protection:

Do not contaminate water with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and disposal:

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET

PROTECT FROM FROST
SHAKE WELL BEFORE USE

Non-printable area

0 2 8 7 8 6 4

2411

Last updated by :

PUBLI H



Zypar[®]

Arylex[®] active

HERBICIDE

TM Trademarks of Corteva Agriscience and its affiliated companies

GROUP 4 | 2 HERBICIDES

Product Registration Number: PCS No. 05800

An oil dispersion formulation containing 5 g/L florasulam + 6.25 g/L halauxifen-methyl (equivalent to as 6 g/L halauxifen acid) and 6 g/L of cloquintocet-mexyl (safener).

ArylexTM is the commonly used name of halauxifen-methyl active substance.

A post-emergence herbicide for use on a variety of winter cereals (wheat, durum wheat, spelt, barley, rye, triticale, oats and these crops under sown with grass) and spring cereals (wheat, durum wheat, spelt, barley, rye, triticale and these crops under sown with grass) and GRASSLAND (established, newly sown leys and for seed production) for the control of a wide range of broad-leaved weeds.

FOR PROFESSIONAL USE ONLY

IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Crops: Barley, Durum Wheat, Rye, Spelt, Triticale, Wheat (Undersown with grass), Oat (Winter, undersown with grass)
Grassland (established, newly sown leys and seed crops)

Maximum Individual Dose: }
Maximum n° of applications: } Full details are given in the Important
Maximum Total Dose: } Information area on the attached leaflet
Latest Time of Application: }
Other Specific Restrictions: }

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

NET CONTENTS : 5 LITRES e

0297939

2512

Last updated by : PUBLI H

SAFETY INFORMATION

ZYPAR®

UFI: 40A7-T0DN-S002-N4HU



WARNING

Causes skin irritation

May cause an allergic skin reaction

Causes serious eye irritation

Very toxic to aquatic life with long lasting effects

Wear protective gloves / clothing / eye / face protection.

IF ON SKIN: Wash with plenty of soap and water.


IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 05800



SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking, and after work.

Environmental protection:

DO NOT CONTAMINATE WATER with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and disposal:

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET

PROTECT FROM FROST

SHAKE WELL BEFORE USE

IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Crops	Maximum Individual Dose	Max No of applications	Maximum Total Dose	Latest Time of Application
Barley, Durum Wheat, Rye, Spelt, Triticale and Wheat	1.0 L/ha (see Other Specific Restriction 2)	1 per crop	1.0 L/ha	30 th June Late boot stage (GS 45)
Barley, durum wheat, rye, spelt, triticale, wheat (Undersown with grass)	1.0 L/ha (see Other Specific Restriction 2)	1 per crop	1.0 L/ha	June 30 th Late boot stage (GS 45)
Oat (winter), Oat (Winter, undersown with grass)	0.5 L/ha	1 per crop	0.5 L/ha	May 31 st (GS 13 – 32)
Grassland (established, newly sown leys, seed crops)	1.0 L/ha (see Other Specific Restrictions)	1 per year	1.0 L/ha (see Other Specific Restrictions)	14 days before grazing/cutting

Other Specific Restrictions:

1. Do not apply more than 0.75 litre product per hectare to any crop before 14th February.
2. The total amount of halauxifen-methyl applied to a winter cereal crop must not exceed 14.04 g/ha (13.5 g a.e/ha of halauxifen-methyl). For autumn sown crops the maximum total must not exceed 7.8 g/ha (7.5 g a.e/ha of halauxifen-methyl) between first leaf unfolded (GS11) and 31st December. For any applications involving halauxifen-methyl made between 1st January and flag leaf swollen (GS 45) the maximum total dose of halauxifen-methyl applied must not exceed 6.24 g/ha (6 g a.e/ha). A minimum interval of 3 months between applications of products which contain halauxifen-methyl must be respected.
3. For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and 14th February in the year of harvest. The total amount of florasulam applied to a cereal crop must not exceed 7.5 g.
4. For applications made to grassland from 1st September to 14th February, do not apply more than 0.75 litre of ZYPAR per hectare.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

ZYPAR is mainly absorbed through the foliage of the target weeds. The ideal timing is when the weeds are small and actively growing.

DO NOT treat crops under stress. Stress can be caused by many factors including frost, drought, water logging, trace element deficiency, disease and pest attack etc. ZYPAR can sometimes induce slight and transient (2-3 weeks) crop yellowing. This has no impact at all to the yield.

When applying ZYPAR to spelt, transient growth inhibition and chlorosis may be observed; however, it should have no adverse effect on yield.

DO NOT apply to crops containing or undersown with clover or any legume containing mixtures.

DO NOT roll or harrow 7 days before or after application. ZYPAR may be used on all soil types.

Weeds not present at the time of application will not be controlled.

WEED RESISTANCE

ZYPAR is a product containing Arylex™ (halauxifen-methyl) active substance and florasulam. It also contains a safener (cloquintocet-mexyl). Arylex™ is a member of the aryloxyacetate family of synthetic auxins (Group 4 according to HRAC classification) and the risk of resistance developing to this active is low. Florasulam belongs to the triazolopyrimidine (Group 2, HRAC classification) and as an ALS inhibitor there is a risk of resistance building to this active ingredient, therefore precautions should be taken to minimize the risk.

The following resistance risk strategies are advised:

1. Do not rely on a single mode of action.
2. Avoid the repeated use of florasulam
 - a. where resistance to other ALS inhibitors has developed
 - b. and to species which are considered high risk species for the ALS herbicides such as Common Poppy and employ alternation or mixtures with herbicides with another mode of action.
3. Follow label recommendations concerning rates and timing of application. Use ZYPAR in a way to maximise control of high risk/difficult to control weed species.
4. Consider the use of cultural control and crop rotation to help control prevent development of resistance biotypes.

CROP SPECIFIC INFORMATION

CEREALS

ZYPAR can be used on a variety of winter cereals (wheat, durum wheat, spelt, barley, rye, triticale and oats) and spring cereals (wheat, durum wheat, barley and rye) for the control of broad leaf weed species.

ZYPAR can be used on a variety of winter cereals (wheat, durum wheat, spelt, barley, rye, triticale and oats) and spring cereals (wheat, durum wheat, barley and rye) under sown with grass for the control of broad leaf weed species.

ZYPAR can be used in cereals grown for seed and malting barley.

Timing

Winter sown cereals

ZYPAR can be applied from 1st September to the following crops: wheat, durum wheat, spelt, barley, rye, triticale (and these crops under sown with grass) from GS 11 (first leaf unfolded) up to GS 45 (flag leaf sheath swollen). The latest calendar date for application is 30th June.

Zypar can be applied from 1st September when winter oats and winter oats (under sown with grass) is at GS13 (three leaves unfolded) up to GS 32 (second node detectable). The latest calendar date for application is 31st May.

Spring sown cereals

ZYPAR can be applied from 15th February when the crop is at GS13 (three leaves unfolded) up to GS 45 (flag leaf sheath swollen) for all cereal types (except oats), including cereals under sown with grass. The latest calendar date for application is 30th June.

The ideal timing for application is when the weeds are small and actively growing. However, some highly susceptible species can be controlled at advanced growth stages for rescue treatments (e.g. cleavers and red dead-nettle).

Undersown cereal crops

ZYPAR may be applied to wheat, barley, winter oats, rye, durum wheat, spelt and winter triticale crops undersown with grass. Application should be made when the grasses have started tillering (GS 21) and when weeds are actively growing.

Weeds controlled

For autumn sown crops the maximum total dose must not exceed 0.75 L/ha when applied from 1st September to 14th February and 1.0 L/ha when applied from 15th February to 30th June, except for winter oats, for which the maximum dose rate is 0.5 L/ha.

Application rates and timing for autumn applications (1st September to 14th February) to winter cereals:

Weed species	Recommended rate L/ha	Timing crop	Maximum growth stage controlled	Susceptibility
Volunteer oilseed rape	0.75	From first leaf unfolded to end of tillering	7 true leaf stage 2-10 cm diameter	S
Cleavers	0.75	From first leaf unfolded to end of tillering	2 true leaf stage 2-6 cm diameter	S

Weed species	Recommended rate L/ha	Timing crop	Maximum growth stage controlled	Susceptibility
Red dead-nettle	0.5	From first leaf unfolded to end of tillering	2 true leaf stage 2-3 cm diameter	S
Scented mayweed, Scentless mayweed	0.75	From first leaf unfolded to end of tillering	4 side shoots visible 1-25 cm diameter	S
Common poppy	0.75	From first leaf unfolded to end of tillering	6 side shoots visible 5 cm diameter	S
Common chickweed	0.75	From first leaf unfolded to end of tillering	3 side shoots visible 1-10 cm diameter	S
Ivy-leaved speedwell	0.75	From first leaf unfolded to end of tillering	2 true leaf stage 1-2 cm diameter	S

Application rates and timing for spring applications (15th February to 30th June) to winter cereals:

Weed species	Recommended rate L/ha	Timing crop	Maximum growth stage controlled	Susceptibility
Volunteer oilseed rape	0.75	From 3 leaves unfolded to second node detectable.	First flowers visible 3-30 cm height	S
Shepherd's purse	0.5	From 3 leaves unfolded to second node detectable.	First flowers visible 4-35 cm height	S
Creeping thistle	1.0	From 3 leaves unfolded to second node detectable.	40 cm height	R
Common fumitory	0.5	From 3 leaves unfolded to second node detectable.	Full flowering 2-22 cm height	S
Common hemp-nettle	1.0	From 3 leaves unfolded to second node detectable.	3-10 cm height	MS
Cleavers	0.75	From 3 leaves unfolded to second node detectable.	First flowers open 11-40 cm height	S

Weed species	Recommended rate L/ha	Timing crop	Maximum growth stage controlled	Susceptibility
Small-flowered cranesbill	0.5	From 3 leaves unfolded to second node detectable.	12-25 cm diameter	S
Red dead-nettle	0.5	From 3 leaves unfolded to second node detectable.	Full flowering 5-18 cm height	S
Scented mayweed, Scentless mayweed	1.0	From 3 leaves unfolded to second node detectable.	2-12 cm height	S
Common poppy	1.0	From 3 leaves unfolded to second node detectable.	10 cm diameter	S
Common chickweed	1.0	From 3 leaves unfolded to second node detectable.	Full flowering 2-25 cm height	S

Application rates and timing for spring applications (15th February to 30th June) to spring cereals

Weed species	Recommended rate L/ha	Timing crop	Maximum growth stage controlled	Susceptibility
Fat-hen	1.0	From 3 leaves unfolded to second node detectable.	3-15cm height	S
Cleavers	0.75	From 3 leaves unfolded to second node detectable.	4-20 cm height	S
Common fumitory	0.5	From 3 leaves unfolded to second node detectable.	6 true leaf stage 4-7 cm height	S
Dead-nettles	0.5	From 3 leaves unfolded to second node detectable.	8 true leaf stage 2-10 cm height	S
Scented mayweed, Scentless mayweed	1.0	From 3 leaves unfolded to second node detectable.	4-20 cm height	S

Weed species	Recommended rate L/ha	Timing crop	Maximum growth stage controlled	Susceptibility
Black bindweed	1.0	From 3 leaves unfolded to second node detectable.	Flower buds visible 3-24 cm height	S
Common chickweed	0.75	From 3 leaves unfolded to second node detectable.	1-15cm diameter	S

Rescue treatment for cleavers to be applied to the crop from third node detectable to flag leaf sheath swollen (winter and spring cereals) and no later than the 30th of June.

Weed species	Recommended rate L pr/ha	Timing crop	Maximum growth stage controlled	Susceptibility
Cleavers	1.0	From third node detectable to flag leaf sheath swollen	Beginning of flowering Up to 90 cm	S

For best results apply ZYPAR to young weeds which will respond quicker when treated at early stages. Vigorous crop competition enhances control of the difficult to control weeds.

GRASSLAND (ESTABLISHED, NEWLY SOWN & SEED PRODUCTION)

ZYPAR can be applied to grassland (established, newly sown, for seed production) from 3 true leaves of the grass (GS 13).

Application timing and Rate of use

For applications from 15th February to 31st August, ZYPAR can be applied at a dose rate of 1 L/ha.

For applications from 1st September to 14th February, do not apply more than 0.75 L/ha.

Application timing is crucial for good control. Applications must be made to weeds that are actively growing at the time of application.

Weeds controlled

Application rates and timing for applications (1st September to 14th February)

Weed species	Recommended rate L/ha	Maximum growth stage controlled	Susceptibility
Volunteer oilseed rape	0.75	7 true leaf stage 2-10 cm diameter	S
Cleavers	0.75	2 true leaf stage 2-6 cm diameter	S
Red dead-nettle	0.5	2 true leaf stage 2-3 cm diameter	S

Weed species	Recommended rate L/ha	Maximum growth stage controlled	Susceptibility
Scented mayweed, Scentless mayweed	0.75	4 side shoots visible 1-25 cm diameter	S
Common poppy	0.75	6 side shoots visible 5 cm diameter	S
Common chickweed	0.75	3 side shoots visible 1-10 cm diameter	S
Ivy-leaved speedwell	0.75	2 true leaf stage 1-2 cm diameter	S

Application rates and timing for applications (15th February to 31st August)

Weed species	Recommended rate L/ha	Maximum growth stage controlled	Susceptibility
Volunteer oilseed rape	0.75	First flowers visible 3-30 cm height	S
Shepherd's purse	0.5	First flowers visible 4-10 cm height	S
Common fumitory	0.5	Full flowering 2-22 cm height	S
Common hemp-nettle	1.0	3-10 cm height	MS
Cleavers	0.75	First flowers open 11-40 cm height	S
Small-flowered cranesbill, cut-leaved cranesbill	0.5	12-25 cm diameter	S
Red dead-nettle	0.5	Full flowering 5-15 cm height	S
Scented mayweed, Scentless mayweed	1.0	4-20cm height	S
Common poppy	1.0	10 cm diameter	S
Common chickweed	1.0	Full flowering 2-25 cm height	S
Fat-hen	1.0	3-15 cm height	S
Black-bindweed	1.0	Flower buds visible 3-24 cm height	S

For best results apply ZYPAR to young weeds which will respond quicker when treated at early stages. Vigorous crop competition enhances control of the difficult to control weeds.

Cutting Interval

Treated grassland can be cut after 14 days following application.

Grazing Interval

Livestock must be kept out of treated areas for at least 14 days following treatment.

Rolling / Harrowing Interval

DO NOT roll or harrow grass for 10 days before or 7 days after application.

WEATHER CONDITIONS

ZYPAR can be used in cold or warm (from 2 to 25 °C), humid or dry conditions. In severe drought conditions there can be a slight reduction in efficacy.

RAINFASTNESS

ZYPAR is rainfast one hour after application.

WATER VOLUME

Using standard or low drift nozzles the recommended spray volume is 100 to 400 litres of water per hectare. The lowest water volume should only be used in open crops on small weeds. On later applications where the crop is dense, the spray volume should be increased to 200-400 litres water per hectare. The minimum recommended pressure is 2 to 3 bars.

For undersown cereal crops, it is recommended to use a minimum spray volume of 150 litres of water per hectare.

For grassland, it is recommended to use a spray volume of 200-400 litres of water per hectare.

Use higher water volumes where the crop is dense.

CROP FAILURE

In the event of a crop failure after a crop has been treated with ZYPAR, consider all herbicides used on the treated field before deciding which new crop to sow.

Autumn applications

In case of a crop failure after an autumn application of ZYPAR at 0.75 L/ha, it is possible to sow the following spring crops, with no cultivation restrictions:

- 1 month after application (no cultivation restrictions): Spring wheat, Spring barley, Maize, Ryegrass.
- 3 months after application (after ploughing): Spring Oilseed rape, Field beans, Peas, Sunflower.

Spring applications

In case of a crop failure after a spring application of ZYPAR at 1.0 L/ha, it is possible to sow:

- 1 month after application (no cultivation restrictions): Spring wheat and Spring barley.
- 2 months after application (after ploughing): Maize.

FOLLOWING CROPS

After an application of ZYPAR there are no restrictions for sowing any succeeding crop after the cereal harvest. However, for sensitive species such as soybean, clover, lentils or sunflower ploughing is recommended prior to drilling.

ADJACENT CROPS

ZYPAR is of low volatility and is therefore not subject to vapour drift.

Avoid spray drift onto non-target crops.

Particular care should be taken to avoid spray drift onto susceptible crops e.g. vines, orchards, sunflower, oil seed rape, legumes, vegetable crops, ornamentals, flax, sugar beet, potatoes etc.

Do not spray in windy weather or only if the wind speed is less than or equal to 3 on the Beaufort scale (i.e. a maximum of 19 km/hour) and a temperature not exceeding 25 °C in the shade.

Apply only with a boom sprayer and use suitable nozzles for herbicides. Use a spray pressure as low as possible and a low height with respect to the treated vegetation, while ensuring the good spray distribution.

The use of drift reduction technology is recommended.

MIXING AND SPRAYING

Application equipment

ZYPAR should be applied through a tractor mounted or trailed horizontal sprayer.

Mixing

Half fill the spray tank with water and add the required amount of ZYPAR. Fill up the spray tank, agitating continuously to ensure thorough mixing, and maintain agitation until spraying is complete. Use only clean water for mixing.

Spray quality

Apply ZYPAR as a MEDIUM / COARSE spray as defined by the BCPC system.

TANK MIXTURES

Where tank mixes are used, and unless directed otherwise, the preferred order of addition of products to the spray tank is as follows: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, solution concentrates. Each product should be added to the half-full sprayer and be fully dispersed before the addition of the next product.

TANK CLEANING

To avoid subsequent injury to crops other than cereals (wheat, durum wheat, spelt, barley, rye, triticale), all spraying equipment must be thoroughly cleaned both inside and out using a proprietary tank cleaner, such as, All Clear Extra spray cleaner at 0.5 % v/v as follows:

1. Immediately after spraying thoroughly and completely rinse all internal surfaces.
2. Rinse inside of tank with clean water and flush through booms and in-line strainers using at least one tenth of the spray tank volume. Drain tank completely.
3. Fill tank with clean water (minimum 10% of tank capacity) and add tank cleaner at the recommended rate and agitate for 15 minutes. Flush the boom and hoses and drain tank completely.

4. Nozzles and filters should be removed and cleaned separately with a tank cleaner at the recommended rate.
5. Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume. Drain tank completely.
6. For disposal of washings, follow Code of Practice for Using Plant Protection Products. Do not spray onto a sensitive crop or land intended for cropping with a sensitive crop.
 1. **Note:** If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

COMPATIBILITIES

For details on compatibilities contact your distributor or Corteva hotline on +44 (0) 800 689 8899 or ukhotline@corteva.com

TRADEMARK ACKNOWLEDGEMENTS

ZYPAR is a trademark of Corteva Agriscience and its affiliated companies. All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

CORTEVA AGRISCIENCE CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<https://www.corteva.ie/content/dam/dpagco/corteva/eu/e/en/files/sds/ZYPAR-IRL-SDS.pdf>

Alternatively contact your supplier