

PROTECT FROM FROST  
SHAKE WELL BEFORE USE

WINGER provides moderate control of powdery mildew on winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.

750ml

app.rox. measure

CONDITIONS FOR USE  
FOR USE AS AN AGRICULTURAL FUNGICIDE

CROP	Max single dose (litres/hectare/crop)	Max no of applications	Max total dose (litres/hectare/crop)	Latest timing of application
Winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.	0.5l/ha	-	1.0	Before beginning of flowering: first anthers visible (GS 61)

500ml

#### Additional Safety Information

##### Operator protection

Wash splashes from skin and eyes immediately.  
Wash hands and exposed skin before meals and work.  
Wear suitable protective clothing, gloves and eye/face protection.  
Harmful to livestock. Keep livestock out of treated areas for at least 7 days.  
In case of contact with eyes, rinse immediately with plenty of water and seek medical advice.  
Do not breathe spray.

##### Storage and disposal

Keep in original container, tightly closed in a safe place.  
Wash out containers thoroughly, preferably using an integrated pressure rinsing device, or manually rinse three times. Add washings to the sprayer at the time of filling. Complete filling to the required volume and continue to agitate throughout the spraying operation.

##### Restrictions

Do not apply more than 3 times to any one crop.  
To protect aquatic organisms, respect an unsprayed buffer zone of 15m to surface water bodies.  
WINGER should only be used up to the full ear emergence (GS 59) stage on all crops.  
Disease control may be reduced if strains of the pathogen less sensitive to WINGER develop.

Lxxxxx PPE xxxxxxx

250ml

# WINGER

WINGER is an emulsifiable concentrate containing 750 grams fenpropidin (81.8% w/w) per litre.

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

WINGER provides moderate control of powdery mildew on winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.

#### FOR PROFESSIONAL USE ONLY

To avoid risks to human health and the environment comply with the instructions for use.

WINGER<sup>®</sup> is an emulsifiable concentrate containing 750 g/l (81.8% w/w) fenpropidin. Also contains hydrocarbons, C10, aromatics, < 1% naphthalene and 2-methylpropan-1-ol.

#### Warning

Harmful if swallowed or if inhaled.  
Causes serious eye irritation.  
May cause respiratory irritation.  
May cause drowsiness or dizziness.  
Suspected of damaging the unborn child.  
May cause damage to organs (Lungs, Nervous system, Eyes) through prolonged or repeated exposure.  
Very toxic to aquatic life with long lasting effects.  
Obtain special instructions before use.

Do not breathe the mist or vapours.  
Wash skin thoroughly after handling.  
Wear Protective gloves/ protective clothing/ eye protection.  
IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTRE/doctor if you feel unwell.  
Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty triple rinsed clean containers which can be disposed of as non-hazardous waste.  
Contains fenpropidin (ISO). May produce an allergic reaction.

PCS No. 05807 UFI: 9YW4-80NW-N00Q-UK8K



Authorisation Holder	Marketing Company	
DHM Agrochemicals Ltd Castle Lodge Kilgobbin Road Dublin 18 Tel: 01 2952377	DHM Agrochemicals Ltd Castle Lodge Kilgobbin Road Dublin 18 Tel: 01 2952377	Net contents: 1 litre Product code / Print date / Batch Number

GROUP 5 FUNGICIDE

#### DIRECTIONS FOR USE

IMPORTANT: this information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

#### CROPS

WINGER may be used on all varieties of winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.

#### DISEASE CONTROL

For moderate control of powdery mildew apply during the early stages of disease development.

Best results will be achieved when not more than 5% of the third leaf from the top of the plant is infected. A repeat spray may be necessary during prolonged or severe attacks.

#### APPLICATION RATE

Apply WINGER at 0.5 litres per hectare.

#### MIXING AND SPRAYING

Make sure the sprayer is set to give an even application at the correct volume.  
Fill the sprayer tank with half the required quantity of clean water and begin agitation. Add the required quantity of WINGER and continue agitation whilst adding the rest of the water.  
Agitate the mixture thoroughly before use and continue agitation during spraying.  
Thoroughly wash all spray equipment with water immediately after use.

#### APPLICATION

##### Volume

Apply WINGER in a recommended 200 litres of water per hectare.

##### COMPATIBILITY

The following products may be tank-mixed with WINGER provided the timing is correct for both WINGER and the partner in the mixture.  
The products should be added separately to the bulk of the water in the spray tank.  
Continuous agitation should be maintained and the products used immediately after mixing.

##### Herbicides

Approved salt formulations of:  
MCPA  
Mecoprop

##### Growth Regulators

Approved formulations containing chlormequat or chlormequat Chloride.

For further information or compatibility, contact DHM Agrochemicals Ltd.

ULBDHWING1L (03-17)

Lxxxxx PPE xxxxxxx