

# TRAILER<sup>®</sup>

## HERBICIDE



# TRAILER<sup>®</sup>

## HERBICIDE

Post-emergence herbicide  
for control of broadleaf  
weeds in cereals

Group 2 HERBICIDE

A water dispersible granules (WG) formulation, containing 750 g/kg of tribenuron-methyl for control of broad-leaved weeds in cereals (winter and spring wheat, winter and spring barley, oats, winter rye, triticale and setaside).

### SAFETY INFORMATION

#### WARNING

May cause damage to organs through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects.

Contains "tribenuron methyl". May produce an allergic reaction.

Do not breathe dust.

Avoid to release to the environment.

Wear protective gloves/ protective clothing/ eye protection/ face protection.

IF ON SKIN: wash with plenty of water.

Collect spillage.

Dispose of contents/container to licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers, which can be disposed of as non-hazardous waste.

Safety data sheet available on request.

Use appropriate containment to avoid environmental contamination.

To avoid risks to human health and the environment, comply with the instructions for use.

FOR PROFESSIONAL USE ONLY.

PCS No. 05858

**Jebagro** Hansoatic Trade Center,  
Kehlweider 11  
20457 Hamburg, Germany



CONTENT:  
**100 g**

Holder: **INDUSTRIAS AFRASA, S.A.** Ciudad de Sevilla, 53. Pol. Ind. fuente del Jarro  
46988 - Paterna (Valencia) SPAIN - Tel. +34 96 132 17 00 - [www.afrasa.es](http://www.afrasa.es)

### IMPORTANT INFORMATION

#### FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

**Crops:** Winter/spring wheat,  
winter/spring barley, oats, winter  
rye, triticale and setaside

**Maximum single dose:** 20 g product per hectare

**Maximum number of treatments:** One per crop for winter/spring  
wheat, winter/spring barley, oats,  
winter rye, triticale

**Latest Time of application:** Before flag leaf sheath extending  
stage (BBCH 39)

**READ ALL OTHER SAFETY PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USE**

**USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE USING PLANT PROTECTION PRODUCTS**

### SAFETY PRECAUTIONS

Precautions marked \* are a legal requirement.

#### Operator Protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

**WEAR SUITABLE PROTECTIVE GLOVES** when handling the concentrate or handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

**WASH ALL PROTECTIVE CLOTHING** thoroughly after use, especially the inside of gloves.

**WASH SPLASHES** from skin immediately.

**WASH HANDS AND EXPOSED SKIN** before meals and at work.

**IF YOU FEEL UNWELL**, seek medical advice (show the label where possible).

#### Environmental Protection:

Do not contaminate surface water with the product or its container. Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads.

Protect aquatic organisms respect an unsplashed buffer zone of 5m to surface water bodies.

#### Storage and Disposal:

KEEP AWAY FROM FOOD, FEED AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

Emergency Information: 24 hour Freephone: +353 01 809 2566

UFI: **D44-C0G5-N00Y-J6UM**

### GENERAL NOTES

Consult label tank cleaning procedures for all tank mix partners and ensure to use the most rigorous procedure recommended.

### NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

Batch number: See packaging

Manufacture date: See packaging

Expiry date: See packaging

### DIRECTIONS FOR USE

**IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.**

#### Weed Control

TRAILER is a foliar acting weed killer, to be applied in post-emergence of crop, for the control of broad-leaved weeds on cereals which can be used on all varieties of wheat, barley, oats, rye and triticale. It can also be used for set-aside. Applications should be made to young actively growing weeds. Activity is faster under good growing conditions and may appear relatively slow under cold or dry weather conditions.

#### Restrictions

Do not spray crops under stress or crops suffering from waterlogging, pest attack, disease or frost.

Do not spray crops undersown with grass mixtures.

Do not apply more than once to any cereal crop.

Do not apply within seven days of rolling the crop.

Special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleaning of equipment is also very important.

#### Weed Resistance

There is a significant risk for the development of weed resistance to TRAILER. Development of resistance can be avoided by alternating with suitable products having a different mode of action. Use also crop rotation and other cultural control measures to prevent and manage herbicide resistance.

#### Susceptible Weeds

The susceptibility rating of weeds in the following table refer to good spray cover and young, actively growing weeds at the timing of treatment.

Weed Species	Common name	Scientific name	Level of susceptibility
Corn chamomille		<i>Anthemis arvensis</i>	S
Fat hen		<i>Chenopodium album</i>	S
Creeping thistle		<i>Cirsium arvense</i>	S
Henbit		<i>Lamium amplexicaule</i>	S
Common poppy		<i>Papaver rhoeas</i>	S
Sow-thistle		<i>Sonchus olearaceus</i>	S
Common chickweed		<i>Stellaria media</i>	S
Saclet pimpernel		<i>Anagallis arvensis</i>	MS
Dead-nettle		<i>Lamium purpureum</i>	MS
Common fumitory		<i>Fumaria officinales</i>	R
Speedwell		<i>Veronica sp.</i>	R

S = Susceptible MS= Moderately Susceptible R = Resistant

#### Soil and Weather

TRAILER can be used on all soil types. Weed control may be reduced in dry conditions.

#### Volume and application

BEFORE USING TRAILER, SPRAYING EQUIPMENT MUST BE CLEAN AND FREE FROM CONTAMINATION WITH OTHER PESTICIDES.

Spray TRAILER in 200-400 litres of water per hectare. Where there are dense weed populations or advanced growth stages the water volume should be 400 litres per hectare.

Use a conventional field crop sprayer at a pressure of 2 - 3 bars and apply as a MEDIUM spray (as defined by BCPC). Care should be taken not to overlap spray swaths.

#### Mixing

Quarter fill the spray tank with clean water, start the agitation and add the required quantity of TRAILER directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

#### Compatibility

TRAILER should be mixed with other herbicides to widen and improve performance for the control of weeds. For information on tank-mixing partners contact your distributor.

#### CROPS

TRAILER can be used on all varieties of the following cereals: wheat (winter & spring), barley (winter & spring), oats, rye and triticale. TRAILER can be used also for setaside.

#### Timing

TRAILER must only be applied one per crop, from the crop without tillers, until before the flag leaf sheath extending stage (from BBCH 20 to BBCH 39).

Not to be applied before end of February in the year of harvest.

#### Dose

Apply TRAILER at a maximum rate 20 g per hectare.

#### Following Crops

After harvest of a cereal crop treated with TRAILER, only cereals, oilseed rape, field beans or grass may be sown in the same calendar year. In the event of crop failure for any reason, only a cereal crop can be sown after an interval of 3 months.

#### WARNINGS

**Take extreme caution to avoid damage by drift onto broad-leaved plants outside the target area or onto surface waters or ditches or land intended for cropping.**

Spraying equipment should not be drained or flushed onto land planted with or intended for planting with trees or crops other than cereals.

#### SPRAY TANK CLEAN-OUT

Poor cleanout practices and insufficient water volumes used for the rinse procedure may result in inadequate removal of product deposits. Subsequent use of application equipment in these circumstances may result in damage to non-cereal crops.

To avoid subsequent damage to crops other than cereals, immediately after spraying trailer thoroughly clear all spray equipment, including inside and outside of lid, using a proprietary sprayer cleaner for use with sulfonylureas according to the following procedure:

1. Immediately after spraying, drain tank completely. Wash any contamination off the outside of the sprayer with clean water.
2. Rinse the inside of the tank with clean water and flush at least one tenth of the spray tank volume through the boom and hoses. Drain tank completely.
3. Half-fill the tank with clean water and add the correct quantity of a proprietary sprayer cleaner, like a liquid alkali detergent (eg Drive\*, Surf\*, OMO\*) at 0.5 L/100 L. Agitate and then flush the boom and hoses with the solution. Top up completely with water and allow to stand for 15 minutes with agitation. Again flush the booms and hoses and drain tank completely (if it is not possible to drain the tank completely, repeat step 3 before going on).
4. Remove nozzles and filters and soak in a bucket containing a proprietary sprayer cleaner at the same concentration as that used for the sprayer.
5. Rinse tank again with clean water and flush at least one tenth of the tank volume through the booms and hoses. Drain tank completely.

CJ-231a-IE-Q05-R00

CJ-231a-IE-Q05-R00