

Enervin® SC

A fungicide for use in potatoes and leeks.

A suspension concentrate containing 200 g/l ametoctradin, 1,2-benzisothiazol-3(2H)-one and 2-methylisothiazol-3(2H)-one

Risk and Safety Information

Warning

May cause an allergic skin reaction.
Toxic to aquatic life with long lasting effects.

If medical advice is needed, have product container or label at hand.

Wear protective gloves/clothing/eye protection.

Avoid breathing mist/spray.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation occurs: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazol-3(2H)-one; 2-methylisothiazol-3(2H)-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCSC No. : 05995



UFI: 1C58-03D2-C00S-NQXT

Supplied by:

BASF Ireland DAC, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG, UK Telephone: 00 44 (0)161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112

Technical Enquiries: 0044 (0)845 602 2553 (office hours)

Authorisation holder:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG, UK Telephone: 00 44 (0)161 475 3000

5 L

® = Registered trademark of BASF

81173638 IE 2014



FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

Crop	Maximum Individual dose	Maximum number of applications	Maximum total dose	Latest time of application
Potato (seed & ware)	1.2 litres product per hectare	4	4.8 litres product per hectare	7 days before harvest
Leek	1.2 litres product per hectare	2	2.4 litres product per hectare	7 days before harvest

Other specific restrictions:

Do not apply by hand held equipment.

Minimum interval between applications is 5 days.

Under high disease pressure, tank mixing of Enervin SC with a contact protectant fungicide should be practiced.

READ ALL PRECAUTIONS BEFORE USE.

PRECAUTIONS

Operator protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

TAKE OFF IMMEDIATELY all contaminated clothing.

WASH SPLASHES from skin immediately.

WHEN USING DO NOT EAT, DRINK OR SMOKE

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible)

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

STORAGE

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

DO NOT RE-USE CONTAINER for any other purpose.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Enervin SC is a protectant fungicide for use in the early stages of foliar late blight (*Phytophthora infestans*) in potato and white tip (*Phytophthora porri*) in leek.

Restrictions/Warnings

For professional use only.

For application using by tractor mounted/trailed sprayers only.

Disease control

Enervin SC will provide protection against the early stages of foliar late blight in potato and white tip in leek. A full programme of correctly timed sprays using other authorised fungicides is likely to be required for longer term control and particularly where disease pressure is high.

Disease Susceptibility

Enervin SC is most effective when applied as a protectant spray before the risk of late blight in potato or white tip in leek occurs.

Enervin SC will give effective protection against phenylamide resistant strains of late blight.

Resistance Management

GROUP	45	FUNGICIDE
--------------	-----------	------------------

Enervin SC contains ametoctradin (a triazolo-pyrimidylamine, QoSI fungicide).

To minimize the risk of resistance developing, and in line with current FRAG-UK advice to use effective doses, always use Enervin SC at the recommended rate (1.2 l/ha) and in mixture with products with a different mode of action at a dose that will give robust control.

In potato no more than 3 consecutive applications of Enervin SC, and no more than a total of 4 applications of Enervin SC per season should be applied.

In leek no more than a total of 2 applications of Enervin SC per season should be applied.

No more than 6 applications of a QoSI fungicide, making up no more than half of the total number of intended late blight sprays per season, may be made.

Enervin SC should be used early in the spray programme, after which application with other fungicides should be made that can offer sufficient protection under high disease

conditions. In this way sub-optimal control of *Phytophthora infestans* is prevented, thereby managing the resistance risk.

General advice:

In potato, Enervin SC will control strains of late blight resistant to phenylamides. No cross resistance to phenylamide resistant strains of blight have been observed.

If an intensive blight control programme is necessary, include products with different modes of action.

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programmes should include fungicides with differing modes of action.

For further advice on resistance management for potato blight (*Phytophthora infestans*) contact your agronomist or specialist advisor and/or visit the FRAG-UK and Potato Council websites.

Crop Specific Information

Potatoes

Enervin SC can be used on all varieties of potato, including seed crops.

Time of application

Enervin SC should be applied as part of a repeat spray programme at 5 – 10 day intervals, depending on disease pressure. As disease pressure and the risk of late blight infection increases, the interval should be shortened. Irrigated crops should be treated as high risk.

Applications of Enervin SC should begin as soon as there is a risk of blight infection, or an official blight warning has been issued. In the absence of any warnings, applications should commence before the crop meets along the rows.

Rate of application

1.2 L/ha Enervin SC in 200-400 L/ha of water.

Leek

Qualified recommendation for a minor use:

There is limited evidence of crop safety and effectiveness for the control of *Phytophthora porri* in leek.

Time of application

Enervin SC should be applied as part of a repeat spray programme at 7 – 10 day intervals, depending on disease pressure. As disease pressure and the risk of white tip infection increases, the interval should be shortened. Irrigated crops should be treated as high risk.

Applications of Enervin SC should begin as soon as there is a risk of white tip.

Rate of application

1.2 L/ha Enervin SC in 350-1000 L/ha water.

Mixing and Application

Never prepare more spray solution than is required.

Three quarters fill the tank with clean water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of Enervin SC to the spray tank while re-circulating. Add the remainder of the water and continue agitation until spraying is complete

When tank mixes are to be used, take due note of any instructions given as to the order of mixing. Each product should be added separately to the spray tank and fully dispersed before the addition of any further product(s).

All tank mixes should be used immediately after mixing.

On emptying the container, rinse container thoroughly by using an integrated pressure rinsing device or by manually rinsing three times. Add washings to tank at time of filling and dispose of container safely

Apply as a medium spray, as defined by BCPC.

Apply in a water volume of 200 – 400 l/ha for potato and 350 – 1000 l/ha for leek, selecting the appropriate water volume for the density of the crop, ensuring good coverage of the foliage and stems.

Tank cleaning

When using tank mixtures always follow the most rigorous cleaning procedure.

Wash sprayer thoroughly immediately after use, using clean water and follow the sprayer cleaning guidance provided by the equipment manufacturer.

For information on compatibilities, contact your distributor, local BASF representative or the BASF Technical Services Hotline 0044 (0) 845 602 2553.

The following does not form part of the product label under S.I. No. 159 of 2012

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of any applicable law.

Additional Product Safety Information

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.basf.ie/Enervin-SC-IE/MSDS

Alternatively, contact your supplier.